

# Antidepressant-like effects of Chaihu-Shugan-San via SAPK/JNK signal transduction in rat models of depression

Yun-Hui Li, Chun-Hu Zhang, Juan Qiu, Su-E Wang, Sui-Yu Hu, Xi Huang, Ying Xie, Yang Wang, Tian-Li Cheng

Institute of Integrated Traditional Chinese and Western Medicine, Key Laboratory of Traditional Chinese Medicine Gan Organ of SATCM, Xiangya Hospital, Central South University, 87 Xiangya Road, 410008 Changsha, PR China

Submitted: 27-06-2013

Revised: 17-08-2013

Published: 24-07-2014

## ABSTRACT

**Background:** Chaihu-Shugan-San (CHSGS), a traditional Chinese medicinal herbal formula, registered in Jingyue Quanshu, has been indicated that oral administration of the extract from it can remit depressive disorder. C-Jun amino-terminal kinase (JNK/SAPK) signal transduction plays a key role in the apoptosis of nerve cells, be reported closely correlated with depression. This study was designed to investigate CHSGS antidepressant-like effects in rat models of depression and probe its possible mechanism. **Materials and Methods:** The classical experimental depression model chronic mild unpredictable stress (CMUS) was used to evaluate the antidepressant-like effects of CHSGS. The extracts were administered orally for 14 days, while the parallel positive control was given at the same time using fluoxetine hydrochloride. The expressions of JNK in the hippocampus were detected by real-time fluorescent quantitation PCR and Western blot assay. **Results:** Intragastric administration of CHSGS for 14 days caused a significant improvement of weight and locomotor activity in the open-field test. In addition, CHSGS treatment inhibited the expressions of JNK in the hippocampus tissue in CMUS rats. **Conclusion:** CHSGS could obviously improve the depressive state of the model rats and its mechanism may be correlated with regulating the expressions of JNK in the hippocampus.

**Key words:** Antidepressant, C-Jun amino-terminal kinase, Chaihu-Shugan-San, fluoxetine, open field test, signal transduction

## INTRODUCTION

Depression is a severe, common mood disorder with high suicide rate and the most disabling medical disease.<sup>[1]</sup> Because the mechanism of depression is quite complex and is not explicit, many currently available synthetic chemical antidepressants have low rates of response and remission and even severe adverse-effect.<sup>[2]</sup> Accordingly, it is necessary to probe and develop more effective antidepressant with lower adverse-effect. Nature plants, such as *Cissampelos sympodialis*,<sup>[3]</sup> *Bacopa monniera*,<sup>[4]</sup> *Terminalia Bellirica* Roxb,<sup>[5]</sup> *Hypericum perforatum*,<sup>[6,7]</sup> *Ginkgo biloba* 36, and *Pueraria lobata*,<sup>[8]</sup>

are an important source of new antidepressant drugs and the safety of nature plants may be better than that of synthetic antidepressants.<sup>[9]</sup> Chaihu-Shugan-San, a traditional Chinese medicinal (TCM) herbal formula, which consisted of seven Chinese herbs and recorded in a medical classic Jingyue Quanshu, has been used as a remedy for relieving depression symptoms caused by liver-Qi stagnation, which results from repression of distress and anger to TCM theory. Recently, it has been reported that CHSGS is effective in the treatment of depression. However, its mechanism is not clear.<sup>[10,11]</sup> It is reported that CHSGS may be antidepressant by regulating the MAPKs signal transduction, such as the expression of BDNF, TrkB, ERK, etc.<sup>[12,13]</sup> C-Jun amino-terminal kinase (JNK/SAPK), one of the signal transduction on MAPKs, plays a key role in the apoptosis of nerve cells and is closely correlated with depression.<sup>[14-16]</sup> This was designed to investigate CHSGS antidepressant-like effects in rat models of depression and examined whether they are correlated with JNK/SAPK signal transduction.

### Address for correspondence:

Dr. Chun-Hu Zhang, Institute of Integrated Traditional Chinese and Western Medicine, Key Laboratory of Traditional Chinese Medicine Gan Organ of SATCM, Xiangya Hospital, Central South University, 87# Xiangya Road, 410008 Changsha, PR China. E-mail: chzxyism@yahoo.com

### Access this article online

#### Website:

www.phcog.com

#### DOI:

10.4103/0973-1296.137367

#### Quick Response Code:



## MATERIALS AND METHODS

### Animals and groups

Forty healthy adult male Sprague-Dawley (SD) rats with similar ethological indexes, weighing 180-220 g, were provided by Experimental Animal Science of Xiangya Medical College of Central South University, and certificate of quality was SCXX (Hunan) 4343232. All rats were housed 5 per cage for 1 week to adapt to the environment under controlled conditions of 12 h light to 12 h dark cycle (lights on from 6:00 a.m. to 6:00 p.m.), background noise ( $40 \pm 10$ ) db, 10% relative humidity and temperature ( $20 \pm 3$ ) °C with food and water available *ad libitum*. Then, the rats were randomly divided into four groups: The normal control group (NC), the model control group (MC), the CHSGS group, and the fluoxetine control group (FC) group, with 10 rats in each group. Except those in the NC, the rest rats were singly housed and exposed on an unpredicted sequence of mild stressor. The experimental procedures were conducted in accordance with the Regulations for the Administration of Affairs Concerning Experimental Animals (1988) and approved by the Animal Experimental Center for Central South University.

### Drugs and reagents

The recipe and dosage of CHSGS used in this study followed the Chinese Pharmacopoeia, first volume 2005. Seven herbs (voucher specimen, no: 20100901-1) of CHSGS were obtained from Xiangya Hospital pharmacy, which were authenticated by the herbal medicine associate Professor Lei Peng, Department of Pharmacy of Xiangya Hospital affiliated Central South University. CHSGS contained Chinese thorowax root 9 g, dried tangerine peel 9 g, Szechwan Lovage Rhizome 9 g, Nutgrass Galingale Rhizome 9 g, Fructus Aurantii 9 g, Paeonia 15 g, and Radix Glycyrrhizae 5 g. CHSGS was boiled twice in distilled water (1:12, w/v) for 30 min. The blended supernatants were then lyophilized (yield = 19.13% (w/w)). The dried extract containing 8 g of crude drugs per gram were made into 1 g/mL of liquid when they were applied. Fluoxetine hydrochloride (FLU) was taken with a form of 20 mg/pellet, and qualified for health-tax credit number of J20080016 which was provided by Eli Lilly (Suzhou, China) Pharmaceutical Co., Ltd. The total RNA extraction reagent (TRIzol) was obtained from Invitrogen Corporation in America and SYBR Premix Ex Tap (perfect real time) kit was bought from Takara Co. Ltd.. The Revert Aid First Strand cDNA Synthesis kit was obtained from Promega (Madison, WI, USA) (A3500 29287) and primers were synthesized by Shanghai Bio-engineering Technology Co., Ltd. JNK (p-JNK) rabbit monoclonal antibody was purchased from Santa Cruz Biotech (Santa Cruz, CA, USA) (SC0564). Rabbit

IgG horseradish peroxidase (HRP)-conjugated rabbit anti-goat polyclonal antibody was purchased from KPL (Maryland, USA), and mouse IgG HRP-conjugated rabbit anti-mouse polyclonal antibody was purchased from Upstate (New York, USA). Radioimmunoprecipitation assay protein lysis buffer was obtained from Santa Cruz Biotechnology. Hybond-Polyvinylidene difluoride was obtained from Amersham (Uppsala, Sweden), and phenylmethyl sulfonyl fluoride was obtained from Sigma.

### Establishment of the CMUS model

All the rats except the normal group were used to establish chronic mild unpredictable stress (CMUS) depression rat model. The animal model was established following modified Willner's method,<sup>[17,18]</sup> which provide the rats CMUS and housed by single cage. The stress regimen consisted of the following stressors: 24 h reversed light/dark, electric shock in foot (10 mA electricity was given every other minute, which last 10 s per time for 30 times), 48 h food deprivation, shaking (once a second, lasting for 15 min), noise (85 db), clipping the tail (clipping last 5 s per time for 10 times), and strange smell. During a period of 28 days, one of the stressors was chosen randomly and done to the rats so that the rats could not expect the stimulus. Every stressor was used for two to three times in total.

### Drugs administration

Rats of the NC group were fed normally, and rats in other groups received different stimulus during the 28 days. From the 15<sup>th</sup> day, all rats were administered with equal volume of normal saline (to the NC group and the MC group) and of corresponding medicinal liquid (5.9 g/kg to the CHSGS group, and 1.8 mg/kg to the FC group) by gastrogavage for 2 successive weeks. The dosage of medicines administered to the rats equaled to that of a 70 kg adult.

### Weights and behavioral tests

#### Weight change

All the rats were weighed on the 0, 15<sup>th</sup>, and 29<sup>th</sup> day during the experiment.

#### Open-field test

The open-field test (OFT) was performed according to the improved method recorded in the literature<sup>[19]</sup> at 7:00 a.m. on the 0, 15<sup>th</sup>, and 29<sup>th</sup> day of the experiment in a quiet room. The procedure of the test was as follows: The rats were individually placed in the central wood cage (100 × 100 cm<sup>2</sup>) with walls 50 cm high and the floor divided into 10 squares (10 × 10 cm<sup>2</sup>). When the hind legs crossed the line of the squares, the rat was considered to have crossed from one square to another (crossing), when the forelegs lift from the floor the rat was considered to have gotten one point (scores of rears). The rats' scores during 3 min were recorded. After each rat was tested in

the OFT, the cage was carefully cleaned with a solution containing 75% ethanol, purified water to remove the scent of the previously evaluated rat, which could modify the spontaneous behavior of the rat.

#### **Sucrose-solution consumption test**

According to the literature,<sup>[20,21]</sup> the test procedures were as follows: In a quiet room 72 h before each OFT. Before the test, the rats were trained to adapt to drinking water with sugar in it. Two bottles were placed in every cage. In the first 24 h, the two bottles were filled with 1% sucrose solution, and in the following 24 h, one of the bottles was filled 1% sucrose solution, and the other was filled with pure water. Then, the basic energy expenditure test and water consumption test were done after 24 h of food and water forbidden. In the same time, the rats were given two bottles of liquid weighed beforehand, one was filled with 1% sucrose solution, and the other with pure water. Further 60 min later, the two bottles were weighed. Then, the total liquid consumption, sucrose solution consumption, and pure water consumption were calculated.

#### **Fluorescence real-time quantitative transcription-PCR analysis of JNK mRNA**

##### **Preparation of the rat brain sample**

Five rats from each group were killed by cutting down their heads on the 29<sup>th</sup> day of the experiment. Then, the brain tissues were immediately taken out on the ice table, and bilateral hippocampuses were isolated. They were placed into RNase-free EP tubes and kept first in liquid nitrogen for 24 h and then at -80°C in a freezer.

RNA was extracted with Trizol. Reverse transcription-polymerase chain reaction (RT-PCR) was accomplished with the Access RT-PCR System. JNK mRNA (211bp; GenBank AB118218.1) was amplified using the forward primer 5'-GGCGGCCAAACAGAAAG-3' and the reverse primer 5'-CTGAGGGCACGGAGGAT-3'. JNK  $\beta$ -actin (201 bp) mRNA was amplified as the control using the forward primer 5'-CGTTGACATCCGTAAAGA-3' and the reverse primer 5'-TGGAAGGTGGACAGTGAG-3'. Each reaction mixture (final volume 25  $\mu$ L) contained 4  $\mu$ L MgCl<sub>2</sub>, 2  $\mu$ L 10 $\times$  RT buffer, 2  $\mu$ L dNTP mixture, 0.5  $\mu$ L recombinant Rnasin, 0.7  $\mu$ L AMV, 1  $\mu$ L Oligo dT, 2  $\mu$ L RNA, and 12.8  $\mu$ L DEPC-treated water. The reaction parameters were: The reverse transcription was performed at 42°C for 15 min, followed by deactivation of reverse transcriptase at 95°C for 5 s.

The real-time quantitative PCR reaction was carried out with ABI 7300 Fluorescence quantitative PCR using reagents of the SYBR Premix Ex Taq (perfect real time) kit, in a reaction volume of 25  $\mu$ L, consisting of 10  $\mu$ L SYBR fluorescent dye, 2  $\mu$ L PCR forward primer and

PCR reverse primer, 2  $\mu$ L RT reactive fluid (cDNA), and 11  $\mu$ L DEPC-treated water. The reaction conditions were as follows: In the first denaturation step, the reaction was heated up to 95°C which was kept for 10 s, and followed by the PCR reaction step consisting of 40 circles, each circle composed of denaturation at the temperature of 95°C for 5 s and annealing at the temperature of 56°C for 30 s. In the whole process of the reaction, the threshold cycle ( $C_T$ ) value of the samples was analyzed with Sequence Detection software version 1.2.3 (Applied Bio-systems Corporation). Further, it was found that the  $C_T$  value decreased as the template concentration increased. Melting curve analysis was carried to assess the characteristics of the PCR. Further,  $\beta$ -actin served as the internal control, the  $2^{-\Delta\Delta C_T}$  method<sup>[11]</sup> was applied to do the relative quantitative analysis of JNK1/2 mRNA. The formula was as follows:  $\Delta\Delta C_T = \text{experimental group } (C_T \text{ assayed samples} - C_T \beta\text{-actin}) - \text{control group } (C_T \text{ assayed samples} - C_T \beta\text{-actin})$ .

##### **Protein extraction and Western blot analysis of JNK**

Radio immunoprecipitation assay protein lysis buffer was used to separate the brain tissue proteins, followed by quantification by the BCA Protein Assay kit (Pierce, Rockford, IL, USA). We analyzed the protein expression of JNK, phosphorylated at p-JNK, using a Bio-Rad electrophoresis apparatus and a protein transfer equipment (MiniProtean; Bio-Rad, Hercules, CA, USA) and Kodak XOMAT AR film (PerkinElmer Life and Analytical Sciences, Inc., Waltham, MA, USA). The antibodies used to probe the Western blot and their dilution was as follows: JNK 1:1000, p-JNK 1:1500.

##### **Statistical methods**

Statistical analyses were carried out using the SPSS 17.0 (SPSS Inc., Brookfield, WI, USA) software package. The results are expressed as the mean value  $\pm$  standard deviation. Data were analyzed two-way repeated measures analysis of variance (ANOVA), because the same rats were subjected to several test sessions. A two-tailed  $P < 0.05$  was considered to be statistical significance.

## **RESULTS**

### **Weight, OFT, and sucrose-solution consumption**

The weight and behaviors of all the rats before the experiment was no statistical significance ( $P > 0.05$ ). After the stress of 14 days chronic mild unpredictable stress, all the rats in the experiment groups had depressed behaviors. Compared with the rats in the NC group, their scores in the OFT decreased, the sucrose solution consumption reduced and the increasing of weight lowered, and the difference was of statistical significance ( $P < 0.05$  or  $< 0.01$ ). After

14 days of intragastric administration, the depression-like behaviors of the rats in the CHSGS group and the FC group had improved, scores of crossing and rears of the OFT increased obviously, the volume of the sucrose solution consumption increased, the increasing of their weight became evident, and the differences between them and the rats in the NC group were of statistical significance ( $P < 0.05$  or  $< 0.01$ ) [Table 1].

### Effects of CHSGS on mRNA of JNK

The molecular cloning of human JNK and rat JNK (SAPK) led to identification of JNK as a member of the MAP kinase group of protein kinases. The melting curves of PCR products of JNK and  $\beta$ -actin had only one peak, which indicated that the amplified gene was a single fragment. The melting temperature of JNK and  $\beta$ -actin was 85.3°C and 84.5°C, respectively. The  $C_T$  values of JNK in different groups and encephalic regions were detected by the amplification of  $\beta$ -actin. With  $\beta$ -actin as internal control, and the JNK mRNA expression of model was supposed to be 1, the relative JNK mRNA expression was calculated by  $2^{-\Delta\Delta C_T}$ . Further, the results are presented in Figure 1.

### Effects of CHSGS on protein expression of JNK and p-JNK

To confirm that the results obtained at the mRNA level corresponded at the protein level, Western blot analysis was carried out. JNK is activated when phosphorylated. To determine the effect of CHSGS on the activation status of JNK, we evaluated its phosphorylation state in the rat brain tissue treated with CHSGS. We detected the total JNK, including the 46-KDa and 54-KDa from the brain tissue of different groups, simultaneously, the 46-KDa p-JNK was

detected. Digital image analysis of the Western blot signals demonstrated significantly higher levels of p-JNK in the MC group relative to NC, and compared with the MC group, the expression of p-JNK was obviously lower in the CHSGS and FC group, the differences between them were of statistical significance ( $P < 0.05$  or  $P < 0.01$ ) [Figures 2 and 3].

## DISCUSSION

Currently, the chronic mild unpredictable stress depression model has become a widely used animal model for studying the pathogenesis of depression and mechanism of antidepressant's action.<sup>[17,22]</sup> In this study, after the establishment of the CMUS model, except those in the NC group, the rest rats underwent a drop in the increasing speed of weight, their scores in OFT and sucrose solution consumption reduced, showed that anhedonia, obviously compared to those in the NC group, loss of appetite, decrease of activity, etc., Those depression-like symptoms improved evidently after treated by the classic antidepressant fluoxetine, suggesting that the animal models were successfully established.<sup>[23]</sup> Meanwhile, after the treatment with administration of CHSGS for 14 days, the depressive-like symptoms were obviously improved.


Depression is one of the major mental disorders associated with symptoms such as regular negative moods, decreased physical activity, feelings of helplessness, sluggish thought, and cognitive function. It has been regarded as "depression syndrome" and "liver qi stagnation" in TCM.<sup>[24-26]</sup> CHSGS consists of seven crude herbal drugs and is used to treat depression-related syndromes associated with qi stagnation. Qi stagnation has been categorized as a term in TCM that

**Table 1: Comparison of weight, OFT, and sucrose-solution consumption among preduplicating, pretherapy rats ( $\pm$ s,  $n=10$ )**


| Groups | Scores of crossing      | Scores of rears        | Total water consumption/ml | 1% sucrose consumption/ml | Weight/g     |
|--------|-------------------------|------------------------|----------------------------|---------------------------|--------------|
| NC     |                         |                        |                            |                           |              |
| Day 0  | 65.6±16.3               | 15.9±6.2               | 16.3±4.1                   | 13.3±4.1                  | 203.6±10.8   |
| Day 15 | 64.5±14.5               | 16.1±5.3               | 17.8±3.7                   | 14.5±3.9                  | 271.1±16.5   |
| Day 29 | 61.5±15.7               | 15.5±4.9               | 20.6±5.4                   | 17.2±4.8                  | 324.5±23.7   |
| MC     |                         |                        |                            |                           |              |
| Day 0  | 66.4±8.2                | 15.7±5.3               | 16.6±4.4                   | 14.1±3.5                  | 204.5±10.3   |
| Day 15 | 27.5±8.9 <sup>Δ</sup>   | 6.8±2.8 <sup>Δ</sup>   | 15.9±3.6 <sup>Δ</sup>      | 6.6±2.3 <sup>Δ</sup>      | 249.8±13.2*  |
| Day 29 | 18.3±6.7                | 4.9±2.1                | 12.3±3.3                   | 7.1±2.8                   | 275.6±15.1   |
| CHSGS  |                         |                        |                            |                           |              |
| Day 0  | 65.8±12.7               | 16.3±4.8               | 17.1±4.9                   | 13.2±3.7                  | 206.9±12.6   |
| Day 15 | 26.4±11.3               | 4.7±2.6                | 16.3±4.5                   | 5.6±2.3                   | 252.6±13.4   |
| Day 29 | 49.6±15.4 <sup>ΔΔ</sup> | 12.3±4.5 <sup>ΔΔ</sup> | 19.1±3.8**                 | 15.3±3.9 <sup>ΔΔ</sup>    | 310.2±20.7** |
| FC     |                         |                        |                            |                           |              |
| Day 0  | 65.8±20.9               | 15.8±3.9               | 16.9±5.1                   | 13.9±3.5                  | 210.6±15.7   |
| Day 15 | 24.9±11.5               | 5.5±2.7                | 16.3±4.3                   | 6.3±3.1                   | 251.9±16.5   |
| Day 29 | 45.6±13.4 <sup>ΔΔ</sup> | 13.2±4.1 <sup>ΔΔ</sup> | 20.3±5.2 <sup>ΔΔ</sup>     | 16.2±5.4 <sup>ΔΔ</sup>    | 308.3±14.7** |

\* $P < 0.05$ ,  $^{\Delta}P < 0.01$ : Compared with the normal at the same time point; \*\* $P < 0.05$ ,  $^{\Delta\Delta}P < 0.01$ : Compared with model control group at the same time point; OFT: Open-field test






**Figure 1:** The expression of JNK mRNA in the hippocampus tissue in post-treatment of varied groups detected by fluorescence real-time quantitative transcription-PCR.  $\Delta P < 0.01$ , compared with the NC group;  $\Delta\Delta P < 0.01$ , compared with the MC group



**Figure 2:** The expression of c-Jun N-terminal kinase (JNK) and phosphorylated-JNK (p-JNK) in the rat hippocampus in posttreatment of varied groups detected by Western blot



**Figure 3:** The expression of p-JNK in the hippocampus tissue in posttreatment of varied groups detected by Western blot.  $\Delta P < 0.01$ , compared with the NC group;  $\Delta\Delta P < 0.01$ , compared with the MC group

refers to insidious, long-standing, serious discomfort that is projected into the body and is manifested by numerous

symptoms, such as insomnia, fatigue, panic, and dyspnea.<sup>[27]</sup> In previous studies, the antidepressant effect of CHSGS was revealed using the CMUS model and it was shown that the mechanism may be by regulating the neuroendocrine,<sup>[11]</sup> but its molecule mechanism is not very clearly.

The hippocampus is one brain structure that has been extensively studied with regard to stress, depression, and antidepressant actions.<sup>[28]</sup> It was reported that volumes of the double-side hippocampus were reduced in patients with major depression compared to healthy controls, and there was a positive correlation between hippocampus atrophy and the time course of the depression.<sup>[29-31]</sup> Chronic stress alone increased the numbers of apoptotic cells in hippocampal subregions and cortex in the adult tree shrew,<sup>[32]</sup> while treatment with the antidepressant tianeptine had an antiapoptotic effect in the stressed animals,<sup>[33]</sup> suggesting that chronic antidepressant treatment may have neuroprotective effect. Some previous studies found that classical antidepressants also protected rat hippocampal neurons in primary culture from the lesion induced by corticosterone.

JNK, initially described as stress-activated protein kinase (SAPK), was identified by their ability to phosphorylate specific sites on the amino terminal transactivation domain of the c-Jun, a transcription factor, following exposure to UV irradiation, growth factors, cytokines, or expression of transforming oncogenes. By phosphorylating these sites, the JNKs stimulate c-Jun's transcriptional activity.<sup>[34,35]</sup> The JNK pathway's major role, involvement in stress responses, has been widely studied. The JNK cascade follows the typical MAPK signaling arrangement. The major target of the JNK pathway is activator protein-1 (AP-1) transcription factor, which is activated mainly by the phosphorylation of c-Jun and other related molecules.<sup>[36]</sup> In the adult mouse, signaling through c-Jun N-terminal kinases (JNKs) links exposure to acute stress to various physiological responses. Inflammatory cytokines, brain injury and ischemic insult, or exposure to psychological acute stressors induce activation of hippocampal JNKs. Acute stress caused activation of JNKs in the hippocampal CA1 and CA3 subfields, and impaired contextual fear conditioning. Conversely, intrahippocampal injection of JNKs inhibitors sp600125 (30  $\mu\text{M}$ ) or D-JNKI1 (8  $\mu\text{M}$ ) reduced activity of hippocampal JNKs and rescued stress-induced deficits in contextual fear. In addition, intrahippocampal administration of anisomycin (100  $\mu\text{g}/\mu\text{L}$ ), a potent JNKs activator, mimicked memory-impairing effects of stress on contextual fear. This anisomycin-induced amnesia was abolished after cotreatment with JNKs selective inhibitor sp600125 without affecting anisomycin's ability to effectively inhibit protein synthesis as measured by c-Fos immunoreactivity. Studies

showed milder and transient activation of the JNKs pathway in the CA1 subfield of the hippocampus during contextual fear conditioning and an enhancement of contextual fear after pharmacological inhibition of JNKs under baseline conditions. Using combined biochemical and transgenic approaches with mutant mice lacking different members of the JNK family (Jnk1, Jnk2, and Jnk3), evidence suggested that JNK2 and JNK3 are critically involved in stress-induced deficit of contextual fear, while JNK1 mainly regulates baseline learning in this behavioral task. Currently studies support the possibility that hippocampal JNKs serve as a critical molecular regulator in the formation of contextual fear.<sup>[37]</sup>

In this study, our experiment results showed that the expressions of JNK mRNA and the protein of p-JNK in the rat of MC groups were increased significantly, and antidepressant Flu and CHSGS could regulate the expression of JNK. Therefore, our results showed that CHSGS could obviously improve the depressive state of the model rats and its mechanism may be correlated with regulating the expressions of JNK in the hippocampus.

## CONCLUSIONS

CHSGS could obviously improve the depressive behavior of the model rats and its mechanism may be correlated with regulating the expressions of JNK in the hippocampus.

## ACKNOWLEDGMENTS

This project was supported by the National Natural Science Foundation of China (No. 30801506) and was partly supported by fund for Key Unit of Traditional Chinese Medicine Gan of SATCM.

## REFERENCES

- Paykel ES. Depression: Major problem for public health. *Epidemiol Psychiatr Soc* 2006;15:4-10.
- Sarko J. Antidepressants, old and new. A review of their adverse effects and toxicity in overdose. *Emerg Med Clin North Am* 2000;18:637-54.
- Almeida RN, Navarro DS, de Assis TS, de Medeiros IA, Thomas G. Antidepressant effect of an ethanolic extract of the leaves of *Cissampelos sympodioides* in rats and mice. *J Ethnopharmacol* 1998;63:247-52.
- Sairam K, Dorababu M, Goel RK, Bhattacharya SK. Antidepressant activity of standardized extract of *Bacopa monniera* in experimental models of depression in rats. *Phytomedicine* 2002;9:207-11.
- Dhingra D, Valecha R. Evaluation of antidepressant-like activity of aqueous and ethanolic extracts of *Terminalia bellirica* Roxb. fruits in mice. *Indian J Exp Biol* 2007;45:610-6.
- Caccia S. Antidepressant-like components of *Hypericum perforatum* extracts: An overview of their pharmacokinetics and metabolism. *Curr Drug Metab* 2005;6:531-43.
- Sánchez-Mateo CC, Bonkanka CX, Prado B, Rabanal RM. Antidepressant activity of some *Hypericum reflexum* L. fil. extracts in the forced swimming test in mice. *J Ethnopharmacol* 2007;112:115-21.
- Yan B, Wang DY, Xing DM, Ding Y, Wang RF, Lei F, *et al.* The antidepressant effect of ethanol extract of *radix puerariae* in mice exposed to cerebral ischemia reperfusion. *Pharmacol Biochem Behav* 2004;78:319-25.
- Schulz V. Safety of St. John's Wort extract compared to synthetic antidepressants. *Phytomedicine* 2006;13:199-204.
- Kim SH, Han J, Seog DH, Chung JY, Kim N, Hong Park Y, *et al.* Antidepressant effect of Chaihu-Shugan-San extract and its constituents in rat models of depression. *Life Sci* 2005;76:1297-306.
- Li YH, Zhang CH, Wang SE, Qiu J, Hu SY, Xiao GL. Effects of Chaihu Shugan San on behavior and plasma levels of corticotropin releasing hormone and adrenocorticotrophic hormone of rats with chronic mild unpredictable stress depression. *Zhong Xi Yi Jie He Xue Bao* 2009;7:1073-7.
- Wang SE, Hu SY, Zhang CH, Qiu J, Li YH. Effect of Chaihu Shugan San and its components on expression of ERK1/2 mRNA in the hippocampus of rats with chronic mild unpredictable stress depression. *Zhong Na Da Xue Xue Bao Yi Xue Ban* 2011;36:93-100.
- Deng Y, Zhang CH, Zhang HN. Effects of chaihu shugan powder on the behavior and expressions of BDNF and TrkB in the hippocampus, amygdala, and the frontal lobe in rat model of depression. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 2011;31:1373-8.
- Sun Y, Yang T, Xu Z. The JNK pathway and neuronal migration. *J Genet Genomics* 2007;34:957-65.
- Brust TB, Cayabyab FS, MacVica BA. C-Jun N-terminal kinase regulates adenosine A1 receptor-mediated synaptic depression in the rat hippocampus. *Neuropharmacology* 2007;53:906-17.
- Li XM, Li CC, Yu SS, Chen JT, Sabapathy, Ruan DY. JNK1 contributes to metabotropic glutamate receptor-dependent long-term depression and short-term synaptic plasticity in the mice area hippocampal CA1. *Eur J Neurosci* 2007;25:391-6.
- Willner P, Towell A, Sampson D, Sophokous S, Muscat R. Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. *Psychopharmacology (Berl)* 1987;93:358-64.
- Grippo AJ, Beltz TG, Weiss RM, Johnson AK. The effects of chronic fluoxetine treatment on chronic mild stress-induced cardiovascular changes and anhedonia. *Biol Psychiatry* 2006;59:309-16.
- Yuan TT, Qiao H, Dong SP, An SC. Activation of hippocampal D1 dopamine receptor inhibits glutamate-mediated depression induced by chronic unpredictable mild stress in rats. *Sheng Li Xue Bao* 2011;63:333-41.
- Benelli A, Filaferrero M, Bertolini A, Genedani S. Influence of Sadenosyl-L-methionine on chronic mild stress-induced anhedonia in castrated rats. *Br J Pharmacol* 1999;127:645-54.
- Forbes NF, Stewart CA, Matthews K, Reid IC. Chronic mild stress and sucrose consumption: Validity as a model of depression. *Physiol Behav* 1996;60:1481-4.
- Isingrini E, Belzung C, Freslon JL, Machet MC, Camus V. Fluoxetine effect on aortic nitric oxide-dependent vasorelaxation in the unpredictable chronic mild stress model of depression in mice. *Psychosom Med* 2012;74:63-72.
- Willner P, Mitchell PJ. The validity of animal models of predisposition to depression. *Behav Pharmacol* 2002;13:169-88.
- Ma Q, Zhou DA, Wang LP. Clinical curative effect and factor analysis of depression treated by acupuncture. *Zhongguo Zhen Jiu* 2011;31:875-8.

25. Zhang HL. Treatment of depression based on differentiation of the Shaoyang Channels. *J Tradit Chin Med* 2010;30:83-92.
26. Han H, Wu LM, Wang MX, Tang JJ, Wang H, Liu RZ, *et al.* Characteristics of traditional Chinese medicine syndromes in post-stroke depression. *Zhong Xi Yi Jie He Xue Bao* 2010;8:427-31.
27. Wei XH, Cheng XM, Shen JS, Wang ZT. Antidepressant effect of Yueju-Wan ethanol extract and its fractions in mice models of despair. *J Ethnopharmacol* 2008;117:339-44.
28. Drevets WC, Price JL, Furey ML. Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Struct Funct* 2008;213:93-118.
29. Bremer JD, Vythilingam M, Nq CK, Vermetten E, Nazeer A, Oren DA, *et al.* Regional brain metabolic correlates of alpha-methylparatyrosine-induced depressive symptoms: Implications for the neural circuitry of depression. *JAMA* 2003;289:3125-34.
30. Sapolsky RM. Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. *Arch Gen Psychiatry* 2000;57:925-35.
31. Sapolsky RM. The possibility of neurotoxicity in the hippocampus in major depression: A primer on neuron death. *Biol Psychiatry* 2000;48:755-65.
32. Lucassen PJ, Fuchs E, Czéh B. Antidepressant treatment with tianeptine reduces apoptosis in the hippocampal dentate gyrus and temporal cortex. *Biol Psychiatry* 2004;55:789-96.
33. Lucassen PJ, Vollmann-Honsdorf GK, Gleisberg M, Czéh B, De Kloet ER, Fuchs E. Chronic psychosocial stress differentially affects apoptosis in hippocampal subregions and cortex of the adult tree shrew. *Eur J Neurosci* 2001;14:161-6.
34. Kyriakis JM, Banerjee P, Nikolakaki E, Dai T, Rubie EA, Ahmad MF, *et al.* The stress-activated protein kinase subfamily of c-Jun kinase. *Nature* 1994;369:156-60.
35. Ichijo H. From receptors to stress-activated MAP kinases. *Oncogene* 1999;18:6087-93.
36. Gupta S, Campbell D, Dériard B, Davis RJ. Transcription factor ATF2 regulation by the JNK signal transduction pathway. *Science* 1995;267:389-93.
37. Sherrin T, Blank T, Hippel C, Rayner M, Davis RJ, Todorovic C. Hippocampal c-Jun-N-terminal kinases serve as negative regulators of associative learning. *J Neurosci* 2010;30:13348-61.

**Cite this article as:** Li Y, Zhang C, Qiu J, Wang S, Hu S, Huang X, *et al.* Antidepressant-like effects of Chaihu-Shugan-San via SAPK/JNK signal transduction in rat models of depression. *Phcog Mag* 2014;10:271-7.

**Source of Support:** Nil, **Conflict of Interest:** None declared.