

# Profiling of components of rhizoma et radix polygoni cuspidati by high-performance liquid chromatography with ultraviolet diode-array detector and ion trap/time-of-flight mass spectrometric detection

Jinfeng Fu<sup>1</sup>, Min Wang<sup>1</sup>, Huimin Guo<sup>1</sup>, Yuan Tian<sup>1</sup>, Zunjian Zhang<sup>1</sup>, Rui Song<sup>1,2</sup>

<sup>1</sup>Key Laboratory of Drug Quality Control and Pharmacovigilance, Ministry of Education, China Pharmaceutical University, <sup>2</sup>Department of Pharmaceutical Analysis, China Pharmaceutical University, Tongjia Lane 24, Nanjing 210009, China

Submitted: 27-07-2014

Revised: 18-09-2014

Published: 10-07-2015

## ABSTRACT

**Introduction:** Rhizoma et Radix Polygoni Cuspidati (Huzhang in Chinese, HZ) is a traditional medicinal plant in China. Many of the components of HZ have been proved to be bioactive while it is difficult to conduct a comprehensive chemical profiling of HZ as a consequence of the absence of efficient separation system and sensitive detective means. We developed a simple and effective method for comprehensive characterization of constituents in HZ. **Objective:** To develop a simple and effective method to characterize the components in HZ and provide useful information for subsequent metabolic studies of HZ. **Materials and Methods:** The components in HZ aqueous extract were characterized by using high performance liquid chromatography with UV diode-array detector (HPLC-DAD) and ion trap/time-of-flight mass spectrometric detection (HPLC-IT/TOF). Stilbenes, anthraquinones, gallates and tannins, naphthalenes and some other compounds were identified and confirmed by diagnostic fragment ions with accurate mass measurements, characteristic fragmentation pathways and relevant published literatures. **Results:** Among the 238 constituents detected in HZ, a total number of 74 constituents were identified unambiguously or tentatively, including 29 compounds reported for the first time in HZ. **Conclusion:** The identification and structure elucidation of these chemicals provided essential data for quality control and further in vivo metabolic studies of HZ. **Key words:** *Polygonum cuspidatum*, HPLC-DAD, HPLC-IT/TOF, qualitative analysis.

**Key words:** *Polygonum cuspidatum*, high-performance liquid chromatography-diode array detector, high-performance liquid chromatography-ion trap/time-of-flight, qualitative analysis

## Access this article online

### Website:

www.phcog.com

### DOI:

10.4103/0973-1296.160455

### Quick Response Code:



## INTRODUCTION

Rhizoma et Radix Polygoni Cuspidati, well-known as Huzhang (HZ) in China and Japanese knotweed in Japan, comes from the stem and root of *Polygonum cuspidatum* Sieb. Et Zucc. This herb has been widely used in Chinese and Japanese folk medicine for the treatment of atherosclerosis, hypertension, cough, suppurative dermatitis, and gonorrhea. HZ contains a variety of

chemical groups such as stilbenes, anthraquinones, flavones, and tannins. Previous pharmacological and clinical studies have indicated that several chemical components in HZ are bioactive.<sup>[1-6]</sup>

Various methods have been used for qualitative analysis of major chemical constituents in HZ. Yi *et al.* developed a high-performance liquid chromatography-electrospray ionization/mass spectrometry (HPLC-ESI/MS) method for simultaneously qualitative and quantitative analysis of 9 compounds.<sup>[7]</sup> Xia *et al.* separated HZ methanol extract by HPLC-diode array detector (HPLC-DAD) and characterized 22 major constituents using HPLC-time-of-flight (TOF/MS).<sup>[8]</sup> Sun *et al.* detected 20 volatile oil in HZ by gas chromatography-MS.<sup>[9]</sup> Electrophoresis and NMR have also been adopted for qualitative assessment of HZ.<sup>[10,11]</sup>

### Address for correspondence:

Asso. Prof. Rui Song, Department of Pharmaceutical Analysis, China Pharmaceutical University, Tongjia Lane 24, Nanjing 210009, China.  
E-mail: songrui1123@gmail.com  
Prof. Zunjian Zhang, Key Laboratory of Drug Quality Control and Pharmacovigilance, Ministry of Education, China Pharmaceutical University, Tongjia Lane 24, Nanjing 210009, China.  
E-mail: zunjianzhangcpu@hotmail.com






Figure 4: The proposed fragmentation pathways of anthraquinone derivatives





Figure 5: Extracted ion chromatograms of ions at m/z 285.04 (a), 283.06 (b), 431.10 (c) and 299.02 (d); and ultraviolet spectra of some anthraquinone derivatives



**Figure S1:** Ultraviolet spectra of some gallates and tannins

analytical speed and accuracy and has been confirmed to be a very powerful tool on the comprehensive identification of both target and nontarget components. Moreover, our previous investigation had clearly indicated the relevance and usefulness of the combination of chromatographic, spectrophotometric, and mass-spectrometric analysis to detect and identify components in complicated samples.<sup>[13]</sup> Therefore, in the present study, HPLC with diode-array and IT/TOF MS detection was used for separation and identification of the components in HZ extract. Compared with previous methods, more components (74) were characterized or tentatively identified, and 29 of these compounds were reported in HZ for the first time.

## MATERIALS AND METHODS

### Reagents and chemicals

Reference standards of gallic acid (HPLC >98%), (+)-catechin (HPLC >98%), (-)-epicatechin (HPLC >98%), emodin (HPLC >98%), rhein (HPLC >98%), aloe-emodin (HPLC >98%), physcion (HPLC >98%) were purchased from the National Institutes for Food and Drug Control (Beijing, China). Emodin-1-O- $\beta$ -D-glucopyranoside (HPLC >98%), emodin-8-O- $\beta$ -D-glucopyranoside (HPLC >98%), resveratrol (HPLC >98%), and polydatin (HPLC >98%) were purchased from Shanghai Yilin Biotech Co. Ltd (Shanghai, China). Methanol of HPLC grade was purchased from Merck (Darmstadt, Germany). Formic acid (analytical reagent) was purchased from the First Chemical

Company of Nanjing (Jiangsu, China). Deionized water was prepared with a Milli-Q system (Millipore, MA, USA).

### Plant materials

The rhizome et radix of *P. cuspidatum* was collected in Hubei Province, China and authenticated by Associate Professor Rui Song (State Key Laboratory of Natural Medicine, China Pharmaceutical University, Nanjing, China). The voucher specimens (No. zzi-P-20130813) were deposited at the Herbarium of China Pharmaceutical University, Nanjing, China.

### Sample preparation

The preparation of HZ aqueous extraction took published literature as reference with some modifications.<sup>[14]</sup> 25 g of HZ was immersed in 500 mL distilled water for 1 h and heated with a heating mantle. After boiling, gentle heating was continued until the volume reduced below 250 mL. The decoction was filtered while hot and condensed to 50 mL and stored at  $-80^{\circ}\text{C}$  for later use.


### High-performance liquid chromatography-diode array detector analysis

The analysis was performed using an Agilent Series 1100 liquid chromatograph (Agilent Technologies, Palo Alto, CA, USA). Chromatography was carried out on a Shimadzu Inertsil C8-3 reversed phase column (150 mm  $\times$  2.1 mm, 3  $\mu\text{m}$ ). The mobile phase was composed of A (methanol) and B (0.1% formic acid aqueous solution) under gradient elution conditions: 5% A at 0–5 min, 5–30% A at 5–25 min, 30–50% A at 25–45 min, 50–90% A at 45–60 min, and





were as follows: Interface voltage,  $-3.5$  kV; detector voltage,  $1.7$  kV; nebulizing gas ( $N_2$ ) flow,  $1.5$  L/min; drying gas ( $N_2$ ) flow,  $100$  kPa; curved desorption line temperature and heat block temperature,  $200^\circ\text{C}$ ; ion accumulated time,  $30$  ms; precursor ion selected width,  $3.0$  amu; collision energy,  $30$ – $80\%$ . Mass spectra and chromatograms were acquired and processed with LC-MS solution version 3.6 (Shimadzu, Japan). Shimadzu's Composition Formula Predictor software was adopted to predict chemical formulas for both precursor and product ions.



**Figure S3:** The proposed fragmentation pathways of naphthalenes

## RESULTS AND DISCUSSION

### Identification of major constituents in Huzhang

Figure 1 shows the typical total ion chromatogram (TIC) of HZ extract acquired in negative mode. In the present study, 238 components were detected in the HZ extract and 74 of them had been confirmed or tentatively identified. Compounds were first classified into several categories based on the characteristic fragment ions obtained from the HPLC-IT/TOF MS analysis. The formulae of detected

components were then predicted by the formula predictor software according to the accurate mass provided by their deprotonated ions  $[\text{M} - \text{H}]^-$  or  $[\text{M} + \text{HCOO}]^-$ . Following that, the fragmentation pathways of these compounds were proposed on account of their  $\text{MS}^n$  behaviors. By comparison with the available standard samples, previously published data and the path of biosynthesis, the components were ultimately identified or putatively characterized. The obtained ultraviolet (UV) spectra of some compounds were hired for auxiliary confirmation.

### Identification of stilbenes derivatives

A total of 16 compounds were assigned to stilbenes derivatives [Tables S1 and S2], among which three components (S1, S13, S16) were first reported in HZ. By comparison with the reference substance, S4 and S10 were identified as polydatin and resveratrol, respectively. A resveratrol-O-diglucoside (S1), which showed  $[M + HCOO]^-$  ion at  $m/z$  597.18 and  $[M - H]^-$  ion at  $m/z$  551.17, was detected in HZ extract. It formed fragment ions at  $m/z$  389.12 ( $[M-H-Glu]^-$ ) and 227.07 ( $[M-H-2Glu]^-$ ). S3 showed the same  $[M + HCOO]^-$  and  $[M - H]^-$  ions with those of polydatin. Meanwhile, the MS<sup>2</sup> spectra of S3 displayed characteristic fragment ions at  $m/z$  227.07 and 185.06, indicating that it was resveratrol-O-glucoside as well. By comparing the UV spectra of S3 with resveratrol [Figure 2], it was tentatively identified as resveratrol-O-glucoside.<sup>[1,15,16]</sup> Both S8 and S9 displayed deprotonated ion at  $m/z$  541.13 and shared fragment ions at  $m/z$  227.07 and

185.06. The neutral loss of 314 Da and the dominant fragment ions at  $m/z$  313.05 and 169.01 were the symbols of the galloyl glucose. Hence, they were tentatively characterized as resveratrol 3-O-D-(6' or 2'-galloyl)-glucopyranoside.<sup>[17]</sup>

Besides the glucosides discussed above, several sulfates were detected in HZ extract. S13 and S16 all gave  $[M - H]^-$  ion at  $m/z$  307.03. The following neutral loss of 80 Da forming ions at  $m/z$  227.07 ( $[M-H-SO_3]^-$ ) and other MS<sup>2</sup> fragmentation ions consistent with those of resveratrol led to the tentative characterization of S13 and S16 as resveratrol-O-sulfates. S5 and S7 both showed deprotonated ion at  $m/z$  469.08 and characteristic MS<sup>2</sup> fragmentation ions at  $m/z$  307.02 ( $[M-H-Glu]^-$ ), 227.07 ( $[M-H-Glu-SO_3]^-$ ) and 185.06 ( $[M-H-Glu-SO_3-C_2H_2O]^-$ ), revealing that they might be sulfonylresveratrol-3-O-glucosides.<sup>[17]</sup> S11, S12, S14, and S15 shared the same precursor ion but almost distinct

**Table S1: Identification of stilbenes derivatives**

Number	$t_R$ (min)	Precursor ion	MS <sup>n</sup> (negative mode)	Identification
S1	20.173	597.18 $[M+HCOO]^-$	MS <sup>2</sup> [597.18]: 227.0680 (44.01); 321.0541 (2.52); 389.1206 (100); 551.1708 (48.79)	<b>Resveratrol-O-diglucoside</b>
S2	25.733	405.12 $[M-H]^-$	MS <sup>2</sup> [405.12]: 243.06 (100)	Piceatannol-O-glucoside
S3	25.907	435.12 $[M+HCOO]^-$ 389.12 $[M-H]^-$	MS <sup>2</sup> [435.12]: 227.07 (100); 389.12 (46.12) MS <sup>2</sup> [389.12]: 143.05 (1.49); 159.08 (1.20); 183.08 (1.05); 185.06 (1.63); 225.0526 (1.05); 227.07 (100)	Resveratrol-4'-O-β-D-glucopyranoside
S4	28.467	435.12 $[M+HCOO]^-$ 389.12 $[M-H]^-$	MS <sup>2</sup> [435.12]: 227.0679 (100); 389.12 (61.91) MS <sup>2</sup> [389.12]: 185.06 (1.17); 227.07 (100)	Polydatin*
S5	29.960	469.08 $[M-H]^-$	MS <sup>2</sup> [469.08]: 185.06 (3.64); 227.07 (100); 307.02 (6.01)	Sulfonylresveratrol-3-O-β-D-glucopyranoside
S6	31.307	469.08 $[M-H]^-$	MS <sup>2</sup> [469.08]: 227.07 (3.78); 241.00 (100); 306.02 (3.51) MS <sup>3</sup> [241.00]: 96.96 (100)	Resveratrol-3-O-β-D-(sulfonyl)-glucopyranoside
S7	32.120	469.08 $[M-H]^-$	MS <sup>2</sup> [469.08]: 183.08 (1.44); 185.06 (5.57); 227.07 (100); 269.08 (1.71); 307.02 (8.36)	Sulfonylresveratrol-3-O-β-D-glucopyranoside
S8	32.467	541.13 $[M-H]^-$	MS <sup>2</sup> [541.13]: 169.01 (17.80); 185.06 (1.94); 227.07 (73.55); 313.05 (100); 379.08 (1.06)	Resveratrol 3-O-D-(6' or 2'-galloyl)-glucopyranoside
S9	33.72	541.13 $[M-H]^-$	MS <sup>2</sup> [541.13]: 169.01 (4.01); 227.07 (1.26); 313.05 (100); 495.24 (2.88)	Resveratrol 3-O-D-(6' or 2'-galloyl)-glucopyranoside
S10	35.293	227.07 $[M-H]^-$	MS <sup>2</sup> [227.07]: 143.05 (48.19); 159.08 (21.72); 183.08 (18.20); 185.06 (100)	Resveratrol*
S11	35.507	469.08 $[M-H]^-$	MS <sup>2</sup> [469.08]: 180.98 (2.53); 222.99 (13.55); 227.07 (23.21); 241.00 (100); 281.08 (2.48); 299.09 (2.87); 307.02 (8.36) MS <sup>3</sup> [241.00]: 96.96 (100); 138.97 (14.51)	Resveratrol-3-O-β-D-(sulfonyl)-glucopyranoside
S12	36.573	469.08 $[M-H]^-$	MS <sup>2</sup> [469.08]: 166.96 (0.27); 227.07 (0.47); 241.00 (100) MS <sup>3</sup> [241.00]: 96.96 (100); 138.97 (9.25); 166.97 (3.89)	Resveratrol-3-O-β-D-(sulfonyl)-glucopyranoside
S13	38.880	307.03 $[M-H]^-$	MS <sup>2</sup> [307.03]: 143.05 (1.23); 157.06 (0.79); 159.08 (1.28); 183.08 (1.03); 185.06 (5.63); 227.07 (100)	<b>Resveratrol-O-sulfate</b>
S14	39.747	469.08 $[M-H]^-$	MS <sup>2</sup> [469.08]: 152.99 (1.21); 180.98 (2.83); 227.07 (1.75); 241.00 (100) MS <sup>3</sup> [241.00]: 96.96 (100); 180.98 (6.55)	Resveratrol-3-O-β-D-(sulfonyl)-glucopyranoside
S15	40.693	469.08 $[M-H]^-$	MS <sup>2</sup> [469.08]: 227.07 (1.93); 241.00 (100); 269.08 (3.96) MS <sup>3</sup> [241.00]: 96.96 (100)	Resveratrol-3-O-β-D-(sulfonyl)-glucopyranoside
S16	42.733	307.03 $[M-H]^-$	MS <sup>2</sup> [307.03]: 143.05 (2.93); 157.06 (2.16); 159.08 (2.16); 183.08 (1.78); 185.06 (7.02); 227.07 (100); 243.06 (1.19)	<b>Resveratrol-O-sulfate</b>

MS: Mass spectrometry. Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

fragment ions with those of S5 and S7. In their MS<sup>2</sup> spectra, the dominant ions were all at *m/z* 241.00, which was 228 Da less than the precursor ion. With the extra MS<sup>3</sup> spectra of the ion at *m/z* 241.00, we predicted it as sulfonyl glucose [Figure 3]. Therefore, these four constituents were believed to be resveratrol-3-O-(sulfonyl)-glucosides.<sup>[18,19]</sup>

## IDENTIFICATION OF ANTHRAQUINONES DERIVATIVES

A total of 24 anthraquinone derivatives were characterized or plausibly identified [Tables S3 and S4]. A3, A8, and A23 were separately characterized as emodin-1-O-β-D-glucopyranoside, emodin-8-O-β-D-glucopyranoside and emodin by direct injection of authentic standards.

In the present study, 11 hydroxyl aloë-emodin derivatives were first reported in HZ. A7, A9, A11, A16, A24 shared the [M – H]<sup>–</sup> ion at *m/z* 285.04, 16 Da more than that of aloë-emodin. The sequential loss of CO and CHO radical further indicated that they were aloë-emodin derivatives [Figure 4]. Therefore, they were tentatively identified as hydroxyl aloë-emodin. Compared with aloë-emodin, the UV spectrum of A9 displayed some changes in the shape and bathochromic shift in the band III [Figure 5], revealing that there was a β-OH on the other side of methylol group.<sup>[12]</sup> Consequently, A9 was

tentatively identified as 6-hydroxylaloë-emodin. Five isomers (A1, A2, A4, A5, A13), which could form the aglycone ion at *m/z* 285.04 after having lost 162 Da and displayed characteristic fragment ions of hydroxyl aloë-emodin, were consequently identified as hydroxyl aloë-emodin-O-glucosides. The precursor ion of A20 produced fragment ion at *m/z* 285.04 after losing 80 Da, which yield a series of characteristic ions of hydroxyl aloë-emodin at *m/z* 257.04, 241.05, and 211.04 in MS<sup>3</sup> spectrum, providing evidence for the assumption that A20 was hydroxyl aloë-emodin-O-sulfate.


Both A6 and A14 showed the [M – H]<sup>–</sup> ion at *m/z* 473.11, which was identical with that of emodin-O-(acetyl)-glucoside. Nevertheless, the MS<sup>2</sup> fragment ion at *m/z* 311.05 formed by direct loss of 162 Da from the precursor ion denied the deduction. Based on this evidence, those two compounds were plausibly identified as acetylemodin-O-glucosides. In addition, an acetylemodin (A21), which showed a deprotonated ion at *m/z* 311.06 and shared most of the fragment ions with emodin, was detected at 55.707 min.

The [M – H]<sup>–</sup> ion and a series of characteristic fragment ions of compound A10 and A17 led to the initial assumption that they were physcion and its isomer. The injection of standard sample of physcion overthrow the inference then as both of them had weaker retention than physcion. Besides, the UV spectrum of A10 was quite different from that of physcion [Figure 5]. Those comprehensive hints indicated that they were possibly two more polar isomers of physcion. Taking the retention times of emodin-1-O-β-D-glucopyranoside and emodin-8-O-β-D-glucopyranoside as reference, A10 was tentatively identified as emodin-1-questin which was first reported in HZ, while A17 was characterized as emodin-8-questin reported in literatures.<sup>[7]</sup>

A11 showed the [M – H]<sup>–</sup> ion at *m/z* 431.10, and shared most of the fragment ions with those of emodin-1 or 8-O-β-D-glucopyranoside. In the UV spectra of A11, the characteristic band III and band V of anthraquinone displayed hypochromatic shift compared with emodin [Figure 5], indicating that the β-OH of emodin was substituted. Consequently, A11 was tentatively identified as emodin-6-O-glucoside.

The formula of A18, which showed [M – H]<sup>–</sup> ion at *m/z* 299.02, was predicted as C<sub>15</sub>H<sub>7</sub>O<sub>7</sub>. In the MS<sup>2</sup> spectrum, the ion at *m/z* 255.03 indicated that a carboxyl was attached to the anthraquinone moiety. Thus, A18 was tentatively characterized as hydroxyl rhein. By comparing the UV spectra of A18 with that of rhein [Figure 5], the substitution position was found to be more likely on the

**Table S2: Structures of identified stilbenes derivatives**

				
Compound	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
<b>S1</b>	Glc	Glc	H	H
	H	Glc	OGlc	H
<b>S2</b>	Glc	OH	OH	H
	H	Glc	OH	H
	H	H	Glc	H
<b>S3</b>	H	Glc	H	H
<b>S4*</b>	Glc	H	H	H
<b>S5/S7</b>	Glc	SO <sub>3</sub> H	H	H
	Glc	H	H	SO <sub>3</sub> H
<b>S6/S11/S12/S14/S15</b>	Glc	Sulfonyl-Glc	H	H
<b>S8/S9</b>	Glc	(2"/6"-Galloyl) Glc	H	H
<b>S10*</b>	H	H	H	H
<b>S13/S16</b>	SO <sub>3</sub> H	H	H	H
	H	SO <sub>3</sub> H	H	H
	H	H	H	SO <sub>3</sub> H

Those components shown in **bold** were reported in Huzhang for the first time.  
Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.



**Table S3: Identification of anthraquinones**

Number	$t_R$ (min)	Precursor ion	MS <sup>n</sup> (negative mode)	Identification
A1	30.907	447.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [447.09]: 195.05 (0.32); 205.04 (0.27); 211.04 (1.72); 239.04 (0.72); 241.05 (1.24); 256.04 (4.07); 257.04 (0.66); 281.04 (1.96); 285.04 (100); 286.04 (11.60); 298.05 (0.60); 309.04 (5.14); 327.05 (5.14)	Hydroxyl aloe-emodin-O-glucoside
A2	39.093	447.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [447.09]: 169.07 (0.38); 197.06 (9.37); 241.05 (100); 242.05 (7.11); 255.06 (0.38); 265.04 (0.66); 283.06 (1.83); 285.04 (2.04); 403.10 (7.83); 404.11 (1.32); 429.08 (0.49); 430.10 (0.33); 448.09 (1.95); 449.11 (0.38) MS <sup>3</sup> [241.05]: 115.05 (5.05); 131.05 (3.01); 141.07 (4.55); 169.07 (28.32); 182.04 (3.01); 197.06 (100); 199.04 (3.01)	Hydroxyl aloe-emodin-O-glucoside
A3	40.373	431.10 [M-H] <sup>-</sup>	MS <sup>2</sup> [431.10]: 210.03 (1.49); 225.05 (10.14); 240.04 (1.93); 241.05 (2.27); 269.04 (100); 270.05 (5.10) MS <sup>3</sup> [269.04]: 180.05 (1.22); 181.06 (19.52); 182.04 (4.05); 197.06 (12.08); 210.03 (16.56); 224.04 (1.86); 225.05 (100); 241.0458 (23.18)	Emodin-1-O-β-D-glucopyranoside*
A4	42.467	447.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [447.09]: 185.06 (0.29); 195.05 (0.24); 198.03 (0.45); 212.04 (0.29); 213.05 (0.81); 225.05 (0.61); 226.03 (1.09); 240.04 (0.33); 241.05 (3.32); 253.05 (0.61); 271.07 (0.52); 281.04 (3.24); 284.03 (1.86); 285.04 (100); 286.04 (6.46); 309.04 (3.67); 327.05 (4.14)	Hydroxyl aloe-emodin-O-glucoside
A5	44.400	447.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [447.09]: 211.04 (0.73); 224.05 (0.62); 225.05 (0.95); 227.03 (0.62); 239.04 (1.44); 240.04 (2.30); 241.05 (1.76); 255.03 (0.95); 256.04 (0.73); 268.04 (2.67); 269.04 (7.68); 283.02 (8.90); 284.03 (100); 285.04 (19.95); 286.04 (0.77)	Hydroxyl aloe-emodin-O-glucoside
A6	44.773	473.11 [M-H] <sup>-</sup>	MS <sup>2</sup> [473.11]: 225.05 (0.81); 269.04 (1.37); 307.0534 (1.37); 311.05 (100); 312.05 (8.08); 353.05 (5.56); 353.06 (4.38) MS <sup>3</sup> [311.05]: 224.05 (18.11); 225.05 (19.69); 239.07 (15.69); 240.04 (13.31); 267.06 (35.34); 268.03 (17.32); 269.04 (100); 283.06 (21.28)	Acetylemodin-O-glucoside
A7	46.467	285.04 [M-H] <sup>-</sup>	MS <sup>2</sup> [285.04]: 197.06 (2.24); 241.05 (100); 242.05 (10.50)	Hydroxyl aloe-emodin
A8	47.493	431.10 [M-H] <sup>-</sup>	MS <sup>2</sup> [431.10]: 210.03 (1.33); 225.05 (9.30); 241.05 (1.56); 265.05 (2.35); 269.04 (100); 270.05 (10.21); 293.04 (6.21); 311.05 (4.10) MS <sup>3</sup> [269.04]: 180.06 (2.84); 181.06 (22.36); 182.04 (4.17); 196.05 (1.39); 197.06 (16.33); 207.04 (2.36); 210.03 (23.41); 224.04 (4.73); 225.05 (100); 241.05 (20.81); 270.0435 (37.09); 271.05 (1.08)	Anthraglycoside B (emodin-8-O-β-D-glucopyranoside)*
A9	50.360	285.04 [M-H] <sup>-</sup>	MS <sup>2</sup> [285.04]: 167.05 (3.56); 182.04 (1.30); 183.05 (2.44); 195.04 (16.88); 198.03 (2.06); 211.04 (100); 212.04 (9.28); 213.05 (3.65); 223.04 (4.40); 224.04 (18.79); 227.03 (1.87); 228.04 (2.81); 239.03 (11.16); 240.04 (1.87); 241.05 (71.47); 242.05 (3.46); 255.03 (13.40); 256.03 (13.12); 257.04 (54.75); 258.05 (1.12); 268.03 (11.47); 287.04 (14.54)	Hydroxyl aloe-emodin
A10	50.707	283.06 [M-H] <sup>-</sup>	MS <sup>2</sup> [283.06]: 240.04 (100); 241.04 (8.03); 268.04 (2.03); 269.03 (0.84); 282.05 (1.18); 284.06 (1.02)	Emodin-1-questin
A11	50.840	431.10 [M-H] <sup>-</sup>	MS <sup>2</sup> [431.10]: 225.05 (2.49); 265.04 (1.18); 269.04 (100); 270.05 (4.73); 282.05 (8.51); 283.06 (1.66); 293.04 (3.67); 311.05 (12.75); 335.06 (1.18)	Emodin-6-O-glucoside
A12	51.307	285.04 [M-H] <sup>-</sup>	MS <sup>2</sup> [285.04]: 167.05 (9.20); 195.04 (14.63); 211.04 (91.23); 212.04 (17.30); 224.06 (15.30); 239.03 (13.29); 241.05 (100); 242.06 (7.86); 255.02 (9.20); 256.04 (10.62); 257.04 (28.78); 268.03 (7.86)	Hydroxyl aloe-emodin
A13	51.413	447.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [447.09]: 197.06 (0.53); 198.03 (0.64); 213.06 (2.14); 217.05 (0.96); 240.04 (0.65); 241.05 (1.23); 253.05 (0.44); 257.04 (4.50); 270.06 (0.36); 271.05 (0.36); 281.04 (2.87); 285.04 (100); 286.04 (5.05); 298.04 (0.48); 309.04 (4.33); 327.05 (4.82)	Hydroxyl aloe-emodin glucoside
A14	51.880	473.11 [M-H] <sup>-</sup>	MS <sup>2</sup> [473.11]: 225.05 (6.89); 253.05 (1.60); 265.04 (2.83); 269.04 (100); 270.06 (5.26); 280.03 (1.89); 293.04 (10.95); 311.05 (4.46); 335.05 (1.60); 395.08 (1.60); 413.08 (6.49) MS <sup>3</sup> [269.04]: 225.05 (100); 241.05 (15.23)	Acetylemodin-O-glucoside

*Contd..*

Table S3: Contd...

Number	$t_R$ (min)	Precursor ion	MS <sup>n</sup> (negative mode)	Identification
A15	53.413	517.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [517.09]: 210.03 (0.04); 225.05 (0.47); 241.05 (0.09); 269.04 (30.96); 270.05 (2.41); 293.04 (1.04); 311.06 (1.47); 431.10 (1.86); 473.11 (100); 474.30 (7.02); 475.11 (4.19)	Emodin-8-O-(6'-O-malonyl)-glucoside
A16	53.547	285.04 [M-H] <sup>-</sup>	MS <sup>2</sup> [285.04]: 167.05 (9.20); 195.04 (14.63); 211.04 (91.23); 212.04 (17.30); 224.04 (15.30); 239.03 (13.29); 241.05 (100); 242.06 (7.86); 255.02 (9.20); 256.04 (10.62); 257.04 (28.78); 268.03 (7.86)	Hydroxyl aloe-emodin
A17	54.280	283.06 [M-H] <sup>-</sup>	MS <sup>2</sup> [283.06]: 240.04 (100); 241.05 (5.23); 268.03 (4.48)	Emodin-8-questin
A18	54.480	299.02 [M-H] <sup>-</sup>	MS <sup>2</sup> [299.02]: 167.05 (24.44); 183.05 (18.06); 193.03 (1.85); 199.04 (1.85); 210.03 (8.03); 211.04 (100); 212.04 (2.26); 227.03 (32.43); 237.01 (1.22); 255.03 (67.42); 256.03 (6.16); 271.02 (5.55); 300.02 (1.85)	<b>6-hydroxylrhein</b>
A19	54.813	511.05 [M-H] <sup>-</sup>	MS <sup>2</sup> [511.05]: 210.03 (1.00); 225.06 (6.46); 241.00 (4.86); 241.05 (1.31); 269.04 (100); 270.05 (20.78); 431.09 (32.02)	<b>Emodin-O-(sulfonyl)-glucoside</b>
A20	55.280	365.00 [M-H] <sup>-</sup>	MS <sup>2</sup> [365.00]: 285.04 (100) MS <sup>3</sup> [285.04]: 195.05 (11.89); 198.02 (3.10); 211.04 (100); 212.04 (4.38); 213.05 (7.18); 223.04 (3.10); 224.05 (9.68); 228.04 (5.00); 239.03 (9.99); 240.05 (3.10); 241.05 (79.78); 255.03 (7.18); 256.03 (9.70); 257.04 (69.91); 268.04 (5.92)	<b>Hydroxy aloe-emodin-O-sulfate</b>
A21	55.707	311.06 [M-H] <sup>-</sup>	MS <sup>2</sup> [311.06]: 195.05 (11.25); 196.05 (13.45); 224.05 (100); 225.05 (53.52); 239.07 (20.10); 240.04 (45.57); 265.13 (11.25); 267.07 (25.60); 268.03 (15.64); 269.04 (12.35)	<b>Acetylemodin</b>
A22	56.533	511.05 [M-H] <sup>-</sup>	MS <sup>2</sup> [511.05]: 222.99 (2.24); 225.06 (6.04); 241.00 (73.67); 241.05 (2.29); 269.04 (100); 270.05 (14.29); 311.05 (6.54); 341.06 (1.53); 431.09 (37.77)	<b>Emodin-O-(sulfonyl)-glucoside</b>
A23	57.547	269.04 [M-H] <sup>-</sup>	MS <sup>2</sup> [269.04]: 179.05 (1.04); 180.06 (3.42); 181.06 (18.97); 182.04 (3.68); 197.06 (14.37); 207.04 (2.39); 210.03 (23.51); 224.04 (3.67); 225.05 (100); 226.06 (1.22); 227.03 (1.04); 241.05 (23.78); 270.05 (58.93); 271.05 (4.15)	Emodin*
A24	59.347	285.04 [M-H] <sup>-</sup>	MS <sup>2</sup> [285.04]: 189.05 (14.03); 213.05 (29.27); 217.05 (16.77); 241.05 (36.21); 257.04 (100)	Hydroxyl aloe-emodin

MS: Mass spectrometry. Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

other side of the carboxyl. Since the C-6 position was easier to be substituted, A18 was consequently qualified as 6-hydroxylrhein.

A19 and A22 exhibited the same deprotonated ion at  $m/z$  511.05 and roughly identical fragment ions despite little difference in intensity. A consecutive loss of  $\text{SO}_3$  and glucoside led to the formation of aglycone ion at  $m/z$  269.04, which was believed to be emodin proved by the diagnostic ions of emodin at  $m/z$  241.05 and 225.05. In the meantime, the fragment ion at  $m/z$  241.00 (sulfonyl glucose) provided further clue that these two compounds were likely to be emodin-O-(sulfonyl)-glucosides.

#### Identification of gallates and tannins



Gallates and tannins can be found in most of the Polygonaceae plants. In the present study, 19 gallates and tannins were detected, and 18 of them were characterized [Tables S5 and S6]. By comparing the retention times, characteristic fragment ions and UV spectra [Figure S1] with references, G3, G12, G15 were identified as gallic acid, (+)-catechin, and (-)-epicatechin,

respectively. Five isomers (G1, G2, G4, G5, G6, and G8), which displayed  $[\text{M} - \text{H}]^-$  ion at  $m/z$  331.07 and shared the same characteristic ions at  $m/z$  169.01 and 125.02 [Figure S2], were characterized as galloyl glucose.

Protocatechuic acid was once detected in HZ.<sup>[20]</sup> In our present study, a protocatechuic acid-O-glucoside (G7) rather than the aglycone itself was found in the herb. The  $[\text{M} - \text{H}]^-$  ion at  $m/z$  315.07 first lost 162 Da to form the deprotonated ion of protocatechuic acid, which produced the ion at  $m/z$  108.02 by a carboxyl crashed down.

G9 and G13 yielded the  $[\text{M} - \text{H}]^-$  ion at  $m/z$  451.12, which could give rise to a further fragment ion at  $m/z$  289.07 after a loss of 162 Da. G13 was eluted between (+)-catechin and (-)-epicatechin, provided evidence that it was more likely to be (-)-epicatechin glucoside under the consideration that (+)-catechin glucoside should have weaker retention than (+)-catechin on reverse phase chromatography. Therefore, G13 was believed to be (-)-epicatechin glucoside while G9 was tentatively

**Table S4: Structures of identified anthraquinones derivatives**

Compound	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>	R <sub>8</sub>
								
<b>A1/A2/A4/A5/A13</b>	H	Glc	CH <sub>2</sub> OH	OH	H	H	H	H
	H	Glc	CH <sub>2</sub> OH	H	OH	H	H	H
	H	Glc	CH <sub>2</sub> OH	H	H	OH	H	H
	H	Glc	CH <sub>2</sub> OH	H	H	H	OH	H
	H	Glc	CH <sub>2</sub> OH	H	H	H	H	OH
A3*	H	Glc	OH	CH <sub>3</sub>	H	H	H	H
<b>A6/A14</b>	COCH <sub>3</sub>	Glc	OH	CH <sub>3</sub>	H	H	H	H
	Glc	OCH <sub>3</sub>	OH	CH <sub>3</sub>	H	H	H	H
A7/A9/A12/A16/A24	H	H	CH <sub>2</sub> OH	OH	H	H	H	H
	H	H	CH <sub>2</sub> OH	H	OH	H	H	H
	H	H	CH <sub>2</sub> OH	H	H	OH	H	H
	H	H	CH <sub>2</sub> OH	H	H	H	OH	H
	H	H	CH <sub>2</sub> OH	H	H	H	H	OH
A8*	Glc	H	OH	CH <sub>3</sub>	H	H	H	H
<b>A10</b>	<b>H</b>	<b>CH<sub>3</sub></b>	OH	CH <sub>3</sub>	H	H	H	H
<b>A11</b>	H	H	OGlc	CH <sub>3</sub>	H	H	H	H
A15	<b>(6'-malonyl) Glc</b>	<b>H</b>	OH	CH <sub>3</sub>	H	H	H	H
A17	<b>CH<sub>3</sub></b>	<b>H</b>	OH	CH <sub>3</sub>	H	H	H	H
								
<b>A18</b>	H	H	COOH	OH	H	H	H	H
<b>A19/A22</b>	Sulfonyl-Glc	H	OH	CH <sub>3</sub>	H	H	H	H
	H	Sulfonyl-Glc	OH	CH <sub>3</sub>	H	H	H	H
<b>A20</b>	Glc	H	CH <sub>2</sub> OH	OH	H	H	H	H
	H	Glc	CH <sub>2</sub> OH	OH	H	H	H	H
	H	H	CH <sub>2</sub> OH	OGlc	H	H	H	H
<b>A21</b>	COCH <sub>3</sub>	H	OH	CH <sub>3</sub>	H	H	H	H
	H	COCH <sub>3</sub>	OH	CH <sub>3</sub>	H	H	H	H
	H	H	OCOCH <sub>3</sub>	CH <sub>3</sub>	H	H	H	H
A23*	H	H	OH	CH <sub>3</sub>	H	H	H	H

Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

**Table S5: Identification of gallates and tannins**

Number	t <sub>R</sub> (min)	Precursor ion	MS <sup>n</sup> (negative mode)	Identification
G1	3.893	331.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [331.07]: 168.01 (4.73); 169.01 (11.86); 211.02 (6.61); 271.04 (100); 272.04 (3.67)	Galloyl glucose
G2	4.267	331.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [331.07]: 125.02 (33.86); 169.01 (100); 170.02 (4.91)	Galloyl glucose
G3	4.560	169.01 [M-H] <sup>-</sup> 339.03 [2M-H] <sup>-</sup>	MS <sup>2</sup> [169.01]: 125.02 (100); 126.03 (7.26) MS <sup>2</sup> [339.03]: 125.02 (7.70); 169.01 (100)	Gallic acid*
G4	5.080	331.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [331.07]: 125.03 (2.61); 169.01 (10.01); 211.02 (8.68); 271.04 (100); 271.19 (1.66); 272.04 (2.92)	Galloyl glucose
G5	7.707	331.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [331.07]: 151.00 (92.92); 169.01 (100); 193.02 (71.67)	Galloyl glucose
G6	8.933	331.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [331.07]: 169.01 (100)	Galloyl glucose
G7	9.240	315.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [315.07]: 108.02 (22.38); 152.01 (49.26); 153.02 (100); 163.04 (16.02)	<b>Protocatwchuic acid-O-glucoside</b>
G8	12.947	331.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [331.07]: 125.02 (19.31); 150.00 (37.65); 151.00 (6.44); 167.04 (7.42); 168.01 (100); 169.01 (25.42); 193.02 (4.12); 313.05 (33.03); 314.05 (3.47)	Galloyl glucose

Contd..

Table S5: Contd...

Number	$t_R$ (min)	Precursor ion	MS <sup>n</sup> (negative mode)	Identification
G9	14.453	451.12 [M-H] <sup>-</sup>	MS <sup>2</sup> [451.12]: 203.07 (1.05); 205.05 (1.42); 245.08 (10.39); 289.06 (100); 290.07 (2.10)	(+)-Catechin-5-O-glucoside
G10	14.760	577.13 [M-H] <sup>-</sup>	MS <sup>2</sup> [577.13]: 245.04 (3.28); 245.08 (3.28); 273.04 (2.89); 287.05 (5.25); 289.07 (40.01); 299.05 (5.64); 407.07 (100); 408.08 (6.36); 425.08 (86.07); 426.08 (2.69); 451.10 (21.27)	Procyanidin B
G11	16.080	577.13 [M-H] <sup>-</sup>	MS <sup>2</sup> [577.13]: 203.07 (1.20); 205.05 (1.44); 245.04 (4.69); 245.08 (10.30); 269.04 (1.09); 273.04 (7.08); 281.04 (3.35); 285.04 (1.17); 287.05 (12.28); 288.06 (1.64); 289.07 (49.53); 290.07 (7.19); 299.05 (12.53); 300.06 (1.17); 339.08 (1.05); 407.07 (100); 408.08 (24.51); 425.08 (57.96); 426.08 (23.52); 449.08 (1.01); 451.10 (11.69); 452.10 (5.21); 578.13 (8.99); 579.13 (9.96)	Procyanidin B
G12	16.733	289.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [289.07]: 109.03 (16.59); 125.02 (10.59); 137.02 (16.75); 146.04 (2.17); 151.0402 (13.57); 159.0411 (11.50); 161.0600 (55.15); 164.0096 (12.76); 165.0185 (12.13); 175.0435 (12.76); 179.0335 (23.72); 187.0381 (30.07); 188.0458 (17.38); 203.0687 (100); 205.0475 (95.94); 245.0771 (93.02); 290.0705 (16.26)	(+)-Catechin*
G13	19.160	451.12 [M-H] <sup>-</sup>	MS <sup>2</sup> [451.12]: 179.04 (3.30); 203.07 (6.26); 205.05 (3.93); 245.08 (20.67); 289.07 (100); 290.07 (5.65)	(-)-Epicatechin-O-glucoside
G14	19.947	577.13 [M-H] <sup>-</sup>	MS <sup>2</sup> [577.13]: 203.07 (1.48); 245.04 (3.08); 245.08 (6.97); 269.04 (1.53); 273.04 (6.76); 281.04 (3.33); 285.04 (1.39); 287.05 (12.02); 289.07 (36.71); 290.07 (3.41); 299.05 (10.40); 300.06 (1.14); 339.08 (1.35); 407.07 (100); 408.08 (25.22); 425.08 (64.44); 425.26 (1.36); 426.08 (31.94); 451.10 (8.16); 452.10 (3.16)	Procyanidin B
G15	21.427	289.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [289.07]: 109.03 (6.06); 123.05 (2.92); 125.02 (16.38); 135.04 (2.63); 137.02 (13.96); 146.04 (4.07); 149.03 (1.78); 151.04 (10.70); 159.04 (6.63); 161.06 (50.68); 162.03 (3.21); 164.01 (12.40); 165.02 (9.41); 167.03 (5.13); 175.04 (11.11); 175.07 (5.42); 177.06 (1.78); 179.03 (21.37); 180.05 (1.50); 185.06 (1.93); 186.06 (1.50); 187.04 (24.36); 188.05 (17.81); 199.07 (2.36); 203.07 (100); 204.07 (2.08); 205.05 (82.12); 206.05 (3.21); 212.04 (4.07); 217.05 (1.78); 227.07 (8.05); 230.06 (2.78); 231.03 (7.19); 245.08 (77.68); 246.08 (11.68); 247.06 (8.69); 271.06 (3.07); 290.07 (21.08)	(-)-Epicatechin*
G16	23.840	729.14 [M-H] <sup>-</sup>	MS <sup>2</sup> [729.14]: 243.03 (1.92); 245.04 (1.51); 245.08 (3.82); 255.03 (1.46); 269.04 (1.59); 271.06 (2.81); 285.03 (2.14); 287.05 (5.82); 289.07 (40.73); 290.07 (4.73); 299.05 (2.58); 303.05 (2.03); 331.05 (1.43); 389.06 (4.14); 407.07 (100); 407.25 (2.61); 407.46 (1.46); 408.08 (19.58); 409.08 (1.57); 425.08 (2.48); 433.09 (2.20); 441.08 (34.46); 442.08 (12.75); 443.08 (1.65); 451.10 (27.97); 452.10 (7.81); 541.11 (1.35); 559.09 (23.75); 560.09 (12.78); 561.10 (2.38); 577.11 (9.08); 578.11 (6.48); 603.11 (8.19); 604.11 (5.52); 730.14 (13.17); 731.14 (12.09)	Procyanidin-B-1-3'-O-gallate/ Procyanidin-B-5-3'-O-gallate
G17	27.787	441.08 [M-H] <sup>-</sup>	MS <sup>2</sup> [441.08]: 137.02 (0.46); 145.02 (0.72); 149.03 (0.55); 161.06 (0.81); 165.02 (1.08); 167.00 (0.55); 169.01 (39.87); 170.02 (2.61); 179.03 (2.92); 187.04 (0.50); 193.01 (15.28); 194.01 (0.64); 203.07 (4.88); 205.01 (1.23); 205.04 (3.64); 211.04 (0.55); 213.02 (0.55); 227.07 (0.77); 245.08 (13.22); 253.04 (3.19); 259.06 (2.38); 271.06 (6.58); 272.06 (1.58); 289.07 (100); 290.07 (17.46); 303.05 (4.12); 331.04 (7.34); 442.08 (9.46); 443.08 (2.38)	(+)-Catechin or (-)-epicatechin gallate
G18	29.787	441.08 [M-H] <sup>-</sup>	MS <sup>2</sup> [441.08]: 169.01 (29.90); 193.01 (20.83); 203.07 (1.62); 205.01 (1.63); 205.04 (1.63); 245.08 (7.43); 253.05 (1.63); 259.06 (2.16); 271.06 (7.03); 287.05 (1.90); 289.07 (100); 289.22 (1.34); 303.05 (2.84); 331.04 (4.73)	(+)-Catechin or (-)-epicatechin gallate
G19	34.147	505.12 [M-H] <sup>-</sup>	MS <sup>2</sup> [505.12]: 161.05 (1.63); 189.05 (6.73); 203.07 (3.55); 205.05 (3.07); 215.03 (59.74); 227.07 (1.80); 233.04 (9.32); 245.08 (13.63); 253.05 (2.13); 271.06 (2.92); 289.07 (100); 290.06 (1.63); 367.08 (4.56)	Not identified

MS: Mass spectrometry. Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.





identified as (+)-catechin-5-O-glucoside.<sup>[21]</sup> Another pair of positional isomers, G17 and G18, gave the deprotonated ion at  $m/z$  441.08. In the MS<sup>2</sup> spectra, the dominant ion at  $m/z$  289.07 and other characteristic fragment ions implied that they were (+)-catechin or (-)-epicatechin derivatives. Since the precursor ion was 152 Da more than the dominant fragment ion at  $m/z$  289.07, thus guiding

us to tentatively identify G17 and G18 as (+)-catechin or (-)-epicatechin gallates. The fragment ion at  $m/z$  169.01 also provided evidence for this assumption.

Three enantiomers of procyanidin B (G10, G11, and G14) were found in the EICs at  $m/z$  577.13. The deprotonated molecule of these compounds could further yield the most



**Table S6: Structures of identified gallates and tannins**

					
	$R_1$		$R_2$		
G1/G2/G4/G5/G6/G8	OH		Glc		
G3*	OH		H		
G7	H		Glc		
	$R_1$	$R_2$	$R_3$	$R_4$	$R_5$
					
G9	H	Glc	H	H	H
G12*	H	H	H	H	H
G17/G18	Galloyl	H	H	H	H
	H	Galloyl	H	H	H
	H	H	Galloyl	H	H
	H	H	H	Galloyl	H
	H	H	H	H	Galloyl
					
<b>G13</b>	Glc	H	H	H	H
	H	Glc	H	H	H
	H	H	Glc	H	H
	H	H	H	Glc	H
	H	H	H	H	Glc
G15*	H	H	H	H	H
G17/G18	Galloyl	H	H	H	H
	H	Galloyl	H	H	H
	H	H	Galloyl	H	H
	H	H	H	Galloyl	H
	H	H	H	H	Galloyl
<b>Possible structures</b>					
G16					
					

Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

abundant ion at  $m/z$  407.07, which might result from Retro Diels-Alder reaction (RDA) and the following loss of  $H_2O$ . The fragment ion at  $m/z$  289.07 can be attributed to the quinone methide (QM) fission cleavage of the type-B interflavan bond resulting from the loss of an (+)-catechin or (-)-epicatechin residue.<sup>[22]</sup> Besides, a type-B galloylated procyanidin (G16) with the precursor ion at  $m/z$  729.14 was detected at 23.840 min. In the  $MS^n$  spectrum, RDA reaction, heterocyclic ring fission, and QM fission cleavage had led to the formation of characteristic ions at  $m/z$  407.07, 451.10, and 289.07 [Figure S2].

### Identification of naphthalene derivatives

Six naphthalene derivatives were detected and characterized in HZ [Tables S7 and S8]. Five of them (N1, N2, N3, N5, N6) formed the same fragment ion at  $m/z$  245.08, which was predicted as torachrysone supported by the characteristic fragment ions at  $m/z$  230.05, 215.03, 187.03, and 159.05 [Figure S3]. N1 was identified as torachrysone-8-O- $\beta$ -D-glucoside.<sup>[23]</sup> N2 and N5 were characterized as two positional isomers of torachrysone-8-O-(acetyl)-glucosides since they directly lost 204 Da (acetyl glucose) and formed the base peak at  $m/z$  245.08. N3 and N6 shared the same  $[M - H]^-$  ion at  $m/z$  487.09, which was 80 Da more than that of N1, indicating that the two compounds were highly likely the sulfate ester of N1, and that the sulfate moiety was connected to glucose since there was no fragment ion which formed by directly loss of hexoside.

The  $MS^2$  spectrum of N4 was entirely different from those of the five components discussed above. The  $[M - H]^-$  ion at  $m/z$  339.02 lost 80 Da and yielded the dominant ion at  $m/z$  259.06 [Figure S3], which was tentatively identified as 2-methoxy-6-acethyl-7-methyljuglone.<sup>[20]</sup> Therefore, N4 was tentatively identified as 2-methoxy-6-acethyl-7-methyljuglone-5-O-sulfate. Other fragment ions at  $m/z$  244.04 ( $[M - H - CH_3]^-$ ) and 231.06 ( $[M - H - CO]^-$ ) provided further evidence for the conclusion.

### Identification of other compounds

Totally, 20 components which could hardly be assigned to categories above were detected, and 10 of them were tentatively identified [Tables S9 and S10]. O1 and O5 were characterized as critic acid<sup>[24]</sup> and 5, 7-dihydroxy-1 (3H)-isobenzofuran-one,<sup>[25]</sup> respectively. O4 and O6 were believed to be glucosides of O5 since the precursor ion of them was 162 Da more than that of O5, and they share the other fragment ions with O5.



O6 displayed the deprotonated ion at  $m/z$  329.09, which produced the ion at  $m/z$  167.03 after losing 162Da, indicating that it was a glucoside. The aglycone was predicted as vanillic acid proved by the ion at  $m/z$

**Table S7: Identification of naphthalene derivatives**

Number	$t_R$ (min)	Precursor ion	MS <sup>n</sup> (negative mode)	Identification
N1	43.373	407.13 [M-H] <sup>-</sup>	MS <sup>2</sup> [407.13]: 215.03 (1.41); 230.06 (13.92); 231.06 (0.47); 245.08 (100); 246.08 (15.42); 247.08 (0.11); 408.13 (3.82); 409.14 (0.88) MS <sup>3</sup> [245.08]: 202.06 (0.37); 215.03 (4.79); 230.06 (100); 231.06 (5.30); 246.08 (6.69)	Torachrysone-8-O-β-D-glucoside
N2	48.587	449.14 [M-H] <sup>-</sup>	MS <sup>2</sup> [449.14]: 215.03 (2.00); 230.05 (11.87); 245.08 (100) MS <sup>3</sup> [245.08]: 215.0336 (2.78); 230.0557 (100)	Torachrysone-8-O-(acetyl)-glucoside
N3	48.707	487.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [487.09]: 159.05 (0.19); 187.03 (0.19); 202.06 (0.19); 215.03 (7.10); 216.04 (0.41); 230.05 (27.01); 231.06 (1.32); 241.00 (1.19); 245.08 (100); 246.08 (10.17); 287.08 (0.32); 407.13 (6.75); 408.13 (1.45); 488.08 (5.34); 489.08 (2.26) MS <sup>3</sup> [245.08]: 215.03 (4.38); 230.05 (100); 231.06 (0.57)	<b>Torachrysone-8-O-(sulfonyl)-glucoside</b>
N4	49.160	339.02 [M-H] <sup>-</sup>	MS <sup>2</sup> [339.02]: 231.06 (2.74); 244.04 (7.23); 259.06 (100); 260.06 (2.23)	<b>2-Methoxy-6-acethyl-7-methyljuglone-5-O-sulfate</b>
N5	51.133	449.14 [M-H] <sup>-</sup>	MS <sup>2</sup> [449.14]: 215.03 (1.62); 230.06 (6.92); 245.08 (100); 246.08 (3.02) MS <sup>3</sup> [245.08]: 215.03 (3.11); 230.06 (100)	Torachrysone-8-O-(acetyl)-glucoside
N6	52.387	487.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [487.09]: 215.03 (2.55); 230.05 (10.96); 245.08 (100) MS <sup>3</sup> [245.08]: 215.03 (4.15); 230.06 (100)	<b>Torachrysone-8-O-(sulfonyl)-glucoside</b>

MS: Mass spectrometry. Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

**Table S8: Structures of naphthalene derivatives**

	
	<b>R</b>
N1	Glc
N2/N5	Acetyl-Glc
<b>N3/N6</b>	Sulfonyl-Glc
<b>Possible structures</b>	
<b>N4</b>	

Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

123.05 ([M-H-Glu-CO<sub>2</sub>]).<sup>[26]</sup> Meanwhile, the occurrence of the two characteristic ions at m/z 239.05 ([M-H-90 Da]<sup>-</sup>) and 209.04 ([M-H-120 Da]<sup>-</sup>) proved that O6 was likely to be vanillic acid-C-glucoside though the exact substitution position remained to be further confirmed.

O7 was characterized as a sulfate since the [M - H]<sup>-</sup> ion lost 80 Da directly and formed ion at m/z 233.04, which was identical with the deprotonated ion of 5-carboxymethyl-7-hydroxy-2-methyl chromone.<sup>[27]</sup> Another sulfate, O8, shared the same fragment ion at m/z 189.05 with O7, indicating they might have the same 2, 5-dimethyl-7-hydroxylchromone nucleus. The most

abundant ion at m/z 231.06 prompted that there might be an acetyl on the heterocyclic nucleus or connected to the hydroxyl group. Since sulfation usually acted on the hydroxyl group, the acetyl was more possibly substituted on the heterocyclic nucleus. Considering the electronic effect and steric hindrance, O8 was tentatively characterized as 3-acetyl-2, 5-dimethylchromone-7-O-sulfate. O9 displayed the [M - H]<sup>-</sup> ion at m/z 269.02 and the fragment ion at m/z 189.05, thus indicating that the compound might be 2,5-dimethylchromone-7-O-sulfate.

The [M - H]<sup>-</sup> at m/z 285.01 of O10 directly lost 80 Da, forming an ion at m/z 205.05, which was predicted as C<sub>11</sub>H<sub>9</sub>O<sub>4</sub><sup>-</sup>. The aglycone was tentatively characterized as 7-hydroxy-4-methoxy-5-methylcoumarin.<sup>[7]</sup> Consequently, O10 was tentatively identified as 4-methoxy-5-methylcoumarin-7-O-sulfate.

## SUMMARY





A relatively comprehensive chemical profiling of HZ was achieved by combination of HPLC-DAD and HPLC-IT/TOF analysis. 74 compounds including stilbenes, anthraquinones, gallates and tannins, naphthalenes along with some other minor components were identified or presumed based on their accurate mass, fragment patterns, and characteristic UV spectra. Besides, 29 of these components were reported in HZ for the first time. Although the substitution positions of some components remained to be further confirmed by other ancillary qualitative methods,

**Table S9: Identification of other compounds**

Number	<i>t<sub>R</sub></i> (min)	Precursor ion	MS <sup>n</sup> (negative mode)	Identification
O1	2.573	191.02 [M-H] <sup>-</sup>	MS <sup>2</sup> [191.02]: 87.01 (1.41); 110.97 (1.21); 111.01 (100); 111.09 (1.61); 173.01 (2.12)	Citric acid
O2	5.787	327.07 [M-H] <sup>-</sup> 373.08 [M+HCOO] <sup>-</sup>	MS <sup>2</sup> [327.07]: 165.02 (100) MS <sup>2</sup> [373.08]: 165.02 (100); 327.07 (25.37) MS <sup>3</sup> [165.02]: 121.03 (100)	<b>5,7-dyhydroxy-1 (3H)-isobenzofuran-one-O-glucoside</b>
O3	10.20	301.09 [M-H] <sup>-</sup> 347.10 [M+HCOO] <sup>-</sup>	MS <sup>2</sup> [301.09]: 139.04 (100) MS <sup>2</sup> [347.10]: 124.02 (6.83); 139.04 (100); 161.04 (26.99); 301.09 (84.94)	Tachioside/isotachioside
O4	10.480	327.07 [M-H] <sup>-</sup>	MS <sup>2</sup> [327.07]: 121.03 (6.33); 165.02 (100); 189.02 (1.01); 207.03 (1.32)	<b>5,7-dyhydroxy-1 (3H)-isobenzofuran-one-O-glucoside</b>
O5	13.267	165.02 [M-H] <sup>-</sup>	MS <sup>2</sup> [165.02]: 77.04 (17.38); 93.04 (49.74); 121.03 (100); 122.04 (28.27); 166.02 (95.92)	5,7-dyhydroxy-1 (3H)-isobenzofuran-one
O6	15.213	329.09 [M-H] <sup>-</sup>	MS <sup>2</sup> [329.09]: 123.05 (1.99); 167.03 (23.66); 209.04 (100); 210.04 (2.69); 239.05 (7.89)	<b>Vanillic acid-C-glucoside</b>
O7	30.240	313.00 [M-H] <sup>-</sup>	MS <sup>2</sup> [313.00]: 147.04 (0.38); 189.05 (100); 190.06 (2.14); 233.04 (55.31); 234.05 (4.97); 314.00 (1.67)	<b>5-carboxymethyl-2-methylchromone-7-O-sulfate</b>
O8	35.320	311.02 [M-H] <sup>-</sup>	MS <sup>2</sup> [311.02]: 189.05 (2.95); 231.06 (100); 232.07 (12.02)	<b>3-acetyl-2,5-dimethylchromone-7-O-sulfate</b>
O9	42.373	269.02 [M-H] <sup>-</sup>	MS <sup>2</sup> [269.02]: 189.05 (100); 189.18 (1.99); 189.26 (0.65); 189.32 (1.50); 189.47 (0.53); 190.06 (3.08) MS <sup>3</sup> [189.05]: 146.04 (100); 174.03 (71.81)	<b>2,5-dimethylchromone-7-O-sulfate</b>
O10	43.240	285.01 [M-H] <sup>-</sup>	MS <sup>2</sup> [285.01]: 161.03 (0.20); 162.03 (0.20); 190.03 (9.42); 205.05 (100); 206.05 (2.93); 286.01 (5.95)	<b>4-methoxy-5-methylcoumarin-7-O-sulfate</b>



MS: Mass spectrometry. Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

**Table S10: Structures of other compounds**

			
	<i>R</i> <sub>1</sub>	<i>R</i> <sub>2</sub>	
<b>O2/O4</b>	Glc	H	
	H	Glc	
O5	H	H	
			
	<i>R</i> <sub>1</sub>	<i>R</i> <sub>2</sub>	<i>R</i> <sub>3</sub>
<b>O7</b>	H	COOH	SO <sub>3</sub> H
<b>O8</b>	COCH <sub>3</sub>	H	SO <sub>3</sub> H
<b>O9</b>	H	H	SO <sub>3</sub> H
O1			
O3			

Contd..

**Table S10: Contd...**

			
	<i>R</i> <sub>1</sub>	<i>R</i> <sub>2</sub>	<i>R</i> <sub>3</sub>
<b>O6</b>	Glc	H	H
	H	Glc	H
	H	H	Glc
<b>Possible structures</b>			
<b>O10</b>			

Those components shown in **bold** were reported in Huzhang for the first time. Those components marked with an asterisk (\*) were confirmed by comparison with authentic reference.

the subtle combination of HPLC-DAD and HPLC-IT/TOF method established in the present study could provide reference method for quality control of HZ and provide a basis for further metabolic studies *in vivo* of HZ. In addition, the high sensitive method could yet be regarded as a new perspective of componential analysis of other HMs, especially those containing components with chromophores.

## REFERENCES

- Jayatilake GS, Jayasuriya H, Lee ES, Koonchanok NM, Geahlen RL, Ashendel CL, et al. Kinase inhibitors from *Polygonum cuspidatum*. J Nat Prod 1993;56:1805-10.
- Park CS, Lee YC, Kim JD, Kim HM, Kim CH. Inhibitory effects of *Polygonum cuspidatum* water extract (PCWE) and its component resveratrol [correction of rasveratrol] on acyl-coenzyme A-cholesterol acyltransferase activity for cholesteryl ester synthesis in HepG2 cells. Vascu Pharmacol 2004;40:279-84.
- Leu YL, Hwang TL, Hu JW, Fang JY. Anthraquinones from *Polygonum cuspidatum* as tyrosinase inhibitors for dermal use. Phytother Res 2008;22:552-6.
- Lee MH, James LT, Wang HY, Chang CC, Lin CC, Lin HY. Extraction of resveratrol from *Polygonum cuspidatum* with magnetic orcinol-imprinted poly (ethylene-co-vinyl alcohol) composite particles and their *in vitro* suppression of human osteogenic sarcoma (HOS) cell line. J Mater Chem 2012;22:24644-51.
- Fu Q, Wang H, Lan Y, Li S, Hashi Y, Chen S. High-throughput and sensitive screening of compounds with deoxyribonucleic acid-binding activity by a high-performance liquid chromatography-tandem mass spectrometry-fluorescence detection technique using palmitate as a fluorescence probe. J Chromatogr A 2014;1323:123-34.
- Zhang H, Li C, Kwok ST, Zhang QW, Chan SW. A Review of the pharmacological effects of the dried root of *Polygonum cuspidatum* (Hu Zhang) and its constituents. Evid Based Complement Alternat Med 2013;2013:208349.
- Yi T, Zhang H, Cai Z. Analysis of Rhizoma Polygoni Cuspidati by HPLC and HPLC-ESI/MS. Phytochem Anal 2007;18:387-92.
- Xia AJ, Zhang H, Jia J, Cai YF, Zhang GQ. Determination of multi-index components and quality control of Polygoni Cuspidati Rhizoma et Radix. Chin Tradit Herb Drugs 2011;09:1761-5.
- Sun J, Chen XQ, Jiang XY, Yu J. Extraction and GC/MS analysis of volatile oil from *Polygonum cuspidatum*. J Chin Mass Spectrom Soc 2006;27:242-5.
- Xiao K, Xuan L, Xu Y, Bai D, Zhong D. Constituents from *Polygonum cuspidatum*. Chem Pharm Bull (Tokyo) 2002;50:605-8.
- Huang S, Zhang GQ, Lou ZY, Wang B, Zhao L, Gu L, et al. Study on HPCE fingerprint of Rhizoma Polygoni Cuspidati. Chin J Pharm Anal 2006;01:24-6.
- Chen XF, Wu HT, Tan GG, Zhu ZY, Chai YF. Liquid chromatography coupled with time-of-flight and ion trap mass spectrometry for qualitative analysis of herbal medicines. J Pharm Anal 2011;01:235-45.
- Song R, Xu L, Xu F, Li Z, Dong H, Tian Y, et al. *In vivo* metabolism study of rhubarb decoction in rat using high-performance liquid chromatography with UV photodiode-array and mass-spectrometric detection: A strategy for systematic analysis of metabolites from traditional Chinese medicines in biological samples. J Chromatogr A 2010;1217:7144-52.
- Chi YC, Lin SP, Hou YC. A new herb-drug interaction of *Polygonum cuspidatum*, a resveratrol-rich nutraceutical, with carbamazepine in rats. Toxicol Appl Pharmacol 2012;263:315-22.
- Fan P, Marston A, Hay AE, Hostettmann K. Rapid separation of three glucosylated resveratrol analogues from the invasive plant *Polygonum cuspidatum* by high-speed countercurrent chromatography. J Sep Sci 2009;32:2979-84.
- Dong J, Wang H, Wan LR, Hashi Y, Chen SZ. Identification and determination of major constituents in *Polygonum cuspidatum* Sieb. et Zucc. by high performance liquid chromatography/electrospray ionization-ion trap time-of-flight mass spectrometry. Chin J Chromatogr 2009;27:425-30.
- Hegde VR, Pu H, Patel M, Black T, Soriano A, Zhao W, et al. Two new bacterial DNA primase inhibitors from the plant *Polygonum cuspidatum*. Bioorg Med Chem Lett 2004;14:2275-7.
- Xiao K, Xuan L, Xu Y, Bai D. Stilbene glycoside sulfates from *Polygonum cuspidatum*. J Nat Prod 2000;63:1373-6.
- Peng W, Qin R, Li X, Zhou H. Botany, phytochemistry, pharmacology, and potential application of *Polygonum cuspidatum* Sieb. et Zucc.: A review. J Ethnopharmacol 2013;148:729-45.
- Kimura Y, Kozawa M, Baba K, Hata K. New constituents of roots of *Polygonum cuspidatum*. Planta Med 1983;48:164-8.
- Xiao K, Xuan LJ, Xu YM, Bai DL. Studies on water-soluble constituents in rhizome of *Polygonum cuspidatum*. Chin Tradit Herb Drugs 2003;6:496-8.
- Qiu X, Zhang J, Huang Z, Zhu D, Xu W. Profiling of phenolic constituents in *Polygonum multiflorum* Thunb. by combination of ultra-high-pressure liquid chromatography with linear ion trap-Orbitrap mass spectrometry. J Chromatogr A 2013;1292:121-31.
- Hua Y, Zhou JY, Ni W, Chen CX. Studies on constituents of *Polygonum cuspidatum*. Nat Prod Res Dev 2001;13:16-8.
- State Administration of Traditional Chinese Medicine. Chinese Material Medica. Vol. 6. Shanghai: Science and Technology Press of Shanghai; 1999. p. 653-9.
- Liu XQ, Yu LM, Wu LJ. Chemical constituents of *Polygonum cuspidatum*. Chin J Chin Mater Med 2003;28:47-9.
- Vechar A, Kuznyetsova Z, Chekalinskaya I. Study of phenolcarboxylic acids of *Polygonum* species. Vol. 6. Proceedings of the Academy of Sciences of the Belorussian SSR, Biological Sciences Series; 1980. p. 71-4.
- Liu X, Wu L, Song G. Study on chemical constituents of *Polygonum cuspidatum* Sieb. J Shenyang Pharm Univ 1999;16:17-20.

**Cite this article as:** Fu J, Wang M, Guo H, Tian Y, Zhang Z, Song R. Profiling of components of rhizoma et radix polygoni cuspidati by high-performance liquid chromatography with ultraviolet diode-array detector and ion trap/time-of-flight mass spectrometric detection. Phcog Mag 2015;11:486-501.

**Source of Support:** Natural Science Foundation of China (No. 81403314), Natural Research Foundation of Jiangsu Province (No. BK2012349), Fundamental Research Funds for the Central Universities (JKPZ2013008) and the Open Project Program of MOE Key Laboratory of Drug Quality Control and Pharmacovigilance (No. MKLDP2013MS06).

**Conflict of Interest:** None declared.