

# Assessment of *In vitro* Antibacterial Activity and Cytotoxicity Effect of *Nigella sativa* Oil

Ayşe Ruveyda Ugur, Hatice Turk Dagi<sup>1</sup>, Bahadır Ozturk<sup>2</sup>, Gulsum Tekin<sup>2</sup>, Duygu Findik<sup>1</sup>

Department of Medical Microbiology, Konya Education and Research Hospital, Departments of <sup>1</sup>Medical Microbiology and <sup>2</sup>Biochemistry, Faculty of Medicine, Selcuk University, Konya, Turkey

Submitted: 12-10-2015

Revised: 30-11-2015

Published: 30-09-2016

## ABSTRACT

**Background:** Methicillin resistance is a serious health concern since it has spread among *Staphylococcus aureus* and coagulase-negative *Staphylococci* (CoNS) that are frequent community and nosocomial pathogens worldwide. Methicillin-resistant strains are often resistant to other classes of antibiotics, making their treatment difficult. *Nigella sativa* oil is known to be active against Gram-positive cocci, yet its *in vitro* cytotoxicity is rarely investigated, is a proper and powerful candidate for treatment of methicillin-resistant isolates. **Objectives:** The aim of this study is to evaluate the *in vitro* antibacterial activity and cytotoxicity effect of *N. sativa* oil. **Materials and Methods:** The minimal inhibitory concentrations (MICs) of *N. sativa* oil were determined by broth microdilution method against four different American Type Culture Collection strains, 45 clinical isolates of methicillin-resistant *S. aureus* (MRSA), and 77 methicillin-resistant CoNS (MRCoNS). The effects of different dilutions (0.25 µg/mL, 0.5 µg/mL, and 1 µg/mL) of *N. sativa* oil on the proliferation of gingival fibroblasts were evaluated. **Results:** The MIC values of *N. sativa* oil against clinical isolates of *Staphylococci* were between <0.25 µg/mL and 1.0 µg/mL. Compared to the control group, there was no cytotoxic effect on the proliferation of the gingival fibroblasts. **Conclusion:** In the present study, the oil of *N. sativa* was very active against MRSA and MRCoNS and had no *in vitro* cytotoxicity at relevant concentrations. These findings emphasize that there is a requirement for further clinical trials on *N. sativa* oil for "safe" medical management of infections caused by methicillin-resistant *Staphylococci*.


**Key words:** Cytotoxic effect, methicillin-resistant *Staphylococci*, microdilution, *Nigella sativa*

## SUMMARY

- The minimal inhibitory concentration (MIC) values of *Nigella sativa* oil against *Staphylococcus aureus* American Type Culture Collection (ATCC) 29213, *Enterococcus faecalis* ATCC 29212, *Escherichia coli* ATCC 25922, and *Pseudomonas aeruginosa* ATCC 27853 standard strains were 0.5 µg/mL,

2 µg/mL, 64 µg/mL, and 64 µg/mL, respectively

- The *N. sativa* oil showed an excellent antibacterial activity against clinical isolates of methicillin-resistant *S. aureus* and methicillin-resistant coagulase-negative *Staphylococci* with very low MIC range of <0.25–1.0 µg/mL
- The *N. sativa* oil exhibited no cytotoxic effect on the proliferation of the gingival fibroblasts.



**Abbreviation used:** ATCC: American Type Culture Collection; CLSI: Clinical and Laboratory Standards Institute; CoNS: Coagulase-negative Staphylococci; DMEM: Dulbecco's modified Eagle's medium; DMSO: Dimethylsulfoxide; FBS: Fetal bovine serum; HGF: Human gingival fibroblast; MIC: Minimal inhibitory concentration; MRCoNS: Methicillin-resistant CoNS; MRSA: Methicillin-resistant *S. aureus*

## Correspondence:

Dr. Ayşe Ruveyda Ugur,  
Department of Medical Microbiology,  
Konya Education and Research Hospital,  
Meram Yeniyol Caddesi No. 97, 42090 Meram,  
Konya, Turkey.  
E-mail: ayserugur@gmail.com  
**DOI:** 10.4103/0973-1296.191459

Access this article online

Website: www.phcog.com

Quick Response Code:



## INTRODUCTION

Methicillin resistance is a serious health concern since it has spread among *Staphylococcus aureus* and coagulase-negative *Staphylococci* (CoNS) that are frequent community and nosocomial pathogens worldwide.<sup>[1,2]</sup> Methicillin-resistant strains are often resistant to other classes of antibiotics making their treatment difficult.<sup>[2]</sup> In the recent years, the need for new antimicrobial agents because of the rise in antibiotic resistance has led to a search for alternative sources of antimicrobials.<sup>[3]</sup> Medicinal plants offer a wide range of biodiversity of great value for pharmacology. It has been known since antiquity that herbs and their essential oils have varying degrees of antimicrobial and therapeutic activity.<sup>[4]</sup> The World Health Organization has been recently supporting countries to integrate traditional medicine with their national health care systems.<sup>[5]</sup>

*Nigella sativa* that belongs to family *Ranunculaceae* is commonly known as black seed or black cumin.<sup>[6]</sup> It has been shown to possess

antimicrobial, immunomodulatory, anti-inflammatory, and antioxidant properties.<sup>[7]</sup> The antimicrobial activity of the oil and its constituents has been frequently studied so far.<sup>[8-10]</sup> Although there have been varying ranges of susceptibility results in the literature, Gram-positive bacteria such as *Bacillus cereus*, *S. aureus*, and *Staphylococcus epidermidis*

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

**Cite this article as:** Ugur AR, Dagi HT, Ozturk B, Tekin G, Findik D. Assessment of *In vitro* antibacterial activity and cytotoxicity effect of *Nigella sativa* oil. *Phcog Mag* 2016;12:471-4.

have been commonly designated as the most susceptible species to *N. sativa* oil.<sup>[8]</sup> Moreover, significant antimicrobial activity against multidrug-resistant clinical bacterial isolates has been also reported.<sup>[11-13]</sup> In this study, we aimed to investigate *in vitro* antibacterial activity and *in vitro* cytotoxicity effect of *N. sativa* oil, which has been so far merely studied.

## MATERIALS AND METHODS

### Antibacterial activity

The *in vitro* antibacterial activity of *N. sativa* oil was evaluated against following strains of *S. aureus* American Type Culture Collection (ATCC) 29213, *Enterococcus faecalis* ATCC 29212, *Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853, 45 clinical isolates of methicillin-resistant *S. aureus* (MRSA), and 77 clinical isolates of methicillin-resistant CoNS (MRCoNS).

The minimal inhibitory concentrations (MICs) of *N. sativa* oil were determined by broth microdilution method according to the Clinical and Laboratory Standards Institute (CLSI) guidelines.<sup>[14]</sup> 10.24 mg of cold pressed *N. sativa* oil (purchased from ZADEF Vital, Konya, Turkey) was dissolved in 10 mL of dimethyl sulfoxide (DMSO) to prepare a stock solution. The serial dilutions from the stock solution were made ranging from 256 µg/mL to 0.25 µg/mL using Mueller-Hinton broth (Becton Dickinson, Sparks, MD, USA) in 96-well microplates. The bacterial suspension containing approximately  $5 \times 10^5$  colony-forming units/mL was prepared from a 24 h culture plate. From this suspension, 100 µL was inoculated into each well. A sterility control well and a growth control well were also studied for each strain. The plates were incubated at 35°C for 24 h. The MICs were read as the lowest concentrations of *N. sativa* that inhibit the appearance of visible growth. These experiments were carried out in duplicate.

### Cell culture

The optimal seeding concentration (10,000 cells/well) for proliferation experiments of the human gingival fibroblasts (HGFs) was determined, and then, cells were allowed to adhere for 19 h in Dulbecco's modified Eagle's medium (DMEM) with 10% fetal bovine serum (FBS). The media were changed to DMEM with 5% FBS containing *N. sativa*. xCELLigence cell index (CI) impedance measurements were performed according to the instructions of the supplier.

### Proliferation experiments

Cytotoxic effect of the selected dilutions of *N. sativa* oil (0.25 µg/mL, 0.5 µg/mL, and 1 µg/mL), equal to the MICs at which the *Staphylococci* were susceptible to, were evaluated on the proliferation of gingival fibroblast cells by a real-time cell analyzer (xCELLigence, Roche Diagnostics GmbH, Penzberg, Germany). The cells were suspended in DMEM with 10% FBS. Then, 200 µL of the suspensions was seeded into wells (10,000 cells/well) of the E-plate 16. The gingival fibroblast cells were observed every 15 min during 95 h. After seeding, cells were held to attach to the E-plate for 19 h; then, the cells were exposed to 100 µL of medium containing dilutions of the *N. sativa* oil.

## RESULTS

### Antibacterial activity

The MIC values of *N. sativa* oil against *S. aureus* ATCC 29213, *E. faecalis* ATCC 29212, *E. coli* ATCC 25922, and *P. aeruginosa* ATCC 27853 standard strains were 0.5 µg/mL, 2 µg/mL, 64 µg/mL, and 64 µg/mL, respectively.

The MIC values of *N. sativa* oil against clinical isolates of *Staphylococci* were between <0.25 µg/mL and 1.0 µg/mL. Of 45 MRSA strains, MICs of 41 isolates were <0.25 µg/mL, two were 0.5 µg/mL, and two were 1 µg/mL. Of 77 MRCoNS strains, MICs of 53 isolates were <0.25 µg/mL, 19 were 0.25 µg/mL, two were 0.5 µg/mL, and three were 1 µg/mL.

### Proliferation experiments


For the *N. sativa* oil applications, cytotoxic effect at concentrations up to 1 µg/mL was not observed on the gingival fibroblasts when compared to the control group. CI of all NS oil concentrations was not significantly different after the treatment [Figure 1].

## DISCUSSION

*N. sativa* seed oil consists of oleoresins and essential oil components, including thymoquinone, dithymoquinone, thymohydroquinone, p-cymene, carvacrol, 4-terpineol, α-thujene, t-anethol, longifolene, thymol, and pinene.<sup>[15]</sup> Thymoquinone (30–52.6%) and p-cymene (7–25.8%) were reported as its major components.<sup>[15,16]</sup> Antimicrobial activity of *N. sativa* oil is attributed mainly to its phenolic constituents of the essential oil compartment. Thus, thymoquinone followed by its related compounds such as thymohydroquinone, dithymoquinone, and thymol along with carvacrol plays major role in antimicrobial activity.<sup>[8,17,18]</sup> Other constituents, oleoresins, linoleic acid, and oleic acid, may also have minor antimicrobial activity.<sup>[15]</sup> Indeed, whole essential oil was reported to have higher antibacterial activity than the combinations of its prominent constituents, suggesting that the minor components potentiate the antimicrobial activity.<sup>[19-22]</sup>

Although our findings regarding the antibacterial activity of *N. sativa* oil against Gram-positive and Gram-negative bacteria were in agreement with other studies in terms of Gram-positive bacteria being more susceptible, our results were inconsistent with most of previously published works in terms of much lower MICs against Gram-positive ATCC strains and clinical isolates of *Staphylococci*. This discrepancy may be explained by several factors such as differences in the extraction methods, antibacterial assay methods used, percentage of active components in the oils, quality and composition of the active constituents, and type of microorganisms selected.<sup>[8,19,23,24]</sup>

In fact, the oil composition and antimicrobial activities of a specific plant may differ depending on geographical locations where it is cultivated and on harvesting periods.<sup>[24-27]</sup> It was reported that genetic differences between *N. sativa* seeds grown in the different countries exist and the genetic polymorphisms took place over time seem to cause distinct



**Figure 1:** Dynamic monitoring of cell adhesion and proliferation using the xCELLigence system. The effects of different dilutions of *Nigella sativa* oil on the proliferation of human gingival fibroblasts at a density of 10,000 cells per well in E-Plates 96 were observed during 95 h

varieties.<sup>[28]</sup> Moreover, differences in the antimicrobial activities of the essential oils may be obtained depending on the species, subspecies, or varieties.<sup>[25]</sup> Indeed, comparison of data belonging to previously published works becomes more complicated because of the lack of a standardized method for investigating the antimicrobial activity of natural compounds such as oils obtained from various herbs. Yet, disk diffusion and broth microdilution methods are most applied techniques to investigate the antimicrobial activity of plant and seed oils.<sup>[8,15,17,29]</sup> The solvents utilized to dissolve these oils or their constituents are also quite diverse.<sup>[8,16,29,30]</sup> CLSI recommends DMSO, ethyl alcohol, polyethylene glycol, and carboxymethyl cellulose as solvents for water-insoluble drugs without mentioning naturally existing antimicrobial compounds.<sup>[14]</sup> The units used for the MIC values that are reported as mg/mL, µg/mL, µL/mL, ppm, µL/well, and %(v/v) also differ between articles making comparison of results very difficult.<sup>[16,19]</sup> Considering vast studies in the literature, it is obvious that there is a need to standardize every step of antimicrobial susceptibility tests for essential oils and their components. However, this subject is beyond our scope.

In the present study, we investigated the antibacterial activity of the whole *N. sativa* oil by broth microdilution method. Our findings presented at least three-fold lower MIC values against *Staphylococci* when compared to previously published works, most of which have studied active components instead of the entire oil of the *N. sativa* seed.<sup>[12,17,30-32]</sup> It was previously reported that the MIC value of 12.5 µg/mL of the *N. sativa* seed extract was the lowest concentration at which all the tested microorganisms (*E. coli*, *B. subtilis*, *S. aureus*, *P. aeruginosa*, *Candida albicans*, *Aspergillus niger*) were inhibited suggesting that further dilutions may possess antimicrobial activity against *Staphylococci*.<sup>[29]</sup> In point of fact, it has been severally shown that *Staphylococci* are more susceptible to *N. sativa* oil and its components than other bacteria.<sup>[8,32,33]</sup>

In the present study, an excellent activity of the *N. sativa* oil was observed with very low MICs against clinical isolates of MRSA and MRCoNS. Our findings were in agreement with literature reporting thymoquinone, the most active constituent, to have substantial antimicrobial activity against MRSA.<sup>[34,35]</sup> Although Hannan *et al.*<sup>[12]</sup> reported higher MIC ranges (0.2–0.5 mg/mL) against MRSA, their results indicated that *N. sativa* has inhibitory effect on MRSA. In another study, the multidrug-resistant *S. aureus* isolates from diabetic wounds were susceptible to various concentrations of *N. sativa* oil.<sup>[11]</sup> Some experiments on animals show that *N. sativa* oil has significant *in vivo* antibacterial activity on *S. aureus* infections.<sup>[9,36]</sup> It is also noteworthy to mention that MICs against methicillin susceptible and methicillin-resistant *Staphylococci* did not differ significantly in our study ( $P > 0.05$ ) as in several other works.<sup>[17,30]</sup> Moreover, a few studies demonstrated that *N. sativa* essential oil and thymoquinone can effectively inhibit *S. aureus* biofilm formation, suggesting that *N. sativa* oil deserves further investigations on methicillin-resistant *Staphylococci*.<sup>[31,37]</sup>

The seed extracts of *N. sativa* are characterized by a low level of toxicity. Although potential toxicity of the *N. sativa* seed oil was investigated in animal experiments to determine LD<sub>50</sub> values, its *in vitro* cytotoxicity effect has been rarely studied. It was reported that the extract of *N. sativa* seed was not toxic when administered to rats intraperitoneally at a daily dose of 50 mg/kg.<sup>[38]</sup> In addition, experimental animals were not affected when *N. sativa* oil at doses of 10 mL/kg was administered orally.<sup>[39,40]</sup> In the present study, *in vitro* cytotoxicity assay of *N. sativa* oil on the fibroblast cells did not represent cytotoxic effect at the relevant dilutions when compared to the control group. Kadan *et al.*<sup>[41]</sup> reported that cytotoxic effect of 50% ethanol/water extract of *N. sativa* on the human hepatocellular carcinoma and the rat L6 muscle cell line exhibited at concentrations higher than 500 µg/mL. Although this concentration was much higher than concentrations we investigated, the results should not

be compared with each other because the subject materials are different in composition and nature.

## CONCLUSION

The oil of *N. sativa* was very active against MRSA and MRCoNS and had no *in vitro* cytotoxicity at concentrations up to 1 µg/mL in the present study. These findings emphasize that there is a requirement for further clinical trials on *N. sativa* oil for “safe” medical management of infections caused by methicillin-resistant *Staphylococci*.

## Acknowledgement

HGFs used in this study have been provided by Dr. Sema Hakkı (Selcuk University Faculty of Dentistry, Konya, Turkey).

The authors declare that they have no competing interests.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Dhawan B, Rao C, Udo EE, Gadepalli R, Vishnubhatla S, Kapil A. Dissemination of methicillin-resistant *Staphylococcus aureus* SCCmec type IV and SCCmec type V epidemic clones in a tertiary hospital: Challenge to infection control. *Epidemiol Infect* 2015;143:343-53.
- Murugesan S, Perumal N, Mahalingam SP, Dilliappan SK, Krishnan P. Analysis of antibiotic resistance genes and its associated SCCmec types among nasal carriage of methicillin resistant coagulase negative *Staphylococci* from community settings, Chennai, Southern India. *J Clin Diagn Res* 2015;9:DC01-5.
- Carson CF, Riley TV. Non-antibiotic therapies for infectious diseases. *Commun Dis Intell Q Rep* 2003;27 (Suppl): S143-6.
- Singh G, Marimuthu P, Murali HS, Bawa AS. Antioxidative and antibacterial potentials of essential oils and extracts isolated from various spice materials. *J Food Saf* 2005;25:130-45.
- [Homepage on the Internet]. WHO Traditional Medicine Strategy 2014-2023. Hong Kong SAR, China: Geneva, Switzerland; December 2013; Available from: [http://www.apps.who.int/iris/bitstream/10665/92455/1/9789241506090\\_eng.pdf?ua=1](http://www.apps.who.int/iris/bitstream/10665/92455/1/9789241506090_eng.pdf?ua=1). [Last cited on 2015 Jan 2].
- Paarakh PM. *Nigella sativa* Linn. A comprehensive review. *Indian J Nat Prod Resour* 2010;1:409-29.
- Ali BH, Blunden G. Pharmacological and toxicological properties of *Nigella sativa*. *Phytother Res* 2003;17:299-305.
- Kokoska L, Havlik J, Valterova I, Sovova H, Sajfrtova M, Jankovska I. Comparison of chemical composition and antibacterial activity of *Nigella sativa* seed essential oils obtained by different extraction methods. *J Food Prot* 2008;71:2475-80.
- Hanafy MS, Hatem ME. Studies on the antimicrobial activity of *Nigella sativa* seed (black cummin). *J Ethnopharmacol* 1991;34:275-8.
- Topozada HH, Mazloum HA, el-Dakhkhny M. The antibacterial properties of the *Nigella sativa* L. seeds. Active principle with some clinical applications. *J Egypt Med Assoc* 1965;48:187-202.
- Emeka LB, Emeka PM, Khan TM. Antimicrobial activity of *Nigella sativa* L. seed oil against multi-drug resistant *Staphylococcus aureus* isolated from diabetic wounds. *Pak J Pharm Sci* 2015;28:1985-90.
- Hannan A, Saleem S, Chaudhary S, Barkaat M, Arshad MU. Anti bacterial activity of *Nigella sativa* against clinical isolates of methicillin resistant *Staphylococcus aureus*. *J Ayub Med Coll Abbottabad* 2008;20:72-4.
- Islam MH, Ahmad IZ, Salman MT. Antibacterial activity of *Nigella sativa* seed in various germination phases on clinical bacterial strains isolated from human patients. *JBPR* 2012;4:8-13.
- Clinical and Laboratory Standard Institute. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-First Informational Supplement, CLSI Document M100-S21. Wayne: Clinical and Laboratory Standard Institute; 2011.
- Singh S, Das SS, Singh G, Schuff C, de Lampasona MP, Catalán CA. Composition, *in vitro*



- antioxidant and antimicrobial activities of essential oil and oleoresins obtained from black cumin seeds (*Nigella sativa* L.). *Biomed Res Int* 2014;2014:918209.
16. Shaaban HA, Sadek Z, Edris AE, Saad-Hussein A. Analysis and antibacterial activity of *Nigella sativa* essential oil formulated in microemulsion system. *J Oleo Sci* 2015;64:223-32.
  17. Halawani E. Antibacterial activity of thymoquinone and thymohydroquinone of *Nigella sativa* L. and their interaction with some antibiotics. *Adv Biol Res* 2009;3:148-52.
  18. Ghosheh OA, Houdi AA, Crooks PA. High performance liquid chromatographic analysis of the pharmacologically active quinones and related compounds in the oil of the black seed (*Nigella sativa* L.). *J Pharm Biomed Anal* 1999;19:757-62.
  19. Burt S. Essential oils: Their antibacterial properties and potential applications in foods – A review. *Int J Food Microbiol* 2004;94:223-53.
  20. Bassolé IH, Juliani HR. Essential oils in combination and their antimicrobial properties. *Molecules* 2012;17:3989-4006.
  21. Mourey A, Canillac N. Anti-histria monocytogenes activity of essential oils components of conifers. *Food Control* 2002;13:289-92.
  22. Anwar S, Ahmed N, Habibatni S, Abusamra Y. Ajwain (*Trachyspermum Ammi* L.) oils. In: Preed VR, editor. *Essential Oils in Food Preservation, Flavor and Safety*. London: Academic Press; 2015. p. 181-92.
  23. Mith H, Duré R, Delcenserie V, Zhiri A, Daube G, Clinquart A. Antimicrobial activities of commercial essential oils and their components against food-borne pathogens and food spoilage bacteria. *Food Sci Nutr* 2014;2:403-16.
  24. Karakaya S, El SN, Karagözü N, Sahin S. Antioxidant and antimicrobial activities of essential oils obtained from oregano (*Origanum vulgare* ssp. *hirtum*) by using different extraction methods. *J Med Food* 2011;14:645-52.
  25. Sarac N, Ugur A. Antimicrobial activities of the essential oils of *Origanum onites* L. *Origanum vulgare* L. subspecies *hirtum* (Link) letswaart, *Satureja thymbra* L. and *Thymus cilicicus* Boiss. and Bal. growing wild in Turkey. *J Med Food* 2008;11:568-73.
  26. McGimpsey JA, Douglas MH, Van Klink JL, Beauregard DA, Perry NB. Seasonal variation in essential oil yield and composition from naturalized *Thymus vulgaris* L. in New Zealand. *Flavour Fragr J* 1994;9:347-52.
  27. D'Antuono LF, Moretti A, Lovato AF. Seed yield, yield components, oil content and essential oil content and composition of *Nigella sativa* L. and *Nigella damascena* L. *Ind Crops Prod* 2002;15:59-69.
  28. Al-Huqail A, Al-Saad F. DNA fingerprinting and genotyping of four black seed (*Nigella sativa* L.) Taxa. *J King Abdul Univ Meteorol Environ Arid Land Agric Sci* 2010;21:93-108.
  29. Bakathir HA, Abbas NA. Detection of the antibacterial effect of *Nigella sativa* ground seeds with water. *Afr J Tradit Complement Altern Med* 2011;8:159-64.
  30. Dadgar T, Asmar M, Saifi A, Mazandarani M, Bayat H, Moradi A, *et al.* Antibacterial activity of certain Iranian medicinal plants against methicillin-resistant and sensitive *Staphylococcus aureus*. *Asian J Plant Sci* 2006;5:861-6.
  31. Chaieb K, Kouidhi B, Jrah H, Mahdouani K, Bakhrouf A. Antibacterial activity of Thymoquinone, an active principle of *Nigella sativa* and its potency to prevent bacterial biofilm formation. *BMC Complement Altern Med* 2011;11:29.
  32. Toama MA, El-Alfy TS, El-Fatratry HM. Antimicrobial activity of the volatile oil of *Nigella sativa* Linneaus seeds. *Antimicrob Agents Chemother* 1974;6:225-6.
  33. Kabbashi AS, Garbi MI, Osman EE, Dahab MM, Koko WS, Abuzeid N. *In vitro* antimicrobial activity of ethanolic seeds extracts of *Nigella sativa* (Linn) in the Sudan. *Afr J Microbiol Res* 2015;9:788-92.
  34. Liu M, Koya S, Furuta H, Matsuzaki S. Growth-inhibiting activity of anthraquinones and benzoquinones against methicillin resistant *Staphylococcus aureus* (MRSA). *Dokkyo J Med Sci* 1996;23:85-93.
  35. Salman MT, Khan RA, Shukla I. Antimicrobial activity of *Nigella sativa* Linn. seed oil against multi-drug resistant bacteria from clinical isolates. *NPR* 2008;7:10-4.
  36. Hosseinzadeh H, Bazzaz BSF, Haghi MM. Antibacterial activity of total extracts and essential oil of *Nigella Sativa* L. seeds in mice. *Pharmacologyonline* 2007;2:429-35.
  37. Manju S, Malaikozhundan B, Vijayakumar S, Shanthi S, Jaishabanu A, Ekambaram P, *et al.* Antibacterial, antibiofilm and cytotoxic effects of *Nigella sativa* essential oil coated gold nanoparticles. *Microb Pathog* 2016;91:129-35.
  38. el Daly ES. Protective effect of cysteine and Vitamin E, *Crocus sativus* and *Nigella sativa* extracts on cisplatin-induced toxicity in rats. *J Pharm Belg* 1998;53:87-93.
  39. Khanna T, Zaidi FA, Dandiya PC. CNS and analgesic studies on *Nigella sativa*. *Fitoterapia* 1993;5:407-10.
  40. Zaoui A, Cherrah Y, Mahassini N, Alaoui K, Amarouch H, Hassar M. Acute and chronic toxicity of *Nigella sativa* fixed oil. *Phytomedicine* 2002;9:69-74.
  41. Kadan S, Saad B, Sasson Y, Zaid H. *In vitro* evaluations of cytotoxicity of eight antidiabetic medicinal plants and their effect on GLUT4 Translocation. *Evid Based Complement Alternat Med* 2013;2013:549345.



Ayşe Ruveyda Ugur

## ABOUT AUTHOR

**Dr. Ayşe Ruveyda Ugur**, is a medical microbiologist at Konya Education and Research Hospital. She completed her medical education (2009) and her specialist training in medical microbiology (2015) at Selçuk University Faculty of Medicine. She has experience and proficiency in medical laboratory management, including diagnostic methods, antimicrobial susceptibility testing, antimicrobial policies, and hospital policies for infection control.