

# Free Radical Scavenging, $\alpha$ -Glucosidase Inhibitory and Anti-Inflammatory Constituents from Indian Sedges, *Cyperus scariosus* R.Br and *Cyperus rotundus* L.

Lavanya Kakarla, Suresh Babu Katragadda<sup>1</sup>, Ashok K Tiwari<sup>2</sup>, K Srigiridhar Kotamraju<sup>3</sup>, K Madhusudana<sup>2</sup>, Anand Kumar D<sup>2</sup>, Mahendran Botlagunta

Biomedical Research Laboratory, Department of Biotechnology, K L E F University, Vaddeswaram, Guntur, Andhra Pradesh, <sup>1</sup>Division of Natural Products Chemistry, CSIR-Indian Institute of Chemical Technology, Uppal Road, Hyderabad, <sup>2</sup>Medicinal Chemistry and Pharmacology Division, CSIR-Indian Institute of Chemical Technology, Uppal Road, Hyderabad, <sup>3</sup>Centre for Chemical Biology, CSIR-Indian Institute of Chemical Technology, Uppal Road, Hyderabad, India

Submitted: 18-12-2015

Revised: 02-02-2016

Published: 30-09-2016

## ABSTRACT


**Background:** *Cyperus scariosus* R. Br and *Cyperus rotundus* L are widely used in ayurvedic preparation for the treatment of diabetes and other diseases. The early literature, so far, does not indicate the presence of any bioactive principle isolated from these plants. **Objective:** To identify free radical scavenging, anti-diabetic and anti-inflammatory principles from these two species. **Materials and Methods:** The bioassay guided fractionation and isolation of active constituents was done by chromatographic techniques. They also evaluated their anti-oxidant activity by DPPH and ABTS. The anti-diabetic activity was screened by  $\alpha$ -glucosidase and  $\alpha$ -amylase assays. Also, the further evaluation of *in vitro* anti-inflammatory activity using THP-1 monocytic cells and *in vivo* anti-inflammatory activity, was confirmed by carrageenan induced rat paw edema as model. **Results:** The activity guided isolation led to isolation of twelve compounds which are: Stigmasterol<sup>[1]</sup>,  $\beta$ -sitosterol<sup>[2]</sup>, Lupeol<sup>[3]</sup>, Gallic acid<sup>[4]</sup>, Quercetin<sup>[5]</sup>,  $\beta$ -amyrin<sup>[6]</sup>, Oleanolic acid<sup>[7]</sup>,  $\beta$ -amyrin acetate<sup>[8]</sup>, 4-hydroxybutyl cinnamate<sup>[9]</sup>, 4-hydroxy cinnamic acid<sup>[10]</sup>, Caffeic acid<sup>[11]</sup> and Kaempferol<sup>[12]</sup> respectively. Among the isolates, the compounds 4 and 5 displayed potent radical scavenging activity with an  $IC_{50}$  values of 0.43 and 0.067  $\mu$ g/ml. The compounds 4, 5 and 10 showed significant anti-diabetic activities. While lupeol<sup>[3]</sup> showed potent IL-1 $\beta$  activity inhibition in THP-1 monocytic cells and also displayed significant ( $p < 0.0025$ ) *in vivo* anti-inflammatory activity. **Conclusion:** In brief, we isolated twelve compounds from both the species. Collectively, our results suggested that aromatic compounds showed good anti-oxidant and anti-diabetic activities.

**Key words:** Alpha amylase, alpha glucosidase, antioxidant activity, IL-1 $\beta$  inflammation, *Cyperus scariosus* and *Cyperus rotundus*.

## SUMMARY

- The study investigates the free radical scavenging,  $\alpha$ -glucosidase inhibitory and anti-inflammatory effects of constituents isolated from Indian sedges viz. *C. scariosus* and *C. rotundus*. The results indicated that phenolic compounds

displayed potent free radical scavenging activity and alpha-glucosidase inhibition activity. While terpene constituent, Lupeol<sup>[3]</sup> showed good IL-1 $\beta$  activity inhibition in THP-1 monocytic cells and also displayed significant ( $p < 0.0025$ ) *in vivo* anti-inflammatory activity in carrageenan induced rat paw edema. However, further studies are required to know the exact molecular mechanism.



**Abbreviations used:** DPPH: 2,2-Diphenyl-1-picrylhydrazyl, ABTS: 2,2-Azinobis(3-ethylbenzothiazoline-6-sulfonic acid), THP-1: Human leukaemia monocytic cell line, IL-1 $\beta$ : Interleukin-1 $\beta$ ,  $IC_{50}$  - Inhibitory concentration 50%.

## Correspondence:

Dr. Mahendran Botlagunta,  
Associate professor,  
Biomedical Research Laboratory,  
Department of Biotechnology,  
K L E F (Koneru Lakshma Education  
Foundation) University, Vaddeswaram,  
Guntur, Andhra Pradesh, India.  
E-mail: bmnchowdary@gmail.com  
DOI: 10.4103/0973-1296.191467

Access this article online

Website: www.phcog.com

Quick Response Code:



## INTRODUCTION

Human body has an antioxidant defence system towards free radical, but over production of free radicals cause imbalance condition between the free radical and antioxidant defence in the body that leads to several diseases including inflammation, aging and diabetes.<sup>[1]</sup> In the case of diabetes,  $\alpha$ -glucosidase is an enzyme that catalyzes the decomposition of disaccharides to glucose for the absorption into the blood stream, in small intestine. Therefore,  $\alpha$ -glucosidase inhibitors could regulate abnormally high stages of plasma glucose after carbohydrate ingestion.<sup>[2]</sup> Presently, synthetic  $\alpha$ -glucosidase inhibitors such as acarbose has been accepted for clinical use in the treatment of type-2 diabetes.<sup>[3]</sup> On the other hand, a number of biological agents have been developed to treat inflammation, including agents that reduce the activity of specific cytokines or their receptors, block lymphocyte trafficking into tissues,

prevent the binding of monocyte-lymphocyte to stimulatory molecules.<sup>[4]</sup> Many of the existing synthetic anti-diabetic and anti-inflammatory drugs can cause several side effects such as flatulence, diarrhoea, cardiovascular diseases, gastrointestinal, and urinary tract infections.<sup>[5,6]</sup>

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

**Cite this article as:** Kakarla L, Katragadda SB, Tiwari AK, Kotamraju KS, Madhusudana K, Kumar DA, Botlagunta M. Free radical scavenging,  $\alpha$ -glucosidase inhibitory and anti-inflammatory constituents from Indian sedges, *Cyperus scariosus* R.Br and *Cyperus rotundus* L.. Phcog Mag 2016;12:488-96.

In recent years, search for novel therapeutic agents and approaches for diabetes and inflammation is still an active research field stimulated by the discovery of new biological targets, and by the possibility of obtaining new bioactive compounds with multiple biological activities. The past decade has witnessed an increasing interest in search of plant based lead compounds for the development of new pharmaceuticals.<sup>[7]</sup> In this regard, plants of the cyperaceae family which comprises a group of more than 5000 different species, distributed in tropical and subtropical areas of the world have attracted the scientific interest because of their wide range of biological activities such as anti-oxidant, antidiabetic, anticancer, antibacterial, antiviral, anti-inflammatory, anti-allergic, and vasodilatory actions.<sup>[7-13]</sup> Among the plants of Cyperaceae family, *Cyperus scariosus* R. Br and *Cyperus rotundus* L, are prominent species which are extensively used in the traditional systems of medicine for the treatment of various ailments. Herbal mixtures of both the species are used to enhance microbial protein synthesis in the rumen of buffaloes. These mixtures are anti-cancer, anti-diabetic, anti-inflammatory activity, suppresses formation and prevent wrinkles, protein oxidation, antinociceptive and, anti-hyperglycemic activities.<sup>[14-17]</sup> In spite of the various pharmacological uses of *C. scariosus* and *C. rotundus* extracts, little is known about their chemical constituents. Previous studies on this species have resulted in the isolation of sesquiterpenes, flavonoids, triterpenoids, and steroids.<sup>[18-20]</sup> Our continuing efforts are directed towards identification of bio active compounds from the Indian medicinal flora. Recently, we reported morphological and chemoprofiling comparisons of various solvent extracts of these two species *C. scariosus* and *C. rotundus* using analytical techniques.<sup>[21]</sup> Consequently, in line with this interest, the current study reports on the presence of free radical scavenging, alpha glucosidase inhibitory and anti-inflammatory activity constituents. These results are reported for the first time in the crude extracts of Indian sedges. Based on these results, we also discussed structure activity relationship of the isolates.

## MATERIALS AND METHODS

### General procedure

The solvents used were all of analytical grade and purchased from local distributor. The thin layer chromatography was performed on precoated silica gel glass plate 60 F<sub>254</sub>. Visualization was performed using chemical spray reagent solution, followed by charring on hotplate. Column chromatography was carried out using silicagel 60-120, 100-200 and 230-400 mesh (qingdao Marine chemical, China). Melting points were recorded on a fisher scientific melting point apparatus. FT-IR spectra was recorded on Thermo Nicolet Nexus 670 FT-IR spectrometer (Thermo fisher). <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were recorded on Bruker spectrometer. ESI-MS was measured on LC-MS-Trap-SL- instrument. DPPH, ABTS, Piroxicam, phorbol-12-Myristate-13- acetate (PMA) were purchased from sigma aldrich.

### Plant material collection, identification and extraction

The whole plants of *C. scariosus* were collected from the Sri Satyadeva nursery, Kadiyam, East Godavari district, Andhrapradesh and *C. rotundus* plants were collected from rice fields. These plants were taxonomically identified by Dr. A. Prasada Rao, Taxonomist, KLU university, Vaddeswaram, Guntur, Andhra Pradesh, India. The voucher specimens were deposited at KLU university, botanical garden, voucher specimen number (KLU-1250 and KLU-1251) for further use. The rhizomes of these herbs were separated, washed, shade dried and grounded to fine powder. The air dried and powdered rhizomes of *C. scariosus* (5 kg) and *C. rotundus* (5 kg), were sequentially extracted with hexane, chloroform, and methanol by cold maceration for 72 hours.

Following extraction, the solvents from each extract were concentrated under reduced pressure to the dryness in a rotary vacuum evaporator.

### Phytochemical chemical investigation of rhizomes of *C. scariosus*

The crude hexane extract of *C. scariosus* (10 gm) was subjected to silica gel column chromatography and eluted successively with hexane: ethyl acetate (100:0, 90:10, 80:20, 70:30, 60:40 and 50:50) mixture in the increasing order of polarity. The column eluted fractions were collected and concentrated. The purity of eluted fractions were analyzed by TLC (solvent system hexane: EtOAc 90:10) and fractions with similar R<sub>f</sub> value were combined to get six major fractions (Viz: F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, F<sub>4</sub>, F<sub>5</sub> and F<sub>6</sub>). The repeated purification of resultant fractions led to the isolation of two compounds [1 and 2]. Similarly, 10 gm's chloroform extract of *C. scariosus* was subjected to column chromatography on a silica gel column, and eluted with a step wise gradient of chloroform: acetone. Initially, the column was eluted with plain chloroform. Then, the column was eluted with chloroform: acetone, (100:0, 90:10, 80:20, 70:30, 60:40 and 50:50) mixture in the increasing order of polarity. The collected fractions purity were analyzed by TLC (mobile phase chloroform: acetone 70:30). These fractions with similar TLC patterns were combined to give six major fractions (Viz: F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, F<sub>4</sub>, F<sub>5</sub> and F<sub>6</sub>), and the repeated purification of resultant fractions led to the isolation of compound 3 in addition to the compounds 1 and 2. Likewise, the methanol extract (10 gm) of *C. scariosus* was loaded on a silica gel column and elution was carried out with a step wise gradient of chloroform: methanol. Initially the column was eluted with plain chloroform, then it was eluted with chloroform: methanol, (100:0, 98:02, 96:04, 95:05 and 90:10) mixture in the increasing order of polarity. All these column fractions were analyzed by TLC (silica gel 60 F<sub>254</sub>, chloroform: methanol 90:10) and fractions with similar TLC patterns were combined to give five major fractions (viz: F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, F<sub>4</sub> and F<sub>5</sub>). The extensive chromatographic purification of resultant fractions led to the isolation of two compounds. [4 and 5]

### Phytochemical chemical investigation of rhizomes of *C. rotundus*

The crude hexane extract of *C. rotundus* (10 gm) was loaded on silica gel column and eluted successively with hexane: ethyl acetate (100:0, 98:02, 96:04, 95:05, 94:06 and 90:10) mixture in the increasing order of polarity. All these column fractions were analyzed by TLC (silica gel 60 F<sub>254</sub>, hexane: EtOAc (90:10) and fractions with similar TLC patterns were combined to give six major fractions (viz: F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, F<sub>4</sub>, F<sub>5</sub> and F<sub>6</sub>). The repeated purification of resultant fractions led to the isolation of three compounds 6, 7 and 8 in addition to the compounds (1 and 2). Similarly, the crude chloroform extract (10 gm) of *C. rotundus* elution was carried out with a step wise gradient of hexane: acetone. Initially, the column was eluted with plain hexane, then it was eluted with hexane: acetone, (100:0, 98:02, 96:04, 95:05 and 90:10) mixture in the increasing order of polarity. The collected fractions banding pattern were analyzed by TLC (mobile phase hexane: acetone 90:10). These fractions with similar TLC patterns were combined to give four major fractions (viz: F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub> and F<sub>4</sub>), which yielded compounds 9 and 10 after repeated purifications. Later, methanol extract (10 gm) of *C. rotundus* was carried out on a silica gel column with a step wise gradient elution of chloroform: methanol. Initially the column was eluted with plain chloroform. The column was, then, eluted with chloroform: methanol, (100:0, 98:02, 96:04, 95:05 and 90:10) mixture in the increasing order of polarity. The column fractions were analyzed by TLC (silica gel 60 F<sub>254</sub>, chloroform: methanol 90:10) and fractions with similar TLC patterns were combined to give five major fractions (Viz: F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, F<sub>4</sub> and F<sub>5</sub>). The repeated purifications of resultant fractions led to the isolation of compounds. [11 and 12]

## Determination of free radicals scavenging antioxidant potentials

### DPPH radical scavenging activity

The scavenging of 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical was determined as per the standard protocols.<sup>[22]</sup> Briefly, in a 96 - well micro plate reaction, mixture was prepared with 25  $\mu$ l compound (1 mg/ml DMSO), 100  $\mu$ l of 0.1M Tris - HCl buffer (pH7.4), 125  $\mu$ l DPPH solutions (0.5 mM in methanol) and incubated in dark for 30 minutes. DPPH decolourization was recorded spectrophotometrically (BioTek synergy4 multi-mode micro plate reader, BioTek Instruments, Inc Winooski, VT, USA) at 517 nm. The Percentages of DPPH scavenging and IC<sub>50</sub> values were calculated.

% inhibition = [(Absorbance control-Absorbance test)/Absorbance control] x 100.

[IC<sub>50</sub>] The values were calculated using linear regression analysis. The IC<sub>50</sub> values signify the concentration of sample, which is appropriate in scavenging 50% of the DPPH free radicals.

### ABTS radical scavenging activity

The ABTS free radical scavenging activity was performed as per the standard protocols.<sup>[22]</sup> The 100 ml stock solution of ABTS (0.5 mM) was prepared by an addition of 1 ml potassium persulfate (6.8 mM PBS, pH 8.0) and stored in dark for 24 hrs. In a 96-well micro plate, 10  $\mu$ l of compound was mixed with 190  $\mu$ l of ABTS. After 15 mins of incubation in dark, decolorized ABTS absorbance was measured spectrophotometrically at 734 nm.

% of inhibition] = [(Absorbance control-Absorbance test)/Absorbance control] x 100.

### Intestinal $\alpha$ -glucosidase inhibition assay

The inhibition of rat intestinal  $\alpha$ -glucosidase enzyme was determined as per the standard protocols.<sup>[23]</sup> 20 $\mu$ l of compound (1 mg/ml DMSO) was incubated with 50  $\mu$ l of crude intestinal  $\alpha$ -glucosidase for 5 mins and then with 50  $\mu$ l of substrate 5 mM p-nitro phenyl- $\alpha$ -D-glucopyranoside and the absorbance was measured at 405 nm. The percentage of enzyme inhibition was calculated by applying the formula mentioned above.

### Pancreatic $\alpha$ -amylase inhibition assay

The inhibition of  $\alpha$ - amylase enzyme was determined as per the standard protocols.<sup>[23]</sup> 40  $\mu$ l of compound was reconstituted in 100  $\mu$ l phosphate buffer (20 mM, p<sup>H</sup> 6.8), containing 6.7 mM sodium chloride in 2 ml eppendorf tubes, and incubated with 200  $\mu$ l porcine pancreatic  $\alpha$ -amylase for 10 minutes. Add 100  $\mu$ l of soluble potato starch solution (0.5%), and exactly after 10 minutes, incubation 400  $\mu$ l of DNS (Di nitro salicylic acid) color reagent was added. The closed tubes were placed in water bath (85-90°C) for 10 minutes to develop color and were cooled. Then, reaction mixture (50  $\mu$ l) was diluted with 175  $\mu$ l of distilled water in a 96 - well micro plate and absorbance (540 nm) was read spectrophotometrically. Percentage of inhibition was calculated by applying above formula.

## IN VITRO SCREENING OF ANTI-INFLAMMATORY ACTIVITY

### Cells and culture

Human acute monocytic leukaemia cell line (THP-1) was procured from National Centre of Cell Sciences (NCCS, Pune, India). The cell line was cultured in Roswell Park Memorial Institute Medium 1640 (RPMI, Sigma, USA). The medium was supplemented with 10% (v/v) heat inactivated fetal bovine serum (Sigma, USA), 1% 100 U/ml penicillin (Sigma, USA) and 1% 100 U/ml streptomycin (Sigma, USA) in 25-cm2

culture flask and incubated at 37°C in an atmosphere of 5% CO<sub>2</sub>. The THP-1 monocytes were treated with PMA (Phorbol- 12 - myristate-13-acetate) in 24 well plates. After 48 hr stimulation, medium was removed, washed twice with PBS and phase contrast images were taken by light microscopy to study the monocyte to macrophage adherence and transformation.

### Enzyme-linked immunosorbent assay for cytokine analysis (ELISA)

The THP-1 monocytes were seeded at a density of 2 x 10<sup>5</sup> cells/ ml in 24 well plates in triplicates. After 48 hrs of stimulation of cells with phorbol 12-myristate 13-acetate (PMA) in the presence or absence of compounds, supernatant was harvested from the cell culture plates into 1.5 ml eppendorf tubes and centrifuged to remove the cell debris. The cell free supernatants were transferred into fresh 1.5 ml eppendorf tubes and transferred them immediately into -80°C until use. On the next day of analysis, samples were carefully thawed and used for the analysis of IL-1 $\beta$  levels using Human Ready-SET-Go ELISA kit (eBiosciences, San Diego, CA, USA) following the manufacturer's instructions.<sup>[24]</sup> Briefly, 100  $\mu$ l of cell free supernatants of test sample, standard, and control was added to 1% BSA- blocked ELISA plate that had been coated overnight with 100  $\mu$ l/ well anti- murine IL-1 $\beta$ . Following sample addition, 100  $\mu$ l/well biotin conjugated IL-1 $\beta$  antibody was added, and then plates were incubated at room temperature for 3hrs. Following detecting antibody addition, 100  $\mu$ l/well (Horse reddish peroxidase) HRP-conjugated avidin was added and incubated at room temperature for 30-45 mins, then aspirated and washed 5-7 times with 250  $\mu$ l/ well of wash buffer with soaking for 5 mins during each wash step. After that 100  $\mu$ l of TMB (Tetra Methyl Benzidine) substrate solution was added to each well and incubated plate at room temperature for 15 mins for the colour development. Following substrate addition, 50  $\mu$ l of 2N H<sub>2</sub>SO<sub>4</sub> was added to each well to stop the reaction. The absorbance measurements were read at 450 nm with a Tecan M 200 multimode reader. The concentrations of cytokines released were obtained and the percentage inhibition of cytokines production was calculated as compared to control conditions.


### Evaluation of *in vivo* anti-inflammatory activity

#### Experimental animals

The male wistar rats weighing between 130 - 150 gm were used for the experiments. They were kept in polypropylene cages under standard laboratory conditions (12:12 hr light/dark) cycle at 24°C. The rats were provided with commercial rat diet (NIN, Hyderabad) and water *ad libitum*. The experiments were conducted after obtaining approval from Institutional Animal Ethical Committee of IICT. The animals were quarantined and acclimatized to laboratory conditions for 7 days prior to study initiation, and they were also observed for general health and suitability for testing during this period.

#### Carrageenan induced rat paw edema

The anti-inflammatory activity of the test compounds was evaluated in wistar rats employing the standard method.<sup>[25]</sup> The animals were fasted overnight and were divided into control, standard, and different test groups. The test compounds were administered by oral route as gum acacia suspension (2% w/v) at the dose of 100 mg/kg. The animals in the standard group, received indomethacin at the dose of 10 mg/kg by oral route, and rats in the control group received the vehicle solution without test compounds. One hour after test drugs administration, rats in all the groups were administered with 0.1 ml of 1% carrageenan in the sub plantar region of right hind paw. The paw volumes were measured before and after 3 hrs after the administration of carrageenan using



**Figure 1:** Isolated compounds in rhizomes of *Cyperus scariosus* R.Br and *Cyperus rotundus* L.

digital plethysmometer (Ugo Basile, Italy). The percentage inhibition of paw volume for treated groups was calculated by comparing with mean paw volume of control group.

**Percentage of edema inhibition** =  $1 - V_t / V_c \times 100$

Where  $V_t$  = Volume of paw edema in drug treated groups

$V_c$  = Volume of paw edema in control groups

## RESULTS AND DISCUSSION

### Isolation of compounds and structure elucidation

The hexane, chloroform and methanol extracts of *C. scariosus* and *C. rotundus* were chromatographed on silica gel (100-200 mesh), and extensive purification of the resultant fractions resulted in the isolation of twelve compounds<sup>[1-12]</sup>. The structures of the isolates were established by the interpretation of the NMR and Mass spectral data. After comparing their physicochemical and spectrometric data with those reported in the literature, they were identified as known compounds and confirmed as: Stigmasterol<sup>[1,26]</sup>  $\beta$ - sitosterol<sup>[2,27]</sup> Lupeol<sup>[3,26]</sup> Gallic acid<sup>[4,28]</sup> Quercetin<sup>[5,29]</sup>  $\beta$ - amyryl<sup>[6,30]</sup> Oleanolic acid<sup>[7,31]</sup>  $\beta$ - amyryl acetate<sup>[8,30]</sup> 4- hydroxy butyl cinnamate<sup>[9,32]</sup> 4- hydroxy cinnamic acid<sup>[10,33]</sup> Caffeic acid<sup>[11,34]</sup> and kaempferol<sup>[12,35]</sup>. Structures of these compounds are shown in [Figure 1].

#### Spectral data:

**Stigmasterol (1):** White solid (200 mg), mp 174-176°C, EIMS: ( $m/z$ ) 412 [ $M^+$ ], IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3424, 2936, 2867, 1640, 1464  $\text{cm}^{-1}$ .  $^1\text{H NMR}$  [300 MHz,  $\text{CDCl}_3$ ]:  $\delta$  5.35 [t,  $J = 6.1\text{Hz}$ , 1H], 5.14 [m, 1H], 4.98 [m, 1H], 3.52 [m, 1H], 2.52 - 2.08 [m, 5H], 1.98 - 1.92 [m, 3H], 1.03 [3H, s], 1.51 [m, 1H], 1.52 [m, 2H], 1.32 - 1.40 [m, 3H], 1.18 [m, 2H], 1.14 [m, 2H], 1.01 [s, 3H], 1.01 [s, 3H], 1.02 [m, 1H], 0.96 [m, 1H], 0.91 [d,  $J = 6.2\text{ Hz}$ , 3H], 0.83 [t,  $J = 7.2\text{ Hz}$ , 3H], 0.82 (d,  $J = 6.6\text{ Hz}$ , 3H), 0.80 (d,  $J = 6.6\text{ Hz}$ , 3H), 0.71 [s, 3H].  $^{13}\text{C NMR}$  [75 MHz,  $\text{CDCl}_3$ ]:  $\delta$  140.71 [C-5], 138.2 [C-22], 72.1 [C-3], 121.7 [C-6], 56.8 [C-14], 56.2 [C-17], 51.1 [C-24], 29.6 [C-25], 42.4 [C-13], 42.6, 40.4, 39.8, 37.5, 36.4, 32.3, 32.1 [C-2], 31.8 [C-7, C-8], 20.9 [C-21], 29.3 [C-16], 18.9 [C-28], 24.4 [C-15], 20.9 [C-21], 21.5 [C-11], 19.3 [C-27], 21.7 [C-19], 12.2 [C-29], 40.6 [C-18].

**$\beta$ -sitosterol (2):** White amorphous powder (150 mg), mp 134-135 °C, EIMS: ( $m/z$ ) 414 [ $M^+$ ], IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3424, 2930, 2852, 1724, 1463, 1378, 1271, 1059, 1023, 963 and 802  $\text{cm}^{-1}$ .  $^1\text{H NMR}$  ( $\text{CDCl}_3$ , 300 MHz):  $\delta$  5.36 (1H, d,  $J = 6.4\text{Hz}$ , H-6), 3.53 (1H, m, H-3), 1.01, 0.68 (3H, s, H-19

and H-18), 0.83 (3H, d,  $J = 6.4\text{ Hz}$ , H-21), 0.81 (3H, d,  $J = 6.4\text{ Hz}$ , H-29) and 0.85 (3H, t,  $J = 7.1\text{ Hz}$ , H-26).  $^{13}\text{C NMR}$  ( $\text{CDCl}_3$ , 75MHz):  $\delta$  37.5 (C-1), 31.9 (C-2), 72.0 (C-3), 42.5 (C-4), 140.9 (C-5), 121.9 (C-6), 32.1 (C-7), 32.1 (C-8), 50.3 (C-9), 36.7 (C-10), 21.3 (C-11), 39.9 (C-12), 42.6 (C-13), 56.9 (C-14), 26.3 (C-15), 28.5 (C-16), 56.3 (C-17), 36.3 (C-18), 19.2 (C-19), 34.2 (C-20), 26.3 (C-21), 36.2 (C-22), 46.1 (C-23), 23.3 (C-24), 12.2 (C-25), 29.4 (C-26), 20.1 (C-27), 19.6 (C-28), 12.0 (C-29).

**Lupeol (3):** Pale yellow powder (100 mg) mp 215-216°C, EIMS: ( $m/z$ ) 426 [ $M^+$ ], IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3308, 2924, 1465, 1379, 1042 and 880  $\text{cm}^{-1}$ .  $^1\text{H NMR}$  (200 MHz,  $\text{CDCl}_3$ ):  $\delta$  4.69 and 4.56 (each 1H, s, H-29), 3.16 (1H, dd, H-3), 2.35 and 1.49 (each 1H, m, 21a), 1.67 (1H, t, H-15a), 1.71 (3H, s, H-30), 1.61 (2H, d, H-12a, 1b), 1.67 (1H, t, H-13), 1.50 (1H, q, H-2b), 1.42 (1H, m, H-22a), 1.41 (2H, m, H-7), 1.50 (1H, q, H-6B), 1.38 (1H, t, H-16b), 1.39 (1H, m, H-21B), 1.25 (1H, q, H-11B), 1.28 (1H, m, H-22b), 1.71 (3H, s, H-26), 1.00 (3H, s, H-23), 0.94 (3H, s, H-27), 0.92 (s, 3H), 0.80, 0.79 and 0.73 (each 3H, H-25, 28, 29) and 0.66 (1H, d, H-5).  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  150.9 (C-20), 109.3 (C-29), 79.0 (C-3), 55.3 (C-5), 50.4 (C-9), 48.0 (C-18), 48.3 (C-19), 25.1 (C-12), 42.8 (C-14), 40.8 (C-8), 40.0 (C-22), 38.8 (C-4), 38.7 (C-1), 38.0 (C-13), 37.1 (C-10), 35.6 (C-16), 34.3 (C-7), 29.1 (C-21), 28.0 (C-23), 27.4 (C-15), 27.4 (C-2), 25.1 (C-12), 28.0 (C-23), 27.4 (C-15), 20.9 (C-11), 19.3 (C-30), 18.3 (C-6), 18.0 (C-28), 15.9 (C-25), 16.1 (C-26), 15.3 (C-24) and 14.6 (C-27).

**Gallic acid (4):** White amorphous solid (20 mg), EIMS (negative mode)  $m/z$  169 [M-H], mp 260°C. IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3369, 3286, 1711, 1620, 1540, 1308 and 1024  $\text{cm}^{-1}$ .  $^1\text{H NMR}$  (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.02 (1H, d,  $J = 0.8$ ).  $^{13}\text{C NMR}$  (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  120.4 (C-1), 109.2 (C-2), 144.3 (C-3), 17.6 (C-4), 144.3 (C-5), 109.2 (C-6) and 169.1 (C-7).

**Quercetin (5):** Pale yellow powder (20 mg), mp 300°C, ESI-MS (negative mode)  $m/z$  301 [M-H]. IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3407, 3318, 1667, 1610, 1521, 1382, 1168, 1131, 1013 and 821  $\text{cm}^{-1}$ .  $^1\text{H NMR}$  (300MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  6.08 (1H, d,  $J = 2.07$ ), 6.29 (1H, d,  $J = 2.07\text{Hz}$ ), 7.64 (1H, d,  $J = 2.2\text{Hz}$ ), 6.78 (1H, d,  $J = 8.4\text{Hz}$ ), 7.54 (1H, dd,  $J = 2.2$  & 8.4Hz).  $^{13}\text{C NMR}$  (75MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  148.2 (C-2), 137.2 (C-3), 177.5 (C-4), 162.6 (C-5), 99.4 (C-6), 165.7 (C-7), 94.6 (C-8), 158.4 (C-9), 104.7 (C-10), 124.3 (C-11), 116.1 (C-12), 146.3 (C-13), 150.3 (C-14), 116.6 (C-15) and 121.8 (C-16).

**$\beta$ -Amyryl (6):** White amorphous powder (8 mg), mp 189-191°C, IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3536 and 1619. EIMS:  $m/z$  426 [ $M^+$ ].  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ ): 5.26 (d,  $J = 6.0\text{Hz}$ , 1H), 3.28 (dd,  $J = 4.9, 10.8\text{Hz}$ , 1H), 2.12-1.86 (m, 2H), 1.84-1.69 (m, 4H), 1.66-1.48 (m, 11H) overlapped with each other, 1.46-1.28 (m, 7H, overlapped), 1.06 (s, 3H), 1.0 (s, 3H), 0.98 (s, 2x3H), 0.96 (s, 3H), 0.86 (s, 3H), 0.78 (s, 3H), 0.75 (s, 3H).  $^{13}\text{C NMR}$  ( $\text{CDCl}_3$ , 75MHz):  $\delta$  138.6, 116.8, 78.0 54.2, 46.8, 48.2, 42.6, 42.4, 43.0, 38.6, 38.4, 37.2, 36.4, 36.2, 34.5, 28.6, 28.4, 27.3, 26.0, 22.4, 22.0, 18.4, 17.6, 16.4, 15.8, 15.2 and 14.8.

**Oleanolic acid (7):** White amorphous powder (5 mg), EIMS:  $m/z$  456 [ $M^+$ ]. IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3423, 2922, 2851, 1691, 1463, 1383 and 1182  $\text{cm}^{-1}$ .  $^1\text{H NMR}$  ( $\text{CDCl}_3$ , 300 MHz):  $\delta$  1.02 (1H, m), 1.57 (1H, m), 3.24 (1H, dd,  $J = 10.5\text{Hz}$ , CH-3), 0.88 (1H, m), 1.58 (1H, m), 1.39 (1H, m), 1.53 (1H, m), 1.36 (1H, m), 1.71 (1H, m), 1.96 (1H, m), 5.28 (1H, s), 1.72 (1H, m), 2.19 (1H, m), 2.12 (1H, m), 1.96 (1H, m), 2.83 (1H, dd,  $J = 13.8, 4.5\text{Hz}$ ), 1.83 (1H, m), 1.32 (1H, m), 1.46 (1H, m), 1.23 (1H, m), 1.82 (1H, m), 2.04 (1H, m), 1.00 (3H, s), 0.79 (3H, s), 0.93 (3H, s), 0.77 (3H, s), 1.15 (3H, s), 1.15 (3H, s), 0.92 (3H, s) and 0.95 (3H, s).  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  15.3 (C-25), 15.4 (C-24), 17.1 (C-26), 18.3 (C-6), 22.9 (C-11), 23.6 (C-30), 23.4 (C-16), 25.9 (C-27), 27.2 (C-2), 28.1 (C-23), 28.4 (C-15), 30.7 (C-20), 32.4 (C-22), 32.6 (C-29), 33.1 (C-7), 33.8 (C-21), 37.1 (C-10), 38.4 (C-1), 38.8 (C-4), 39.3 (C-8), 41.0 (C-18), 41.6 (C-14), 45.9 (C-19), 46.7 (C-17), 47.6 (C-9), 55.2 (C-5), 79.6 (C-3), 122.6 (C-12), 143.6 (C-13), 182.9 (C-28).

**$\beta$ -Amyrin acetate (8):** colourless needles (6 mg), IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3490, 2986, 1606, 1482 and 1066  $\text{cm}^{-1}$ . The EI-MS:  $m/z$  468 [M<sup>+</sup>]. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz):  $\delta$  5.18 (t,  $J$  = 3.5 Hz, 1H), 4.50 (t,  $J$  = 8.0 Hz), 2.05 (s, 3H), 1.64 (m), 1.88 (m), 0.84 (m), 1.53 (m), 1.58 (m), 1.80 (m), 1.76 (m), 1.98 (m), 2.31 (m), 1.93 (m), 1.60 (m), 1.32 (m), 1.08 (m), 0.88 (s), 0.96 (s), 0.86 (s), 0.97 (s), 1.13 (s), 0.83 (s), 0.88 (s), 0.87 (s). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz):  $\delta$  171.1 (C-31), 145.2 (C-13), 121.6 (C-12), 81.0 (C-3), 55.2 (C-5), 47.6 (C-9), 47.2 (C-18), 46.8 (C-19), 39.8 (C-4), 38.3 (C-1), 37.7 (C-8), 34.7 (C-21), 33.4 (C-29), 31 (C-20), 26.9 (C-16), 26.1 (C-15), 23.4 (C-11), 21.4 (C-32), 18.3 (C-6), 16.8 (C-26), 16.7 (C-25) and 15.6 (C-24).

**[4-] hydroxy butyl cinnamate (9):** white amorphous powder (5 mg), IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : IR 3388, 2921, 1675  $\text{cm}^{-1}$ . Mass  $m/z$  220 [M<sup>+</sup>]. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz):  $\delta$  6.84 (1H, d,  $J$  = 8.6), 7.4 (1H, d,  $J$  = 8.6 Hz), 7.63 (1H, d,  $J$  = 15.6 Hz), 6.30 (1H, d,  $J$  = 15.6 Hz), 4.18 (2H, t,  $J$  = 6.7 Hz), 1.69 (2H, m), 0.88 (3H, t,  $J$  = 6.9 Hz). The <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz):  $\delta$  127.4 (C-1), 115.6 (C-2), 129.9 (C-3), 165.9 (C-4), 129.9 (C-5), 115.8 (C-6), 156.3 (C-7), 167.4 (C-9), 64.64 (C-10), 25.98 (C-11), 22.68 (C-12) and 14.1 (C-13).

**[4-] hydroxy cinnamic acid (10):** white solid (6 mg), IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3371, 2929, 1672  $\text{cm}^{-1}$ . Mass  $m/z$  163 [M-H]. The <sup>1</sup>H NMR (CD<sub>3</sub>OD), 300 MHz):  $\delta$  6.84 (1H, d,  $J$  = 8.4 Hz), 7.43 (1H, d,  $J$  = 8.4 Hz), 7.64 (1H, d,  $J$  = 16.05 Hz), 6.26 (1H, d,  $J$  = 16.05 Hz). The <sup>13</sup>C NMR (CD<sub>3</sub>OD, 75 MHz):  $\delta$  114.2 (C-1), 115.2 (C-2), 129.7 (C-3), 159.0 (C-4), 159.0 (C-4), 129.7 (C-5), 115.2 (C-6), 145.4 (C-7), 125.7 (C-8) and 169.5 (C-9).

**Caffeic acid (11):** yellow solid (10 mg), mp 137-140°C, ESI-MS  $m/z$  179 [M-H]. IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3407, 3229, 2923, 1645 and 1449  $\text{cm}^{-1}$ . <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300 MHz):  $\delta$  6.97 (1H, d,  $J$  = 1.8), 6.73 (1H, d,  $J$  = 8.1 Hz), 6.85 (1H, dd,  $J$  = 1.8 & 8.1 Hz), 6.12 (1H, d,  $J$  = 15.8 Hz), 7.48 (1H,  $J$  = 15.8 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz):  $\delta$  128.0 (C-1), 115.2 (C-2), 145.9 (C-3), 146.5 (C-4), 117.2 (C-5), 123.2 (C-6), 144.9 (C-7), 116.5 (C-7), 116.5 (C-8), 171.5 (C-9).


**Kaempferol (12):** Yellow powder (5 mg), mp 286°C. ESI-MS:  $m/z$  285 [M-H]. <sup>1</sup>H NMR (DMSO- $d_6$ , 300 MHz):  $\delta$  6.27, 6.46 (1H,  $J$  = 2.1 Hz), 8.0 (1H, d,  $J$  = 8.7 Hz), 6.99 (1H, d,  $J$  = 8.7 Hz). <sup>13</sup>C NMR (DMSO- $d_6$ , 75 MHz):  $\delta$  145.9 (C-2), 135.5 (C-3), 175.5 (C-4), 156.4 (C-5), 99.06 (C-6), 167.3 (C-7), 93.8 (C-8), 160.5 (C-9), 101.5 (C-10), 121.7 (1'), 129.1 (2'), 115.4 (3'), 159.2 (4'), 115.4 (5'), 129.1 (6').

## BIOLOGICAL ACTIVITIES OF ISOLATED COMPOUNDS

### DPPH and ABTS radical scavenging activity

In present days, oxidative stress leads to several diseases in human beings such as atherosclerosis, diabetes, cancer, inflammation, cancer, alzheimer's,

and parkinson's.<sup>[36]</sup> The oxidative stress is mainly caused by free radicals. These species include both oxygen and non oxygen radicals. The oxygen radical include peroxy, superoxide anion, hydroxyl, peroxy nitrile, and nitric oxide radical. The non oxygen radical includes H<sub>2</sub>O<sub>2</sub>, hypochlorous acid, and singlet oxygen. These free radicals (ROS) produced by biochemical reaction in human body cause the structural and functional changes in human metabolism. The various plant extracts and different types of secondary metabolites have been shown not only as free radical scavenging and antioxidant activities. This characteristic hydroxyl functional group scavenge free radicals, and prevent the oxidation of biological molecules by converting the more ROS by donating hydrogen atom into inactive species. Therefore, antioxidant and free radical scavenging constituents are more useful in the prevention and treatment of oxidative stress induced disorders and diseases. In our study, all these isolates were evaluated for their DPPH (1, 1-diphenyl-2-picrylhydrazyl) and ABTS + [2, 2'- azino - bis (3-ethyl benzothiazoline-6-sulfonic acid)] free radical scavenging activities. The values obtained for antioxidant assays were represented as percentages of radical scavenging activities at various concentrations. IC<sub>50</sub> values denote the concentration of sample required to scavenge 50% DPPH radicals [Figure 2a]. In our experiment, we used a well known anti oxidant ascorbic acid as standard. As shown in [Table 1], kaempferol<sup>[12]</sup> caffeic acid<sup>[11]</sup> quercetin,<sup>[5]</sup> and gallic acid<sup>[4]</sup> were displayed potent DPPH scavenging activities with IC<sub>50</sub> values of 3.85, 1.24, 1.1 and 0.43  $\mu\text{g/ml}$ , respectively. While the remaining compounds, stigmasterol<sup>[1]</sup>, sitosterol<sup>[2]</sup> and lupeol<sup>[3]</sup> have not shown DPPH inhibition at 25  $\mu\text{g/ml}$ . The pattern and potentials in decreasing order of DPPH scavenging activity was observed as follows: **gallic acid > quercetin > caffeic acid > kaempferol**. Similarly, all these compounds were evaluated for ABTS free radical scavenging activity using trolox as standard. The results represents the percentage inhibition of ABTS at 10  $\mu\text{g/ml}$  [Table 1 and Figure 2b]. The analysis of the results indicated that quercetin<sup>[5]</sup>, kaempferol<sup>[12]</sup>, cinnamic acid<sup>[10]</sup>, caffeic



**Figure 2:** Concentration dependent (a) DPPH and (b) ABTS free radical scavenging pattern of compounds isolated from *C. scariosus* and *C. rotundus*.

**Table 1:** Concentration required for scavenging DPPH and ABTS free radicals by 50% (IC<sub>50</sub>).

Sample	% inhibition of DPPH at 25 $\mu\text{g/ml}$	DPPH IC <sub>50</sub> $\mu\text{g/ml}$	% inhibition of ABTS at 10 $\mu\text{g/ml}$	ABTS IC <sub>50</sub> $\mu\text{g/ml}$
Stigmasterol	NA	--	NA	--
Lupeol	NA	--	NA	--
Kaempferol	87.8 $\pm$ 0.8	3.85 $\pm$ 0.4	97.61 $\pm$ 1.1	0.45 $\pm$ 0.05
Caffeic acid	84.3 $\pm$ 0.4	1.24 $\pm$ 0.1	96.72 $\pm$ 0.9	0.48 $\pm$ 0.2
Gallic acid	83.3 $\pm$ 1.0	0.43 $\pm$ 0.1	94.13 $\pm$ 2.0	0.56 $\pm$ 0.01
4- hydroxy cinnamic acid	22.4 $\pm$ 2.3	--	99.4 $\pm$ 0.5	0.45 $\pm$ 0.01
Quercetin	85.6 $\pm$ 1.2	1.1 $\pm$ 0.2	98.21 $\pm$ 0.2	0.067 $\pm$ 0.01
$\beta$ - sitosterol	NA	--	34.83 $\pm$ 2 $\phi$ .0	--
Ascorbic acid	84.2 $\pm$ 0.6	2.17 $\pm$ 0.2		
Trolox			93.64 $\pm$ 0.9	0.1 $\pm$ 0.01

Note: NA= Not active, Values represent mean  $\pm$  SD (n=3)

acid<sup>[11]</sup> and gallic acid<sup>[4]</sup> displayed potent ABTS scavenging activities with IC<sub>50</sub> values of 0.067, 0.45, 0.45, 0.48, and 0.56 µg/ml, respectively. The order of the inhibitory potentials was observed as follows: **quercetin > kaempferol = cinnemic acid > caffeic acid > gallic acid**. However, the compounds stigmasterol<sup>[1]</sup>, sitosterol<sup>[2]</sup> and lupeol<sup>[3]</sup> has not shown ABTS inhibition at 10 µg/ml. These observations revealed the fact that that presence phenolic hydroxyl groups positively affect compounds radical scavenging potential, and absence of phenolic hydroxyl group drastically reduced its DPPH and ABTS radical scavenging capacity. The phenolic compounds are being major contributors to the antioxidants behaviour of plants and the antioxidant activity of phenols is mainly due to their redox properties, which allow them to act as reducing agents, hydrogen donors, and singlet oxygen.<sup>[37]</sup> Furthermore, free radical scavenging and antioxidant activity of phenols mainly depends on the number and position of hydrogen-donating hydroxyl groups on the aromatic ring of the phenol molecules. This characteristic of hydroxyl functional group scavenge free radicals and prevent the oxidation of biological molecules by converting the more ROS by donating hydrogen atom into inactive species.<sup>[38]</sup> Therefore, antioxidant and free radical scavenging constituents are more useful in the prevention and treatment of oxidative stress induced disorders and diseases.

#### *α*-glucosidase and *α*-amylase inhibitory activities


*C. scariosus* and *C. rotundus* are widely used in ayurvedic preparation for the treatment of diabetes and other diseases. However, there has been relatively little information pertaining to its active constituents and their alpha-glucosidase activities. Thus, in this study, we have tested all the compounds isolated from *C. scariosus* and *C. rotundus* for *α*-glucosidase inhibition activity using a standard protocols. In our experiment, we used a well known anti diabetic drug acarbose as standard. Our results represent that percentage inhibition of *α*-glucosidase at 20 µg/ml [Table 2

and Figure 3a]. Among all the compounds quercetin<sup>[5]</sup> and lupeol<sup>[3]</sup> showed [24.7%] and [13.1%] inhibition of alpha glucosidase activity at 20 µg/ml, respectively, whereas as the other compounds showed mild activity. Similarly, we also measured *α*-amylase inhibitory potentials of all the isolates. The results represent the percentage inhibition of *α*-amylase at 40 µg/ml [Table 2 and Figure 3b]. The results indicated that gallic acid<sup>[4]</sup> and 4- hydroxy cinnemic acid<sup>[10]</sup> showed alpha amylase activity with the 31.2% and 23.5% inhibitory potentials. Comparing our experimental results revealed the fact that phenolics and flavanoids have shown antidiabetic activity. The results from earlier studies suggest that flavanoids and phenolics isolated from several medicinal plants could have potential to treat type-II diabetes mellitus.<sup>[39]</sup> These compounds are capable not only for oxidative stress but also of inhibiting carbohydrate hydrolyzing enzymes to prevent hyperglycemia.<sup>[40]</sup>


**Table 2:** Percentage inhibition of *α*-glucosidase and *α*-amylase activities

Sample	% inhibition of AGI at 20 µg/ml	% inhibition of alpha amylase at 40 µg/ml
Stigmasterol	4.0 ± 0.7	12.3 ± 1.3
Lupeol	13.1 ± 1.9	07.3 ± 1.3
Kaempferol	NA	05.4 ± 0.8
Caffeic acid	4.8 ± 1.0	09.7 ± 0.5
Gallic acid	2.7 ± 0.9	31.2 ± 0.2
4- hydroxy cinnemic acid	6.3 ± 0.7	23.5 ± 1.5
Quercetin	24.7 ± 1.0	05.6 ± 0.6
β- sitosterol	NA	10.2 ± 1.0
Acarbose	61.6 ± 0.4	40.0 ± 1.0


Note: NA= Not active, Values represent mean ± SD (n=3)



**Figure 3:** (a). *α*-glucosidase (b) *α*-amylase inhibitory potential of isolated compounds.



**Figure 4:** Phase contrast images of control, standard and test compounds in Thp-1 cells captured by light microscopy. (a). THP-1 monocytic cells. (b). THP-1 cells treated with PMA to induce monocyte to macrophage differentiation. (c). THP-1 cells treated with stigmasterol without PMA. (d). THP-1 cells + PMA with stigmasterol. (e). THP-1 cells treated with sitosterol without PMA. (f). THP-1 cells + PMA with sitosterol. (g). THP-1 cells treated with lupeol without PMA. (h). THP-1 cells + PMA with lupeol. (i). THP-1 cells treated with piroxicam (standard) without PMA. (j). THP-1 cells + PMA with piroxicam.




**Figure 5:** Represents % inhibition of IL-1β activity in ThP-1 monocytic cells treated with stigmasterol, sitosterol and lupeol.

## In vitro anti-inflammatory screening of compounds 1, 2 and 3 using PMA- induced production of IL-1 $\beta$ model

### Measurement of IL-1 $\beta$ production by ELISA

During acute or chronic inflammation excessive IL-1 $\beta$  is produced as pro inflammatory cytokine, which results in the systemic inflammatory response syndrome, severe tissue damage, and septic shock. The effects of stigmasterol<sup>[1]</sup>,  $\beta$ - sitosterol,<sup>[2]</sup> and lupeol<sup>[3]</sup> isolated from *C. scariosus* on modulating the production levels of IL-1  $\beta$ , was tested by stimulating human monocytic cell line THP-1, using PMA, and the results were compared with standard drug piroxicam. The THP-1 monocytic cells were treated with 100 ng/ml PMA in the presence or absence of isolated compounds and standard piroxicam 10  $\mu$ M/ml. After 48 hours PMA stimulation, the monocytes differentiated into macrophages and cytokines

released into the media. Following stimulation period, supernatant was collected and the respective wells washed twice with phosphate buffer saline (PBS), and phase contrast images were taken by light microscopy. The phase contrast micrographs of control cells and treated cells are given in [Figure 4]. The cells and nuclei of the control cells are normal with clear-cut membranes, while nuclei are elliptical. Whereas, the cells stimulated with PMA, the monocytes differentiated into macrophages, and some of the phagocytic changes observed in the cultured media. Whereas, the cells treated with test compounds stigmasterol,<sup>[1]</sup>  $\beta$ - sitosterol<sup>[2]</sup> and lupeol<sup>[3]</sup> are showing distorted membranes and nucleus with blebs. After morphological changes observation, from the collected supernatant we estimated IL-1  $\beta$  levels by using sandwich ELISA as per the manufacturers instructions. Among the three tested compounds lupeol exhibited 47% promising result by down regulating levels of IL-1 $\beta$ , whereas stigmasterol (1),  $\beta$ - sitosterol<sup>[2]</sup> were found



**Figure 6:** *In vivo* anti-inflammatory activity in rats. (a). Oral feeding (b). Treated with carrageenan (c). Treated with stigmasterol (d). Treated with sitosterol (e). Treated with lupeol. (f). Treated with indomethacin (g). Measurement of paw volume by using Plethysmometer.

**Table 3:** Effect of compounds 1,2 and 3 of *C. scariosus* on carrageenan-induced paw edema in rats.

Groups	Dose (mg/kg)	Paw volume (0hr)	Paw volume (3hr)	Paw volume difference	% inhibition at 3hr
Carrageenan (Control)	--	1.205 $\pm$ 0.012	2.545 $\pm$ 0.03	1.34 $\pm$ 0.033	--
Stigmasterol	100	1.17 $\pm$ 0.044	2.635 $\pm$ 0.044	1.46 $\pm$ 0.026	0.0 $\pm$ 0.0
$\beta$ - sitosterol	100	1.18 $\pm$ 0.040	2.54 $\pm$ 0.099	1.36 $\pm$ 0.063	2.42 $\pm$ 1.59
Lupeol	100	1.23 $\pm$ 0.008	2.41 $\pm$ 0.014	1.18 $\pm$ 0.007(a)	11.94 $\pm$ 0.52(a)
Indomethacin (Standard)	10	1.18 $\pm$ 0.018	1.92 $\pm$ 0.0359	0.73 $\pm$ 0.035(b)	44.9 $\pm$ 2.66(b)

Note: Values represent mean  $\pm$  SD (n=4). Unpaired 't' test with welch's correction was applied for Statistical analysis. <sup>a</sup>p < 0.0025 and <sup>b</sup>p < 0.0001, when compared with paw volume of control group at the end of 3hr.

to be equally active and showed mild activity in inhibition of IL-1 $\beta$  as compared to lupeol [Figure 5]. The three compounds activity results were compared with standard drug piroxicam (10  $\mu$ M), which exhibited 75% inhibition of IL-1 $\beta$  concentration. Around 53 % of IL-1 $\beta$  inhibition was observed with 10  $\mu$ M/ml treatment of lupeol.<sup>[3]</sup> Our results revealed that among the three compounds, lupeol has shown to decrease the generation of proinflammatory cytokine IL-1 $\beta$ . The earlier studies showed that phytosterols and triterpenoids decrease the generation of proinflammatory cytokines such as tumour necrosis factor- $\alpha$  and interleukin-1 $\beta$  in lipopolysaccharide in treating macrophages.<sup>[41,42]</sup>

### In vivo anti-inflammatory activity of compounds 1, 2 and 3

In traditional medicine, rhizomes of *C. scariosus* were used to treat inflammation. Till today there is no scientific proof for justification of traditional use of rhizomes in treatment of inflammation. Hence, the present study focussed to study anti-inflammatory activity of compounds<sup>[1-3]</sup> isolated from *C. scariosus*. They were studied for *in vivo* anti inflammatory activity by using carrageenan-induced rat paw edema as the method. The Carrageenan is a strong chemical used for the release of inflammatory and pro inflammatory mediators (prostaglandins, leukotrienes, histamine, bradykinin, TNF- $\alpha$ , etc). To test the anti inflammatory activity, male wistar rats were divided into three groups such as control, standard, and different test groups. The percentage reduction in paw volume after treatment with respective test compounds was considered as anti-inflammatory activity. It was found that stigmasterol 1 and  $\beta$ -sitosterol<sup>[2]</sup> did not display anti-inflammatory activity. However, lupeol<sup>[3]</sup> reduced significantly ( $p < 0.0025$ ) the paw volume, when compared with the control group of animals [Figure 6 and Table 3]. Therefore, it is inferred that the lupeol<sup>[3]</sup> has anti-inflammatory effect in rats at a dose of 100 mg/kg body weight with marked inhibition 12% inflammation at the third hour. Comparing these results with earlier studies, lupeol and its derivatives were shown to exhibit higher anti-inflammatory activity than commonly used non steroidal anti-inflammatory drug indomethacin in rat and mouse models of inflammation.<sup>[43]</sup>

## CONCLUSION

In this study, we investigated anti-oxidant, anti-diabetic, and anti-inflammatory constituents from the Indian sedges which resulted in the isolation of twelve compounds. Biological activities of these isolates revealed that aromatic compounds gallic acid<sup>[4]</sup> quercetin<sup>[5]</sup> caffeic acid<sup>[11]</sup>, and kaempferol<sup>[12]</sup> showed potent antioxidant activity and anti-diabetic activity. While lupeol<sup>[3]</sup> showed potent IL-1  $\beta$  activity inhibition in THP-1 monocytic cells, and also displayed significant ( $p < 0.0025$ ) *in vivo* anti-inflammatory activity in carrageenan induced rat paw edema.

## Acknowledgment

The authors would like to thank the management of KLEF University and Director, CSIR- Indian Institute of Chemical Technology for providing facilities. One of the authors, Lavanya Kakarla is thankful to DST, New Delhi for financial support under Women scientist (WOS-A) scheme.

## Financial support and sponsorship

This work was financially supported by Department of Science and Technology, New Delhi, India, under the Women Scientist (WOS-A) project grant (SR/WOS-A/LS-1160/2014 (G)).

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

1. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods impact on human health. *Pharmacognosy Reviews* 2010;4:118-16.
2. Oh J, Jo SH, Kim JS, Ha KS, Lee JY, Choi HY, YU SY, et al. Selected tea and tea pomace extracts inhibit intestinal  $\alpha$ -glucosidase activity in *in vitro* and postprandial hyperglycemia *in vivo*. *Int J Mol Sci* 2015;16:8811-25.
3. Derosa G, Maffioli P.  $\alpha$ -glucosidase inhibitors and their use in clinical practice. *Archives of Medical Science* 2012;8:899-6.
4. Dinarello CA. Anti inflammatory agents present and future. *Cell* 2010;140:935-50.
5. Dabhi AS, Bhatt NR, Shah MJ. Voglibose an alpha glucosidase inhibitor. *Journal of Clinical and Diagnostic Research* 2013;7:3023-27.
6. Apetz N, Munch G, Govindaraghavan S, Gyengesi E. Natural compounds and plant extracts as therapeutics against chronic inflammation in Alzheimer's disease-a translational perspective. *CNS Neurol Disord drug targets* 2014;13:1175-91.
7. Uddin SJ, Mondal K, Shilpi JA, Rahman MT. Antidiarrhoeal activity of *Cyperus rotundus*. *Fitoterapia* 2006;77:134-36.
8. Alam MA, Jahan R, Rahman S, Das AK, Rahmatullah M. Antinociceptive and anti-hyperglycemic activity of methanol leaf extract of *Cyperus scariosus*. *Pak J Pharm Sci* 2011;24:53-6.
9. Tsoyi K, Jang HJ, Lee YS, Kim YM, Kim HJ, Seo HG. (+) – nootkatone and (+) – valence from rhizomes of *Cyperus rotundus* increase survival rates in septic mice due to heme oxygenase-1 induction. *Journal of Ethnopharmacology* 2011;137:1311-7.
10. Pirzada AM, Aii HH, Naeem M, Latif M, Bukhari AH, Tanveer A. *Cyperus rotundus* L Traditional uses, phytochemistry and pharmacological activities. *J Ethnopharmacol* 2015;174:540-60.
11. Seo WG, Pae HO, Oh GS, Chai KY, Kwon TO, Yun YG, et al. Inhibitory effects of methanol extract of *Cyperus rotundus* rhizomes on nitric oxide and superoxide productions by murine macrophage cell line, Raw 264.7 cells. *J Ethnopharmacol* 2001;76:59-64.
12. Nagarajan M, Kuruvilla GR, Kumar KS, Venkatasubramanian P. Abhava pratinidhi dravya: A comparative phytochemistry of *Ativisha*, *Musta* and related species. *J Ayurveda Integr Med* 2015;6:53-63.
13. Jin JH, Lee DU, Kim YS, Kim HP. Anti-allergic activity of sesquiterpenes from the rhizomes of *Cyperus rotundus*. *Archives of Pharmacol Research* 2011;34:223-28.
14. Wadhwa M, Bakshi MPS. Herbal feed additives - impact on the rumen environment in buffaloes. *Ind J Animal Nutri* 2006;23:102-9.
15. Park SE, Shin WT, Park C, Hong SH, Kim GY, Kim SO, et al. Induction of apoptosis in MDA-MB-231 human breast carcinoma cells with an ethanolic extract of *Cyperus rotundus* L by activating caspases. *Oncol Rep* 2014;32:2461-70.
16. Gilani AU, Janbaz KH. Studies on protective effect of *Cyperus scariosus* extract on acetaminophen and CCl<sub>4</sub>-induced hepatotoxicity. *General Pharmacology* 1995;26:627-31.
17. Gupta SK, Sharma RC, Aggarwal OP, Arora RB. Anti-inflammatory activity of the oil isolated from *Cyperus scariosus*. *Indian Journal of Experimental Biology* 1972;10:41-2.
18. Thomas D, Govindhan S, Baiju EC, Padmavathi G, Kunnumakkara AB, Padikkala J. *Cyperus rotundus* L prevents non-steroidal anti-inflammatory drug-induced gastric mucosal damage by inhibiting oxidative stress. *J Basic Clin Physiol Pharmacol* 2015;26:485-90.
19. Sayed HM, Mohamed MH, Farag SF, Mohamed GA, Omobuwajo OR, Proksch P. Fructose-amino acid conjugate and other constituents from *Cyperus rotundus* L. *Nat Prod Res* 2008;22:1487-97.
20. Ahn JH, Lee TW, Kim KH, Byun H, Ryu B, Lee KT, et al. 6-Acetoxy cyperene, a patchoulane-type sesquiterpenes isolated from *Cyperus rotundus* rhizomes induces caspase-dependent apoptosis in human ovarian cancer cells. *Phytother Res* 2015;10:doi: 10.1002/ptr.5385
21. Kakarla L, Babu KS, Mahendran B. Morphological and chemo profile (Liquid chromatography mass spectroscopy and Gas chromatography and mass spectroscopy) comparisons of *Cyperus scariosus* R.Br and *Cyperus rotundus* L. *Pharmacognosy Magazine* 2015;11:439-47.
22. Tiwari AK, Reddy KS, Radhakrishnan J, Kumar DA, Zehra A, Agawane SB. Influence of antioxidant rich fresh vegetable juices on starch induced postprandial hyperglycemia in rats. *Food and Function* 2011;2:521-28.
23. Reddy SD, Siva B, Poornima B, Kumar DA, Tiwari AK, Ramesh U, et al. New free radical scavenging neolignans from fruits of *Piper attenuatum*. *Pharmacogn Mag* 2015;11:235-41.
24. Kahlenberg JM, Dubyak GR. Differing caspase-1 activation states in monocyte versus macrophage models of IL-1 $\beta$  processing and release. *J Leukocyte Biology* 2004;76:676-84.
25. Winter CA, Risley EA, Nuss GW. Carrageenan induced edema in hind paw of the rats-an assay for anti-inflammatory drugs. *Proc Soc Exp Biol Med* 1962;11:544-7.

26. Galipalli S, Patel NK, Prasanna K, Bhutani KK. Activity guided investigation of *Carissa carandas* (L.) roots for anti-inflammatory constituents. *Nat Prod Res* 2015;29:1670-2.
27. Deore SL, Khadabadi SS. Isolation and characterization of phytoconstituents from *Chlorophytum borivilianum*. *Pharmacognosy Research* 2010;2:343-9.
28. Yuandani Jantan I, Ilangkovan M, Husain K, Chan KM. Inhibitory effects of compounds from *Phyllanthus amarus* on nitric oxide production, lymphocyte proliferation and cytokine release from phagocytes. *Drug Des Devel Ther* 2016;10:1935-45.
29. Tao YW, Xu WD, Tian Y, Shi JG. Steroids and aromatic derivatives from *Euphorbia micractina*. *Zhongguo Zhong Yao Za Zhi* 2015;40:4639-44.
30. Dosoky NS, Moriarity DM, Setzer WN. Phytochemical and biological investigations of *Conradina canescens*. *Nat Pro Commun* 2016;11:25-8.
31. Kuete V, Nana F, Ngameni B, Mbaveng AT, Keumedjio F, Ngadjui BT. Anti microbial activity of the crude extract, fractions and compounds from stem bark of *Ficus ovata* (Moraceae). *J Ethnopharmacol* 2009;124:556-61.
32. Arambewela LS, Kumaratunga A, Arawwawala M, Owen NL, Du L. Volatile oils of *Alpinia calcarata* grown in Sri Lanka. *J Essent Oil Res* 2005;17:124-5.
33. Jung J, Lee JH, Bae KH, Jeong CS. Anti gastric actions of eugenol and cinnamic acid isolated from *Cinnamomi ramulus*. *Yakugaku Zasshi* 2011;131:1103-10.
34. Lincha VR, Zhao BT, Woo MH, Yang IJ, Shin HM. Effects of constituent compounds of *Smilax china* on nicotine-induced endothelial dysfunction in human umbilical vein endothelial cells. *Bio Pharm Bull* 2016;39:984-92.
35. Lin Z, Chen L, Qiu Q, Guo S. Isolation and identification of anti proliferative compounds from the roots of *Tetragium hemsleyanum* against MDA-MB-435S cell lines. *Pak J Pharm Sci* 2016;29:1171-5.
36. Durackova Z. Some current insights into oxidative stress. *Physiol Res* 2010;59:459-69.
37. Zhu D, Sanchez-Ferrer A, Nystrom L. Antioxidant activity of individual steryl ferulates from various cereal grain sources. *J Nat Prod* 2016;79:308-16.
38. Zafar S, Ahmed R, Khan R. Biotransformation; a green and efficient way of antioxidant synthesis. *Free Radic Res* 2016;27:1-10.
39. De souze E, Zanatta L, Seifriz I, Creczynski-Pasa TB, Pizzolatti MG. Hypoglycemic effect and antioxidant potential of kaempferol-3-7-O-a-di rhamnoside from *Bauhinia forficata* leaves. *Journal of Natural Products* 2004;67:829-32.
40. Hanamura T, Hagiwara T, Kawagishi H. Structural and functional characterization of polyphenol isolated from acerola (*Malpighia emarginata* DC.) fruit. *Bioscience, Biotechnology, and Biochemistry* 2005;69:280-86.
41. Gabay O, Sanchez C, Salvat C, Chevy F, Breton M, Nourissat G. et al. Stigmasterol a phytosterol with potential anti-osteoarthritic properties. *Osteoarthritis and cartilage* 2010;18:106-16.
42. Fernandez MA, De las Heras B, Garcia MD, Saenz MT, Villar A. New insights into the mechanism of action of the anti-inflammatory triterpene lupeol. *The Journal of Pharmacy and Pharmacology* 2001;53:1533-39.
43. Sudhakar V, Ashok Kumar S, Varalakshmi P, Sujatha V. Protective effect of lupeol and lupeol linoleate in hypercholesterolemia associated renal damage. *Molecular and Cellular Biochemistry* 2008;317:11-20.



**Dr. Mahendran Botlagunta**

#### ABOUT AUTHOR

**Dr. Mahendran Botlagunta**, working as a Associate Professor in the department of Biotechnology, K L E F University, Guntur, Andhra Pradesh. His research focuses on interdisciplinary areas of cancer, Inflammation, nanomedicine and bioinformatics. His works are all well reported in international journals.