

Decoction of *Dictamnus Dasycarpus* Turcz. Root Bark Ameliorates Skin Lesions and Inhibits Inflammatory Reactions in Mice with Contact Dermatitis

Beodeul Yang, Hong-Bum Lee, Sura Kim, Young Chul Park¹, Koanhoi Kim², Hyungwoo Kim

Division of Pharmacology, School of Korean Medicine, Pusan National University, Gyeongnam, ¹Department of Microbiology and Immunology, School of Medicine, Pusan National University, Gyeongnam, ²Department of Pharmacology, School of Medicine, Pusan National University, Gyeongnam, Korea

Submitted: 25-07-2016

Revised: 08-09-2016

Published: 19-07-2017


ABSTRACT

Background: The root bark of *Dictamnus dasycarpus* Turcz. (Dictamni Radicis Cortex) has been widely used to treat skin diseases in Korea, and its anti-inflammatory efficacies were recently reported. **Objective:** The paper aims to investigate the inhibitory effects of decoction of Dictamni Radicis Cortex (DDRC) in mice with contact dermatitis (CD). **Materials and Methods:** We investigated the effects of DDRC on skin lesion characteristics such as crust, scales, incrustation and petechiae, the erythema and melanin indexes, skin thickness, histopathologic changes, and cytokine production in 1-fluoro-2,4-dinitrofluorobenzene (DNFB)-induced CD mice. **Results:** Topical application of DDRC ameliorated crust, scales, incrustation, and induced by DNFB. In addition, DDRC lowered the erythema index significantly ($P < 0.05$). DDRC effectively inhibited enlargement of skin thickness ($P < 0.05$). Histopathologic observation showed that DDRC inhibited epidermal hyperplasia, hyperkeratosis, and spongiotic changes. Finally, DDRC decreased production levels of IFN- γ , TNF- α and IL-6 induced by repeated application of DNFB ($P < 0.05$). **Conclusion:** These data suggest that DDRC can be used in the treatment of inflammatory skin diseases including CD. Moreover, these results are closely related to the decreasing production of TNF- α , IFN- γ and IL-6 in inflamed tissues.

Key words: *Dictamnus dasycarpus*, traditional Chinese medicine, inflammation, contact dermatitis

SUMMARY

- DDRC ameliorated skin lesions such as crust, scales, incrustation and petechiae, and lowered erythema index on skin surface in CD mice
- DDRC inhibited enlargement of dorsal skin and prevented epidermal hyperplasia, hyperkeratosis, and spongiotic changes in inflamed tissues
- DDRC reduced the levels of TNF- α , IFN- γ , and IL-6 in inflamed tissues of CD mice
- DDRC did not affect spleen/body weight ratio in CD mice.



Abbreviations used: DDRC: decoction of Dictamni Radicis Cortex, CD: contact dermatitis, DNFB: 1-fluoro-2,4-dinitrofluorobenzene, AOO: acetone and olive oil, DEX: dexamethasone, CBA: cytometric bead array

Correspondence:

Prof. Hyungwoo Kim, Division of Pharmacology, School of Korean Medicine, Pusan National University, Yangsan, Gyeongnam, Korea.
E-mail: kronos7@pusan.ac.kr
DOI: 10.4103/0973-1296.211034

Access this article online

Website: www.phcog.com

Quick Response Code:



INTRODUCTION

Contact dermatitis (CD), which affects roughly 20% of the population in the United States, is the most general form of occupational skin disease.^[1,2] CD has great socio-economic impact and can lead to decreased work performance and quality of life.^[3] CD is a T-cell-mediated type IV hypersensitivity in which exposure to allergens or sensitization is followed by inflammation on the second exposure, and therefore, allergic reaction occurs.^[4] Skin diseases including CD can be caused by both external and internal factors. External factors can be work environment, cosmetics, or other personal care products, while internal factors are psychological and genetic.^[3,5] Although external factors are the major cause of CD, psychological and genetic factors also play a role by enhancing contact hypersensitivity.^[5] Although the exact cause of allergic reaction is unknown, several of its mechanisms have been elucidated. Type IV hypersensitivity is highly associated with mast cells,

which act as effector cells by secreting pro-inflammatory substances such as histamines, interleukin (IL), tumor necrosis factor (TNF)-alpha, and interferon (IFN)-gamma.^[4-6] CD results in increased dermal weight and

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Yang B, Lee HB, Kim S, Park YC, Kim K, Kim H. Decoction of *Dictamnus Dasycarpus* Turcz. Root bark ameliorates skin lesions and inhibits inflammatory reactions in mice with contact dermatitis. Phcog Mag 2017;13:483-7.




Figure 3: Effect of DDRC on erythema and melanin indexes on skin surface in CD mice. The erythema and melanin indexes were measured using a dermo-spectrophotometer on day 15. (A) Erythema index; (B) melanin index. NOR: nontreated normal mice, CTL: nontreated CD mice, DDRC: 60 μ g/day of DDRC-treated CD mice, DEX: 30 μ g/day of dexamethasone-treated CD mice. All values are presented as the mean \pm SD. [#] $P < 0.05$ vs. nontreated normal mice (NOR). ^{*} $P < 0.05$ vs. nontreated CD mice (CTL).




Figure 4: Effect of DDRC on skin thickness in CD mice. The skin thickness was measured using Vernier calipers on day 15. Abbreviations are the same as in [Figure 3]. All values are presented as the mean \pm SD. [#] $P < 0.05$ vs. nontreated normal mice (NOR). ^{*} $P < 0.05$ and vs. nontreated CD mice (CTL).

Measurement of dorsal skin thicknesses

After observing skin lesions and measuring the erythema and melanin index, the dorsal skins were resected and their thicknesses measured using Vernier calipers (Mitutoyo, Tokyo, Japan).

Histopathologic examination

Tissues obtained from experimental animals were fixed in 4% neutral formalin overnight, then washed, dehydrated with ethyl alcohol sequentially, treated with xylene and embedded in paraffin. Skin tissues (4 μ m) were then resected, after which sections were stained with hematoxylin and eosin (H and E) and observed using a light microscope (200 \times).

Measurement of cytokine production

Cytokine levels in inflamed tissues were measured according to the cytometric bead array (CBA) method using a Mouse Inflammation CBA kit (BD Biosciences, San Jose, CA, USA). At the final step of the experiment, resected dorsal skin tissues were lysed and homogenized with protein extraction solution (Intron Bio, Daejeon, Korea) using a bullet blender (Next Advance, NY, USA) to obtain tissue lysates. Fifty micrograms of each lysate was then used to measure the levels




Figure 5: Effects of DDRC on histopathologic changes in CD mice. Skin tissues were stained with H and E and observed using a light microscope. (A) NOR; (B) CTL; (C) 60 μ g/day of DDRC; (D) 30 μ g/day of DEX. Yellow bars indicate the vertical length of the epidermis. Filled arrows indicate the hyperkeratotic area. All observations were made at a magnification of 200 \times .

of interferon-gamma (IFN- γ), tumor necrosis factor-alpha (TNF- α), interleukin-6 (IL-6), and interleukin-10 (IL-10). All experimental procedures were conducted according to the manufacturer's protocols.

Statistical analysis

A Student's t-test was used for all statistical comparisons, and Prism 5 version 5.01 (Graph Pad Software Inc., La Jolla, CA, USA) was used for all analyses. All data are presented as the mean \pm SD. P less than 0.05 were considered significant.

RESULTS

DDRC ameliorated skin lesions in CD mice

DNFB treatment induced skin lesions of CD such as crust, scale, incrustation, petechiae, and pigmentation in the CTL group [Figure 2B]. Treatment with DDRC ameliorated these symptoms and decreased the crust, scale, and incrustation [Figure 2C].

DDRC lowered erythema index on skin surface in CD mice

A marked increase in erythema index was observed in the CTL group. Treatment with DDRC reduced the erythema index significantly, while DEX did not [Figure 3A]. Conversely, neither DDRC nor DEX affected the melanin index [Figure 3B].

DDRC inhibited enlargement of dorsal skin in CD mice

The inhibitory effects of DDRC on swelling and enlargement of skin tissues were evaluated by measuring the effects on skin thicknesses in mice with CD. Repeated application of DNFB increased the levels of skin thickness to more than twice those observed in the normal group. These increases were effectively inhibited by DDRC and DEX [Figure 4].




Figure 6: Effects of DDRC on cytokine production in CD mice. The levels of TNF- α , IFN- γ , IL-6 and IL-10 in skin tissues were measured using the cytometric bead array method. A total of 50 μ g of tissue lysates were used to measure the cytokine levels. Abbreviations are the same as those used in [Figure 3]. (A) TNF- α ; (B) IFN- γ ; (C) IL-6; (D) IL-10. All values are presented as the mean \pm SD. [#] $P < 0.05$ and ^{###} $P < 0.001$ vs. nontreated normal mice (NOR). ^{*} $P < 0.05$ vs. nontreated CD mice (CTL).




Figure 7: Effects of DDRC on spleen/body weight ratio in CD mice. Body weights and spleen weights were measured on day 15, at which time the spleen/body weight ratio was calculated. Abbreviations are the same as in [Figure 3]. All values are presented as mean \pm SD. [#] $P < 0.05$ and ^{##} $P < 0.01$ vs. nontreated normal mice (NOR). ^{**} $P < 0.01$ vs. nontreated CD mice (CTL).

DDRC prevented epidermal hyperplasia, hyperkeratosis, and spongiotic changes in inflamed tissues

There were no abnormal changes in the normal group [Figure 5A]. Repeated painting with DNFB induced significant epidermal hyperplasia (yellow bar), as well as hyperkeratosis (filled arrow), which are hallmarks of skin inflammation. In addition, spongiotic changes were observed in the CTL group [Figure 5B], while DDRC treatment inhibited epidermal hyperplasia, hyperkeratosis, and spongiotic changes [Figure 5C]. Treatment with DEX most effectively prevented abnormal changes in inflamed tissues of CD mice [Figure 5D].

DDRC reduced the levels of TNF- α , IFN- γ , and IL-6 in inflamed tissues of CD mice

Marked increases in TNF- α , IFN- γ , IL-6 and IL-10 production were observed in the CTL group. These increases in TNF- α , IFN- γ , and IL-6 production were effectively reduced by topical application of DDRC. Neither DDRC nor DEX affected the IL-10 level [Figure 6].

DDRC did not affect spleen/body weight ratio in CD mice

The effects of DDRC on enlargement of the spleen were estimated based on the spleen/body weight ratio. The spleen/body weight ratio in the CTL group doubled compared with the NOR group. The DDRC group showed no change in spleen/body weight ratio in CD mice. However, when compared with the CTL and NOR groups, this ratio was significantly decreased in the DEX group [Figure 7].

DISCUSSION

In this experiment, sensitization with DNFB and increased concentration of DNFB for challenge led to increased susceptibility to CD. Sensitization

of the ears induced activation of immune cells, so synthesis of pro-inflammatory factors such as cytokines and chemokines could occur. The immune cells activated in the initiation phase then circulate throughout the body with lymph and blood circulation.^[12] During the elicitation phase, effector T cells from lymph nodes and blood were recruited at the challenged site.^[4] The animal model used in this experiment was designed to mimic human CD, especially allergic contact dermatitis. For this reason, mice were sensitized on days 1, 2, and 3 (initiation phase) and challenged on days 7, 9, 11, and 13 (elicitation phase) on different sites (ear and shaved back).

Dorsal skin appearance data show that both DDRC and DEX relieved crust, scales, incrustation, and petechiae compared with the control group [Figure 2]. The erythema index was distinctly higher in all challenged groups than in the normal group, and only DDRC led to a significant reduction in the erythema index [Figure 3A]. Conversely, the melanin index remained unchanged [Figure 3B]. In addition, the control group had the thickest skin due to inflammation, whereas the DDRC and DEX groups had thinner skin than the control group, with the skin of the DEX group being slightly thinner than that of the normal group [Figure 4]. Although these data indicate that DDRC and DEX significantly relieved skin inflammation, the results observed for DEX indicate that corticosteroid use results in a thinner skin barrier. This results in aesthetic and functional problems, with vessels underneath the skin becoming more visible. Moreover, the thinner skin barrier makes it more vulnerable to other foreign substances. Notably, topical application of corticosteroids to relieve allergic reaction has been shown to result in skin atrophy,^[13] as has steroid injection.^[14] Therefore, patients with dermal inflammation may need to compromise or change their treatment to alternative or complementary agents.

The control groups had thicker layers because of their inflammatory response, which was a result of keratinocyte proliferation.^[15] In addition, repeated application of DNFB induced spongiotic changes, which are a hallmark of inflammatory skin diseases and large vesicles in control group [Figure 5B]. The histopathologic changes observed in the present study indicate that both DDRC and DEX prevented epidermal hyperplasia, hyperkeratosis, and spongiotic changes induced by DNFB [Figure 5]. In CD, keratinocytes can be activated by TNF- α and IFN- γ and play a major role as hapten-presenting cells associated with CD8⁺ T cells.^[4,16] In addition,

IL-6 is known to induce the proliferation and migration of keratinocytes, leading to skin diseases such as psoriasis.^[17] Once activated, keratinocytes secrete cytokines including IL-1, IL-3, IL-6, IL-8, and TNF- α .^[18,19] As shown in Figure 6, there were marked increases in TNF- α , IFN- γ , IL-6, and IL-10 in the control group. DDRC had an inhibitory effect on TNF- α , IFN- γ , and IL-6 production, indicating that DDRC acts as an anti-inflammatory agent in skin tissue. Considering the proliferative effect of TNF- α , IFN- γ , and IL-6, these data indicate that DDRC can inhibit epidermal hyperplasia via regulation of pro-inflammatory cytokines such as TNF- α , IFN- γ , and IL-6 in inflamed tissues.

The spleen/body weight ratio of the control group was almost double that of the normal group. The DDRC group showed almost the same level as that of the control group, but the DEX group showed a reduced weight ratio relative to the normal group, indicating that corticosteroids led to immune suppression Figure 7. Based on these data, the continuous use of corticosteroid can lead to immune suppression and bring other unexpected side effect. It has also been reported that dexamethasone induces spleen atrophy via the apoptosis of splenocytes.^[20] Conversely, DDRC did not affect spleen/body weight ratio, implying that the action mechanisms of DDRC are somewhat different from those of corticosteroids, especially in the framework of general immune weakness.

Taken together, these data indicate that DDRC can prevent epidermal hyperplasia, hyperkeratosis, and spongiotic changes via regulation of the production levels of pro-inflammatory cytokines. In addition, these anti-inflammatory actions of DDRC inhibit increases in skin thickness and ameliorate skin lesions and the erythema index.

CONCLUSIONS

We demonstrated the anti-inflammatory effects of DDRC in CD mice in this study. DDRC inhibited the production of TNF- α , IFN- γ , and IL-6, resulting in reduced epidermal hyperplasia, hyperkeratosis, and spongiotic changes. These anti-inflammatory reactions of DDRC finally prevented enlargement of skin thickness and ameliorated skin lesions. However, in contrast to dexamethasone, DDRC had no effect on the spleen/body weight ratio. Overall, these results indicate that DDRC can be used to treat patients with inflammatory skin diseases with relative safety.

Financial support and sponsorship

This research was supported by a National Research Foundation of Korea grant funded by the Korean government (MSIP; grant no. 2015R1A2A2A04005619 and 2014R1A5A2009936).

Conflict of interest

There are no conflicts of interest

REFERENCES

1. Mathias CG. The cost of occupational skin disease. *Arch Dermatol* 1985;121:332-4.
2. Diepgen TL. Occupational skin diseases. *J Dtsch Dermatol Ges* 2012;10:297-313.
3. Bathe A, Diepgen TL, Mattered U. Subjective illness perceptions in individuals with occupational skin disease: a qualitative investigation. *Work* 2012;43:159-69.
4. Saint-Mezard P, Krasteva M, Chavagnac C, Bosset S, Akiba H, Kehren J, *et al.* Afferent and efferent phases of allergic contact dermatitis (ACD) can be induced after a single skin contact with haptens: evidence using a mouse model of primary ACD. *J Invest Dermatol* 2003;120:641-7.
5. Kaplan DH, Igyplan BZ, Gaspari AA. Early immune events in the induction of allergic contact dermatitis. *Nat Rev Immunol* 2012;12:114-24.
6. Kim H, Kim M, Kim H, Lee GS, An WG, Cho SI. Anti-inflammatory activities of *Dictamnus dasycarpus* Turcz., root bark on allergic contact dermatitis induced by dinitrofluorobenzene in mice. *J Ethnopharmacol* 2013;149:471-7.
7. Lyons JJ, Milner JD, Stone KD. Atopic dermatitis in children: clinical features, pathophysiology, and treatment. *Immunol Allergy Clin North Am* 2015;35:161-83.
8. Wen M, Wei C, Hu Z, Srivastava K, Ko J, Xi S, *et al.* Efficacy and tolerability of anti-asthma herbal medicine intervention in adult patients with moderate-severe allergic asthma. *J Allergy Clin Immunol* 2005;116:517-24.
9. Kim CM, Shin MG, An DG, Lee KS. *The Encyclopedia of Oriental Herbal Medicine*. Seoul: Jeongdam 1997;1648-51.
10. The Textbook Compilation Committee of Korean herbology *Bonchohak*. Seoul: Younglimsa 2010;228-9.
11. Yoon JS, Sung SH, Kim YC. Neuroprotective limonoids of root bark of *Dictamnus dasycarpus*. *J Nat Prod* 2008;71:208-11.
12. Morita H, Yamagata M, Inohara S, Sagami S. Flow cytometric analysis of protein kinase C in lymph node cells during the induction phase of contact hypersensitivity reaction. *J Dermatol* 1989;16:352-4.
13. Klopot A, Baida G, Bhalla P, Haegeman G, Budunova I. Selective activator of the glucocorticoid receptor compound a dissociates therapeutic and atrophogenic effects of glucocorticoid receptor signaling in skin. *J Cancer Prev* 2015;20:250-9.
14. van Vendeloo SN, Etema HB. Skin depigmentation along lymph vessels of the lower leg following local corticosteroid injection for interdigital neuroma. *Foot Ankle Surg* 2016;22:139-41.
15. Cesinaro AM, Nannini N, Migaldi M, Pepe P, Maiorana A. Psoriasis vs allergic contact dermatitis in palms and soles: a quantitative histologic and immunohistochemical study. *APMIS* 2009;117:629-34.
16. Sosroseno W. The immunology of nickel-induced allergic contact dermatitis. *Asian Pac J Allergy Immunol* 1995;13:173-81.
17. Kawakami M, Kaneko N, Anada H, Terai C, Okada Y. Measurement of interleukin-6, interleukin-10, and tumor necrosis factor-alpha levels in tissues and plasma after thermal injury in mice. *Surgery* 1997;121:440-8.
18. Albanesi C. Keratinocytes in allergic skin diseases. *Curr Opin Allergy Clin Immunol* 2010;10:452-6.
19. Kim J, Krueger JG. The immunopathogenesis of psoriasis. *Dermatol Clin* 2015;33:13-23.
20. Nakao R, Yamamoto S, Yasumoto Y, Oishi K. Dosing schedule-dependent attenuation of dexamethasone-induced muscle atrophy in mice. *Chronobiol Int* 2014;31:506-14.