

# Cytotoxic, Antimitotic, and Antiproliferation Studies on *Rasam*: A South Indian Traditional Functional Food

Agilandeswari Devarajan<sup>1,2</sup>, M. K. Mohan Maruga Raja<sup>2</sup>

<sup>1</sup>Department of Pharmaceutics, Hillside College of Pharmacy and Research Centre, Bengaluru, Karnataka, <sup>2</sup>Department of Pharmacy, Centre for Research and Development, PRIST University, Thanjavur, Tamil Nadu, India

Submitted: 11-04-2017

Revised: 18-05-2017

Published: 11-10-2017


## ABSTRACT

**Background:** *Rasam* is a traditional South Indian food, prepared using tamarind juice as a base, with a variety of spices. *Rasam*, with all its ingredients medicinally claimed for various ailments, is a functional food. Systematic consumption of traditional functional food provides an excellent preventive measure to ward off many diseases. **Objective:** To study *rasam* for cytotoxic, antimitotic, and antiproliferation potential beyond its culinary and nutritional effect. **Materials and Methods:** Brine shrimp lethality assay, onion root tip inhibition assay, and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay in Calu-6, HeLa, MCF-7 cell lines for four stage-wise samples in the preparation of *rasam* (RS1, RS2, RS3, and RS4) were studied. **Results:** RS4, the end product of *rasam* showed high lethality with an LC<sub>50</sub> value of 38.7 µL/mL. It showed maximum antimitotic activity in a dose-dependent manner compared to other samples with an IC<sub>50</sub> value of 189.86 µL/mL. RS4 also showed an IC<sub>50</sub> value of 350.22 and 410.15 µL/mL in MCF-7 and Calu-6 cell lines, respectively. **Conclusion:** From this study, we suggest that *rasam* is a classic example of traditional functional food and it can treat breast and lung cancer on chronic use.

**Key words:** Brine shrimp lethality, Calu-6, HeLa, MCF-7, onion root tip inhibition, *Saaru*

## SUMMARY

- *Rasam*, a South Indian traditional functional food, showed high lethality (LC<sub>50</sub> = 38.7 µL/mL) against brine shrimps
- *Rasam* also showed potential antimitotic activity (IC<sub>50</sub> = 189.86 µL/mL) by inhibiting the onion root tips
- *Rasam* showed an IC<sub>50</sub> value of 350.22 and 410.15 µL/mL against MCF-7 and Calu-6 cell lines respectively
- *Rasam*, when consumed on daily dietary basis, can treat breast and lung cancer.



**Abbreviations used:** SS 316: Stainless Steel 316 grade; MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; DMEM: Dulbecco's modified Eagle medium; FBS: Fetal bovine serum media; TPVG: Trypsin phosphate versene glucose; EDTA: Ethylene diamine tetra acetic acid; PBS: Phosphate buffered saline; DMSO: Dimethyl sulfoxide.

## Correspondence:

Prof. Agilandeswari Devarajan,  
Department of Pharmaceutics, Hillside  
College of Pharmacy and Research Centre,  
#9, Raguhuvanahalli, Kanakapura Main Road,  
Bengaluru - 560 062, Karnataka, India.  
E-mail: agilandeswaridevarajan@gmail.com  
**DOI:** 10.4103/pm.pm\_138\_17

## Access this article online

Website: www.phcog.com

## Quick Response Code:



## INTRODUCTION

The view that food can have an expanded role that goes well beyond providing a source of nutrients truly applies to traditional functional foods. The systematic consumption of traditional functional food provides an excellent preventive measure to ward off many diseases. Epidemiological randomized clinical trials carried out in different countries have demonstrated numerous health effects related to functional food consumption such as reduction of cancer risk, improvement of heart health, stimulation of immune system, decrease of menopause symptoms, improvement of gastrointestinal health, maintenance of urinary tract health, anti-inflammatory effects, reduction of blood pressure, maintenance of vision, antibacterial effect, antiviral effect, reduction of osteoporosis, and anti-obese effect.<sup>[1]</sup> Traditional functional foods can help prevent chronic disease or optimize health, therefore reducing health-care costs and improving the quality of life.

Spices play very important role in digestive function and the Indian tradition has a long history of use of spice in food as medicines to

prevent and treat diseases.<sup>[2]</sup> Another epidemiological study suggested that curcumin, the bioactive compound of turmeric, as one of the most prevalent nutritional and medicinal compounds used by the Indian population, is responsible for the significantly reduced (4.4 times) prevalence of cardiovascular diseases, metabolic diseases, neurodegenerative diseases, and cancer in India compared to the United States of America.<sup>[3]</sup> It is also estimated that an adult in India consumes 80–200 mg/day of curcumin and 50 g of garlic in 1 week. Hence, there

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

**Cite this article as:** Devarajan A, Mohan Maruga Raja MK. Cytotoxic, antimitotic, and antiproliferation studies on *Rasam*: A South Indian traditional functional food. *Phcog Mag* 2017;13:S452-7.

is a realistic possibility to reach a therapeutic dose by daily dietary consumption.<sup>[4,5]</sup> The whole world realized only in the 20<sup>th</sup> century that food plays a major role in disease prevention, but centuries before ancient India has realized the importance of food in health and wellness. *Rasam* is a very popular South Indian traditional spice soup. It is consumed on daily basis in every South Indian home. It is also called as *rasam* or *chaaru* or *saaru* in different South Indian languages. In a traditional South Indian meal, *rasam* is preceded by a *sambar* rice course and is followed by curd rice. *Rasam* is traditionally prepared using tamarind juice as a base, with a variety of spices which are considered to be good for health.<sup>[6]</sup> The main spices used in *rasam* preparation are coriander, garlic, curry leaves, tamarind, cumin, black pepper, mustard, turmeric, red chili, and asafetida.<sup>[7]</sup> *Rasam* is a functional food because all ingredients used in the preparation of *rasam* are medicinally claimed for various ailments. *Sambar*, another South Indian traditional dish, has shown preventive effect against colon cancer.<sup>[8]</sup> There is a need to understand traditional systems and visualize the future of medicine and health care. The linkage between “the past” and “the future” of medicine is much more important and can give us “new directions” for better understanding health, disease, and possible solutions.<sup>[9]</sup>

A study on *rasam*, which is being consumed from time immemorial, is only an approach of “drug rediscovery.” In view of all the above facts, *rasam* was studied for cytotoxic, antimitotic, and antiproliferation potential beyond its culinary and nutritional effect.

## MATERIALS AND METHODS

### Materials

All ingredients of *rasam* were purchased from Arokya Organic Shop, Vellore, Tamil Nadu. All utensils used for the preparation of *rasam* were of Stainless Steel 316 grade (SS 316). Brine shrimp eggs were purchased from Ocean Star International Inc., Snowville, UT, USA. Onion bulbs were purchased from Nutrisiree Organics, Bengaluru, Karnataka. MCF-7 (ATCC HTB-22, passage number 11), HeLa (ATCC CCL2, passage number 13), CALU-6 (ATCC HTB-56, passage number 19) cell lines were procured. 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reagent, Dulbecco's modified Eagle medium (DMEM), fetal bovine serum (FBS) media, trypsin phosphate versene glucose (TPVG) solution, ethylene diamine tetra acetic acid (EDTA), phosphate buffered saline (PBS), dimethyl sulfoxide (DMSO), trypsin, acetocarmine, and cedar wood oil were obtained from Sigma-Aldrich (Bengaluru, India) and HiMedia Ltd., (Mumbai, India). All other chemicals and solvents were obtained from SD Fine Chemicals (Mumbai, India) and were of analytical grade.

### Preparation of *rasam*

*Rasam* was prepared in five stages as mentioned below.

1. Tamarind fruit pulp mixture (T1): 6.88 g of tamarind fruit pulp was immersed in 450 mL of water for 10 min, and then it was hand crushed for 45 times and strained. The strained liquid was rinsed with 5 mL water, to which 0.4 g of turmeric powder and 4 g of sea salt were added
2. Tomato fruit mixture (T2): 82.44 g of fresh tomato fruits were cut and hand crushed for 60 times. The crushed fruit was rinsed with 5 mL of water
3. Spice mixture (T3): 1.33 g of pepper drupes was crushed in an SS 316 mortar and pestle for 85 times. 2.67 g of cumin fruits was added over to the crushed pepper drupes and crushed for 100 times. To the above-crushed mixture, 0.82 g of chili pepper was added and crushed for 50 times; then, 9.63 g of garlic cloves was added and crushed for 90 times
4. All mixture (T4): Tomato fruit mixture (T2) was rinsed with 10 mL of water and spice mixture (T3) was rinsed with 10 mL of water. Both

rinsing were added to tamarind fruit pulp mixture (T1), which was designated as sample RS1

5. Final product (T5): 4 ml of Indian sesame oil was heated at 60°C for 2 min. After 5 seconds 0.82 g of mustard seeds were added. After 3 s, 1.53 g of whole chili pepper was added. After 2 s, 0.61 g of curry leaves was added, which was designated as sample RS2. Immediately, all mixture (T4) was rinsed with 20 mL of water and added. The whole liquid was allowed to boil for 5 min. After 5 min, 1.50 g of coriander leaves was added; this was designated as sample RS3. When the liquid frothed, 0.05 g of asafetida was added and the heating was switched off to yield the final product, which was designated as sample RS4.

The stage-wise samples RS1, RS2, RS3, and RS4 in the preparation of *rasam* were studied to evaluate the significance of the traditional processing.

### Cytotoxicity study

Cytotoxicity was studied by brine shrimp lethality bioassay as per the method of Meyer *et al.*, 1982.<sup>[10]</sup> Brine shrimps (*Artemia salina*) were hatched from eggs in a conical-shaped vessel (1 L), filled with artificial seawater (prepared using sea salt 38 g/L and adjusted to pH 8.5 using 1 N NaOH) with constant aeration for 36 h at room temperature (20°C ± 5°C) under light. After hatching, active nauplii, free from egg shells, were collected from brighter portion of the hatching chamber and used for the assay. Ten nauplii were drawn through a glass capillary and placed in vials each containing 4.5 mL of brine solution (24% of NaCl in water). In every vial, 0.5 mL of water or sample was added to the brine solution and maintained at room temperature (20°C ± 5°C) under light. The number of surviving nauplii after 24 h was counted. Experiments were conducted at different concentrations (20, 40, 80, 120, 160, 200, and 400 µL/mL) of the samples (RS1, RS2, RS3, and RS4) diluents being water. Each concentration of the sample was studied in six vials along with a control group.

The %lethality was determined from the number of surviving nauplii in control and sample using the below-mentioned formula.

$$\% \text{ lethality} = \frac{\text{Control} - \text{Sample}}{\text{Control}} \times 100$$

Control is number of surviving nauplii in control, and sample is number of surviving nauplii in sample. LC<sub>50</sub> values were calculated from percentage lethality versus concentration best-fit line graph.

### Antimitotic study

Antimitotic study was studied by onion root tip inhibition assay as per the method of Rai *et al.*, 2007.<sup>[11]</sup> The old roots of *Allium cepa* bulbs (56.46 ± 4.14 g; values are expressed as mean ± standard deviation [SD]) were removed and grown in the dark over a small beaker containing 35 mL of tap water (the water was changed every 24 h) at 20°C ± 5°C until the root tips have grown to approximately 2–3 cm (in approximately 2–3 days). The bulbs with root tips grown up to 2–3 cm were selected. Ten root tips grown above 2 cm were kept as such, and rest of them were trimmed off. The bulbs with only 10 root tips were placed over water or sample solutions, and incubation was carried out at 20°C ± 5°C. Samples RS1, RS3, and RS4 each at a concentration of 1, 10, 100, and 200 µL/mL were studied in triplicate. The oil sample (RS3) would form a layer on the surface and never allow the growth of the root tip; hence, RS3 was omitted for this study.

In each bulb, root length and newly grown root tips were recorded at 0, 24, 48, and 72 h. One root in each bulb was cut using a scalpel between 8.00 to 13.00 h IST at 0, 24, 48, and 72 h. Extreme root tips (2–3 mm) of root were cut and put in a test tube containing water. The test tube was

treated with 1.5 mL of 0.1 N HCl and incubated at 60°C for 12 min. The 0.1 N HCl was drained and the root tips were washed with water 4 times to remove the acid traces. Then, five drops of acetocarmine was added and incubated at 60°C for 24 min. The red turned root tip was transferred to a clean glass slide and a cover slip was placed over. Very gentle pressure was applied with thumb over the cover slip to provide uniform spread of the cells. The number of mitotic and total meristematic cells was counted in 5–6 fields using high power ( $\times 100$ ) light microscope. For control and all samples, 500 cells were counted and cells manifesting different stages of mitosis, i.e., interphase and prophase (P), metaphase (M), anaphase (A), and telophase (T), were recorded. The mitotic index at 0, 24, 48, and 72 h was calculated using the following formula.

$$\text{Mitotic index} = (P + M + A + T) / (\text{Total number of cells}) \times 100$$

P, M, A, and T is prophase, metaphase, anaphase, and telophase, respectively.

IC<sub>50</sub> values were calculated from the concentration versus percentage of inhibition best-fit line graph. The values are expressed as mean  $\pm$  SD. GraphPad Instat Version 4 software (GraphPad Software, USA) was used. Data were subjected to the one-way analysis of variance to determine the significance of changes followed by Dunnett's multiple comparisons.

### Antiproliferation study

MCF-7, HeLa, and Calu-6 stock cells were cultured in DMEM supplemented with 10% inactivated FBS, penicillin (100 IU/mL), streptomycin (100  $\mu$ g/mL), and amphotericin B (5  $\mu$ g/mL) in humidified atmosphere of 5% CO<sub>2</sub> at 37°C until confluent. The cells were dissociated with TPVG solution (0.2% trypsin, 0.02% EDTA, 0.05% glucose in PBS). The viability of the cells was checked and centrifuged.

MTT assay was carried out by the modified method of Lau *et al.*, 2004.<sup>[12]</sup> The monolayer cell culture was trypsinized and the cell count was adjusted to  $1 \times 10^5$  cells/mL using DMEM containing 10% FBS. To each well of the 96-well microtiter plate, 100  $\mu$ L of the diluted cell suspension ( $5 \times 10^4$  cells/well) was added and incubated for 24 h at 37°C, 5% CO<sub>2</sub>. After 24 h, the supernatant was flicked off and the monolayer was washed with the medium once.<sup>[13]</sup> All samples were mixed with DMEM supplemented with 2% inactivated FBS to obtain different concentrations, ranging from 166.67 to 500  $\mu$ L/mL. Samples were added onto the partial monolayer in microtiter plates. The plates were then incubated at 37°C for 72 h in 5% CO<sub>2</sub> atmosphere. After 72 h, the test solutions in the wells were discarded and 50  $\mu$ L of MTT (5 mg/10 mL of MTT in PBS) was added to each well. The plates were gently shaken and incubated for 4 h at 37°C in 5% CO<sub>2</sub> atmosphere. The supernatant was removed, 100  $\mu$ L of DMSO was added, and the plates were gently shaken to solubilize the formed formazan. The absorbance was measured using a Tecan microplate reader at a wavelength of 590 nm. The percentage inhibition was calculated using the following formula.

$$\text{Percentage inhibition} = (\text{ABS of control} - \text{ABS of samples}) / (\text{ABS of control}) \times 100$$

ABS is absorbance and IC<sub>50</sub> values were calculated from the concentration versus percentage of inhibition best-fit line graph.

## RESULTS AND DISCUSSION

Brine shrimp lethality assay is a rapid, reliable and has been used for over 30 years in cytotoxic, phototoxic, pesticidal, trypanocidal, enzyme inhibition, and ion regulation activities.<sup>[14]</sup> The nauplii after hatching at 0 and 24 h are shown in Figure 1. RS4, the final product of *rasam*, showed high lethality (38.7  $\mu$ L/mL) than RS1 (165.6  $\mu$ L/mL), RS2 (387.6  $\mu$ L/mL), and RS3 (124.2  $\mu$ L/mL) [Table 1]. The percentage of lethality was found to be directly proportional to the concentration of the samples. Stagewise preparation analysis shows 3.2-fold increase of





Figure 1: Brine shrimp nauplii after hatching at 0 and 24 h

cytotoxicity in RS4 as compared to of RS1, RS2, and RS3. It is evident that the process in preparation of *rasam* plays an important role in increasing the physiological action of the final product (RS4). Brine shrimp lethality bioassay has good correlation with the human solid tumor cell lines;<sup>[15]</sup> hence, it can be suggested that the sample RS4 is bioactive, with cytotoxic and antitumor activity.

Most of the plant-derived anticancer drugs affect the microtubule dynamics of the cell and induce persistent modification of biological processes and signaling pathways that ultimately lead to apoptotic death.<sup>[16]</sup> Onion root tip inhibition assay is a quick, efficient, simple but sensitive antimitotic bioassay and also known to give a similar result to the *in vitro* animal cytotoxicity test.<sup>[17]</sup> The study showed that at 72 h, RS1, RS3, and RS4 at all concentrations (1, 10, 100, and 200  $\mu$ L/mL) significantly inhibits root growth [Table 2]. Figure 2 shows the effect of RS1 (200  $\mu$ L/mL), RS3 (200  $\mu$ L/mL), and RS4 (200  $\mu$ L/mL) on root tip growth at 72 h in comparison to the control group. It is evident that all samples affect cell division in long duration. There was no increase in root length for RS1 (200  $\mu$ L/mL) and RS3 (200  $\mu$ L/mL) after 24 h. The final product RS4 at 200  $\mu$ L/mL showed no increase in root length even after 0 h, suggesting maximum root inhibition. RS3 (200  $\mu$ L/mL) and RS4 (100 and 200  $\mu$ L/mL) significantly prevented the growth of new root tips after 0 h [Table 2]. However, after 72 h, all samples at different concentrations also significantly inhibited the growth of new root tips. RS4 (200  $\mu$ L/mL) not only significantly inhibits the root length but also prevents the growth of a new root tip after 0 h. The number of cells in prophase was nearly double the metaphase in RS1, RS3, and RS4 treated groups. The number of cells in telophase varied from 9 to 12 within the counted 500 cells of all sample-treated groups. The number of dividing cells affected and the mitotic index clearly confirms that RS4 (200  $\mu$ L/mL) at 24 h significantly inhibits mitosis [Table 3]. RS3 (100 and 200  $\mu$ L/mL) and RS4 (10 and 100  $\mu$ L/mL) showed lesser number of dividing cells and lower mitotic index after 72 h. The IC<sub>50</sub> value (189.86  $\mu$ L/mL) of the final product of RS4 (*rasam*) only confirms maximum inhibition of mitosis in a dose-dependent manner compared to RS1 and RS3 [Table 3], and also, the IC<sub>50</sub> value of RS4 was found to be 2.2 times lower than its constituents (RS1 and RS3). Antitumor drugs that interact with microtubules and tubulin are known to block mitosis and induce cell death by apoptosis.<sup>[18]</sup> Hence, it can be suggested that the





**Figure 2:** Effect of RS1, RS3, and RS4 on onion root tip growth at 72 h

**Table 1:** The lethality effect of RS1, RS2, RS3, and RS4 on brine shrimps

Control/sample	Concentration	Number of nauplii at 0 h	Number of surviving nauplii after 24 h	Percentage lethality	LC <sub>50</sub> (µL/mL)
Water	0.1 mL/mL	60	60	0	NA
RS1	20 µL/mL	60	58	3.3	165.6
	40 µL/mL	60	51	15.0	
	80 µL/mL	60	41	31.7	
	120 µL/mL	60	38	36.7	
	160 µL/mL	60	31	48.3	
	200 µL/mL	60	27	55.0	
RS2	400 µL/mL	60	16	73.3	387.6
	20 µL/mL	60	59	1.7	
	40 µL/mL	60	56	6.7	
	80 µL/mL	60	53	11.7	
	120 µL/mL	60	51	15.0	
	160 µL/mL	60	47	21.7	
RS3	200 µL/mL	60	44	26.7	124.2
	400 µL/mL	60	29	51.6	
	20 µL/mL	60	47	21.7	
	40 µL/mL	60	43	28.3	
	80 µL/mL	60	37	38.3	
	120 µL/mL	60	31	48.3	
RS4	160 µL/mL	60	26	56.7	38.7
	200 µL/mL	60	20	66.7	
	400 µL/mL	60	11	81.7	
	20 µL/mL	60	36	40.0	
	40 µL/mL	60	29	51.7	
	80 µL/mL	60	20	66.7	
	120 µL/mL	60	13	78.3	
	160 µL/mL	60	9	85.0	
	200 µL/mL	60	4	93.3	
	400 µL/mL	60	0	100	

NA: Not available

**Table 2:** Effect of RS1, RS3, and RS4 on root length and newly grown root tips at 0, 24, 48, and 72 h

Control/sample	Concentration	Root length (cm)				Total number of newly grown root tip at			
		0 h (n=30)	24 h (n=27)	48 h (n=24)	72 h (n=21)	0 h	24 h	48 h	72 h
Water	1 mL/mL	2.46±0.20	3.82±0.33	5.02±0.32	6.41±0.43	0	10.33±1.53	14.33±0.58	22.67±0.58
RS1	1 µL/mL	2.31±0.17	3.62±0.28*	4.78±0.35*	5.51±0.30**	0	9.67±1.53 (NS)	13.67±0.58 (NS)	21.67±1.16 (NS)
	10 µL/mL	2.43±0.18	3.51±0.24**	4.36±0.28**	5.12±0.32**	0	9.33±0.58 (NS)	12.67±1.16 (NS)	20.33±0.58**
	100 µL/mL	2.54±0.17	3.14±0.24**	3.67±0.38**	4.02±0.27**	0	8.67±0.58 (NS)	11.33±1.16**	18.67±0.58**
	200 µL/mL	2.44±0.17	2.91±0.29**	2.90±0.30**	2.92±0.25**	0	8.33±0.58 (NS)	8.67±1.16**	8.67±1.16**
RS3	1 µL/mL	2.41±0.16	3.52±0.23**	4.29±0.35**	5.11±0.18**	0	9.33±1.53 (NS)	12.33±0.58 (NS)	20.67±0.58*
	10 µL/mL	2.31±0.16	2.71±0.36**	3.28±0.20**	3.85±0.20**	0	9.33±0.58 (NS)	11.33±0.58**	18.67±0.58**
	100 µL/mL	2.35±0.18	2.41±0.13**	2.49±0.18**	2.46±0.15**	0	8.33±0.58 (NS)	8.33±0.58**	8.33±0.58**
	200 µL/mL	2.59±0.25	2.60±0.26**	2.58±0.26**	2.58±0.26**	0	7.33±1.16**	7.33±1.16**	7.33±1.16**
RS4	1 µL/mL	2.57±0.22	3.46±0.20**	4.53±0.35**	4.93±0.16**	0	8.33±1.16 (NS)	9.67±1.53**	12.33±0.58**
	10 µL/mL	2.46±0.19	3.25±0.28**	3.83±0.28**	4.02±0.26**	0	8.33±0.58 (NS)	8.33±0.58**	8.33±0.58**
	100 µL/mL	2.53±0.25	2.61±0.27**	2.60±0.27**	2.58±0.28**	0	1.67±1.58**	1.67±0.58**	1.67±0.58**
	200 µL/mL	2.48±0.18	2.48±0.19**	2.48±0.20**	2.46±0.20**	0	0.33±0.58**	0.33±0.58**	0.33±0.58**

Values are expressed as mean±SD; \*P<0.05; \*\*P<0.01; NS (P>0.05). NS: Not significant; SD: Standard deviation

**Table 3:** Effect of RS1, RS3, and RS4 on total number of dividing cells, mitotic index at 0, 24, 48, and 72 h, and their IC<sub>50</sub> values after 72 h

Control/ sample	Concentration	Cells manifesting different stages of mitosis				Mitotic index				Percentage of inhibition at 72 h	IC <sub>50</sub> (µL/mL)
		0h	24h	48h	72h	0h	24h	48h	72h		
Water	1 mL/mL	252.33±11.02	247.67±10.26	248.67±17.50	243.67±14.30	50.47±2.20	50.13±2.08	49.73±3.5	48.73±2.86	0	NA
RS1	1 µL/mL	248.33±13.05	250.33±11.06 (NS)	244.67±9.45 (NS)	239.33±9.07 (NS)	49.67±2.61	50.07±2.21 (NS)	48.93±1.89 (NS)	47.87±1.82 (NS)	1.76	380.22
	10 µL/mL	242.33±6.51	252.67±20.79 (NS)	238.67±10.50 (NS)	238.67±10.50 (NS)	48.47±1.30	50.53±4.16 (NS)	47.73±2.10 (NS)	47.33±2.10 (NS)	2.87	
	100 µL/mL	248.67±7.02	250.67±13.58 (NS)	228.67±9.29 (NS)	228.67±9.29 (NS)	49.73±1.41	50.13±2.72 (NS)	45.73±1.86 (NS)	45.73±1.86 (NS)	6.16	
	200 µL/mL	251.67±13.01	247.33±5.03 (NS)	226.33±21.50 (NS)	216.33±10.60*	50.33±2.60	49.47±1.01 (NS)	45.27±4.3 (NS)	43.27±2.12*	11.2	
RS3	1 µL/mL	252.33±6.51	252.67±7.64 (NS)	246.33±14.50 (NS)	244.67±14.57 (NS)	50.47±1.30	50.53±1.53 (NS)	46.27±3.0 (NS)	48.93±2.91 (NS)	0.41	440.64
	10 µL/mL	244.67±11.72	225.33±13.58 (NS)	231.33±5.86 (NS)	227.67±8.33 (NS)	48.93±2.34	45.07±2.72 (NS)	46.27±1.17 (NS)	45.53±1.67 (NS)	6.57	
	100 µL/mL	256.67±10.02	223.67±9.29 (NS)	216.33±4.73*	214.33±5.03**	51.33±2.0	44.73±1.86 (NS)	43.27±0.95*	42.87±1.01**	12.02	
	200 µL/mL	242.33±14.05	219.67±14.98 (NS)	206.33±6.66**	206.33±4.51**	48.47±2.81	43.93±3.0*	41.27±1.33**	41.27±0.90**	15.31	
RS4	1 µL/mL	235.67±6.43	234.33±10.79 (NS)	214.33±6.81*	208.67±3.51**	50.27±2.4	49.53±2.53 (NS)	47.27±2.73 (NS)	46.53±1.62 (NS)	4.51	189.86
	10 µL/mL	254.67±7.37	231.33±11.72 (NS)	164.33±7.64**	142.67±10.41**	47.13±1.29	46.87±2.16 (NS)	42.87±1.36*	41.73±0.70**	14.36	
	100 µL/mL	237.33±11.68	203.33±8.51**	152.67±9.71**	113.33±6.81**	50.93±1.47	46.27±2.34 (NS)	32.87±1.53**	28.53±2.08**	41.45	
	200 µL/mL					47.47±2.34	40.67±1.7**	30.53±1.94**	22.67±1.36**	53.48	

Total number of cells counted was 500; values are expressed as mean±SD; \* P<0.05; \*\* P<0.01; NS (P>0.05). NS: Not significant; SD: Standard deviation

**Table 4:** Percentage of inhibition and IC<sub>50</sub> values of RS1, RS2, RS3, and RS4 in MCF-7, HeLa, and Calu-6 cell lines

Samples	Concentration (µL/mL)	MCF-7 cell lines			HeLa cell lines			Calu-6 cell lines		
		ABS at 590 nm	Percentage of inhibition	IC <sub>50</sub> (µL/mL)	ABS at 590 nm	Percentage of inhibition	IC <sub>50</sub> (µL/mL)	ABS at 590 nm	Percentage of inhibition	IC <sub>50</sub> (µL/mL)
Control	-	0.7368	0	0	0.5568	0	0	0.7632	0	0
RS1	166.67	0.6079	17.50	>500	0.4912	11.78	>500	0.6574	13.86	>500
	200	0.5897	19.97		0.4717	15.28		0.6203	18.72	
	250	0.5468	25.79		0.4543	18.41		0.5558	27.18	
	333.33	0.5121	30.50		0.3957	28.93		0.5343	29.99	
	500	0.4826	34.50		0.3727	33.06		0.5012	34.33	
RS2	166.67	0.5961	19.10	>500	0.4779	14.17	>500	0.6687	12.38	>500
	200	0.5758	21.85		0.4488	19.40		0.6427	15.79	
	250	0.5517	25.12		0.4072	26.87		0.6198	18.79	
	333.33	0.5281	28.33		0.3631	34.79		0.5958	21.93	
	500	0.5017	31.91		0.3245	41.72		0.5524	27.62	
RS3	166.67	0.5996	18.62	>500	0.4687	15.82	>500	0.5779	24.28	>500
	200	0.5682	22.88		0.4427	20.49		0.5488	28.09	
	250	0.5417	26.48		0.4198	24.60		0.5072	33.54	
	333.33	0.5101	30.77		0.3958	28.92		0.4631	39.32	
	500	0.4814	34.66		0.3524	36.71		0.4245	44.38	
RS4	166.67	0.5562	24.51	350.22	0.4892	12.14	>500	0.6452	15.46	410.15
	200	0.5303	28.03		0.4565	18.01		0.6050	20.73	
	250	0.4982	32.38		0.4224	24.14		0.5894	22.77	
	333.33	0.3843	47.84		0.4020	27.80		0.4663	38.90	
	500	0.3125	57.59		0.3786	32.00		0.3611	52.69	

ABS: Absorbance

exhibited antimetabolic activity of RS4 may be due to its interaction with microtubules.

RS4 showed an  $IC_{50}$  value of 350.22 and 410.15  $\mu\text{L}/\text{mL}$  in MCF-7 and Calu-6 cell lines, respectively [Table 4]. RS1, RS2, and RS3 showed an  $IC_{50}$  value more than 500  $\mu\text{L}/\text{mL}$  in MCF-7, HeLa, and Calu-6 cell lines, respectively. The results showed that the final product RS4 may be active against breast and lung cancers. However, the mechanism for such an effect needs further evaluation.

The postmodern “preventive medicine” concept of the Western medicine has absolutely recognized that food plays an important role in the incidence of many diseases. Dietary choice remains the basis for maintaining a healthy lifestyle and well-being, especially relating to cardiovascular disease, diabetes, obesity, hypertension, some cancers, circulatory diseases, and stroke, despite remarkable advances in medicine and pharmaceutical drug development.<sup>[19,20]</sup> According to the National Cancer Registry Programme of the India Council of Medical Research, more than 1300 Indians die every day due to cancer. Between 2012 and 2014, the mortality rate due to cancer increased by approximately 6%. The risk of cancer incidence may also be due to the deviation from traditional functional food toward fast, junk, and westernized foods.

The different ingredients used in *rasam* have been individually attributed to various pharmacological effects in preclinical and clinical studies. It is trouble-free to ascertain that the *rasam*'s effects are due to the antioxidant effect of tamarind fruit pulp;<sup>[21,22]</sup> antioxidant and anticarcinogenic effect of turmeric;<sup>[23]</sup> antioxidant and anticancer activity of chili pepper;<sup>[21]</sup> antioxidant activity of cumin;<sup>[21]</sup> anticancer and antioxidant effects of garlic bulbs;<sup>[21,24-27]</sup> antioxidant and bioavailability enhancing effect of black pepper;<sup>[28,29]</sup> and antioxidant activity of coriander leaves.<sup>[30]</sup> However, if all ingredients and/or active constituents of *rasam* were scientifically formulated together using available technology, it may not yield the desired physiological result; however, somehow in the preparation of *rasam*, the traditional processing naturally ensures higher cytotoxic, antimetabolic, and antiproliferation activity in the final product. The  $LC_{50}$  and  $IC_{50}$  values of *rasam* may not be very significant compared to active pharmaceutical agents that are administered in a fixed dose but consuming *rasam* as daily diet can ensure healing effect. The real challenge lies not in proving whether *rasam* is functional foods having health benefits, but in defining what these benefits are and developing the methods to expose them by scientific means.

## CONCLUSION

*Rasam* is a South Indian traditional functional food that can treat breast and lung cancer on chronic use.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Choudhary R, Tandon RV. Consumption of functional food and our health concerns. *Pak J Physiol* 2009;5:76-83.
- Weiss RS. *Recipes for Immortality: Healing, Religion, and Community in South India*. 1<sup>st</sup> ed. Wellington: Oxford University Press; 2009.
- Calabrese V, Cornelius C, Dinkova-Kostova AT, Calabrese EJ, Mattson MP. Cellular stress responses, the hormesis paradigm, and vitagenes: Novel targets for therapeutic intervention in neurodegenerative disorders. *Antioxid Redox Signal* 2010;13:1763-811.
- Tapsell LC, Hemphill I, Cobiac L, Patch CS, Sullivan DR, Fenech M, et al. Health benefits of herbs and spices: The past, the present, the future. *Med J Aust* 2006;185:S4-24.

- Sainani GS, Desai DB, Gorhe NH, Natu SM, Pise DV, Sainani PG, et al. Dietary garlic, onion and some coagulation parameters in Jain community. *J Assoc Physicians India* 1979;27:707-12.
- Todiwala C. *Mr Todiwala's Spice Box: 120 Recipes with Just 10 Spices*. 1<sup>st</sup> ed. London: Octopus Publishing Group; 2016.
- Srinivasan K. Spices as influencers of body metabolism; an overview of three decades of research. *Food Res Int* 2005;38:77-86.
- Prasad VG, Reddy N, Francis A, Nayak PG, Kishore A, Nandakumar K, et al. Sambar, an Indian dish prevents the development of dimethyl hydrazine-induced colon cancer: A preclinical study. *Pharmacogn Mag* 2016;12 Suppl 4:S441-5.
- Patwardhan B, Khambholja K. Drug discovery and ayurveda: Win-win relationship between contemporary and ancient sciences. In: Kapetanovic I, editor. *Drug Discovery and Development – Present and Future*. Croatia: In Tech; 2011. p. 9-10.
- Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL, et al. Brine shrimp: A convenient general bioassay for active plant constituents. *Planta Med* 1982;45:31-4.
- Rai KM, Basavaraju YB, Sadashivamurthy B. Biological assay and antimetabolic activity of novel analogues of  $\beta$ -apocropodophyllin. *Indian J Pharm Sci* 2007;69:116-8.
- Lau CB, Ho CY, Kim CF, Leung KN, Fung KP, Tse TF, et al. Cytotoxic activities of *Coriaria versicolor* (Yunzhi) extract on human leukemia and lymphoma cells by induction of apoptosis. *Life Sci* 2004;75:797-808.
- Mallone R, Mannering SI, Brooks-Worrell BM, Durinovic-Belló I, Cilio CM, Wong FS, et al. Isolation and preservation of peripheral blood mononuclear cells for analysis of islet antigen-reactive T cell responses: Position statement of the T-cell workshop committee of the immunology of diabetes society. *Clin Exp Immunol* 2011;163:33-49.
- Solis PN, Wright CV, Anderson MM, Gupta MP, Phillipson JD. A microwell cytotoxicity assay using *Artemia salina* (brine shrimp). *Planta Med* 1993;59:250-2.
- Anderson JE, Goetz CM, McLaughlin JL, Suffness M. A blind comparison of simple bench-top bioassays and human tumour cell cytotoxicities as antitumor prescreens. *Phytochem Anal* 1991;2:107-11.
- Mollinedo F, Gajate C. Microtubules, microtubule-interfering agents and apoptosis. *Apoptosis* 2003;8:413-50.
- Babatunde B, Anabuik F. *In vivo* cytogenotoxicity of electronic waste leachate from Iloabuchi electronic market, Diobu, Rivers State, Nigeria on *Allium cepa*. *Challenges* 2015;6:173-87.
- Jordan MA, Wilson L. Microtubules as a target for anticancer drugs. *Nat Rev Cancer* 2004;4:253-65.
- Zarraga IG, Schwarz ER. Impact of dietary patterns and interventions on cardiovascular health. *Circulation* 2006;114:961-73.
- Retelny VS, Neuendorf A, Roth JL. Nutrition protocols for the prevention of cardiovascular disease. *Nutr Clin Pract* 2008;23:468-76.
- Parthasarathy VA, Chempakam B, Zachariah TJ, editors. *Chemistry of Spices*. 1<sup>st</sup> ed. Oxfordshire: CAB International; 2008.
- Caluwe EDe, Halamova K, Damme PV. *Tamarindus indica* L. – A review of traditional uses, phytochemistry and pharmacology. *Afrika Focus* 2010;23:53-83.
- Itokawa H, Shi Q, Akiyama T, Morris-Natschke SL, Lee KH. Recent advances in the investigation of curcuminoids. *Chin Med* 2008;3:11.
- Banerjee SK, Dinda AK, Manchanda SC, Maulik SK. Chronic garlic administration protects rat heart against oxidative stress induced by ischemic reperfusion injury. *BMC Pharmacol* 2002;2:16.
- Hassan HT. Ajoene (natural garlic compound): A new anti-leukaemia agent for AML therapy. *Leuk Res* 2004;28:667-71.
- Tilli CM, Stavast-Kooy AJ, Vuerstaek JD, Thissen MR, Krekels GA, Ramaekers FC, et al. The garlic-derived organosulfur component ajoene decreases basal cell carcinoma tumor size by inducing apoptosis. *Arch Dermatol Res* 2003;295:117-23.
- Terrason J, Xu B, Li M, Allart S, Davignon JL, Zhang LH, et al. Activities of Z-ajoene against tumour and viral spreading *in vitro*. *Fundam Clin Pharmacol* 2007;21:281-9.
- Vijayakumar RS, Surya D, Nalini N. Antioxidant efficacy of black pepper (*Piper nigrum* L.) and piperine in rats with high fat diet induced oxidative stress. *Redox Rep* 2004;9:105-10.
- Khajuria A, Thusu N, Zutshi U. Piperine modulates permeability characteristics of intestine by inducing alterations in membrane dynamics: Influence on brush border membrane fluidity, ultrastructure and enzyme kinetics. *Phytomedicine* 2002;9:224-31.
- Melo EA, Bion FM, Filho JM, Guerra NB. *In vivo* antioxidant effect of aqueous and etheric coriander (*Coriandrum sativum* L.) extracts. *Eur J Lipid Sci Technol* 2003;105:483-7.