A multifaceted peer reviewed journal in the field of Pharmacognosy and Natural Products www.phcog.com | www.phcog.net

# Evaluation of Hypolipidemic and Antioxidant Effects in Phenolrich Fraction of *Crataegus pinnatifida* Fruit in Hyperlipidemia Rats and Identification of Chemical Composition by Ultraperformance Liquid Chromatography Coupled with Quadropole Time-of-flight Mass Spectrometry

#### Feng Shao, Lifei Gu<sup>1</sup>, Huijuan Chen<sup>2</sup>, Ronghua Liu, Huilian Huang, Lanying Chen, Ming Yang

Key Laboratory of Modern Preparation of TCM, Ministry of Education, Jiangxi University of Traditional Chinese Medicine, Nanchang, Jiangxi, <sup>1</sup>Department of Complex Prescription of Traditional Chinese Medicine, China Pharmaceutical University, Nanjing, Jiangsu, <sup>2</sup>Department of Pharmacy, The People's Hospital of Bozhou, Anhui, China

Submitted: 12-09-2016

Revised: 03-11-2016

Published: 13-11-2017

#### ABSTRACT

Background: Hawthorn (Crataegus pinnatifida) fruit has enjoyed a great popularity as a pleasant-tasting food associated with hypolipidemic and antioxidant effects. Objective: Our aim was to screen the effective fraction of hawthorn fruit in the treatment of hyperlipidemia rats. Materials and Methods: In this study, ethanol extract of hawthorn fruit (Fr.1) and four fractionated extracts (Fr.2, Fr.3, Fr.4, and Fr.5) were compared to total phenol content evaluated using Folin-Ciocalteu method, and hypolipidemic and antioxidant effects were assessed in hyperlipidemic rats. Results: Total phenol content of Fr.4 was higher than other fractions by at least 2 fold. Furthermore, this fraction possessed the strongest hypolipidemic and antioxidant effects in hyperlipidemic rats. On this basis, 15 phenolic compounds and four organic acids in Fr.4 were positively or tentatively identified using ultra-performance liquid chromatography coupled with quadropole time-of-flight mass spectrometry. In addition, 5-O-caffeoyl quinic acid butyl ester was first reported in hawthorn fruit. **Conclusion:** Phenol-rich fraction in hawthorn fruit exhibited satisfactory hypolipidemic and antioxidant effects, and this could be exploited for further promotion of functional foods. Key words: Antioxidant effect, Crataegus pinnatifida fruit, hypolipidemic effect, phenolic compounds, ultra performance liquid chromatography coupled with quadropole time-of-flight mass spectrometry

#### **SUMMARY**

 Phenol-rich fraction in hawthorn fruit possesses most potent hypolipidemic and antioxidant effects in hyperlidemia rats.



Abbreviationsused:UPLC-Q-TOF-MS/MS:Ultraperformanceliquidchromatographycoupledwithquadropoletime-of-flightmassspectrometry;TC:Totalcholesterol;TG:Triglyceride;LDLC:Low-densitylipoprotein-cholesterol;HDLC:High-densitylipoprotein-cholesterol;GSH-Px:Glutathioneperoxidase;SOD:Superoxidedismutase;MDA:Malondialdehyde;CAT:Catalase;NO:Nitricoxide;NOS:Nitricoxide•OH:Hydroxyl radicals.•OH:Hydroxyl radicals.•OH:Superoxideanions,

#### Correspondence:

Prof. Ming Yang, Key Laboratory of Modern Preparation of TCM, Ministry of Education, Jiangxi University of Traditional Chinese Medicine, Jiangxi, Nanchang 330004, China. E-mail: youngming501@163.com **DOI:** 10.4103/pm.pm\_402\_16



# INTRODUCTION

Hawthorn (*Crataegus pinnatifida*), from *Rosaceae* family, is a deciduous tree mainly distributed and cultivated in temperate areas, including China, Korea, and Russia.<sup>[1]</sup> The eatable and medicinal history of hawthorn fruit in China could be dated back to about 300 AD.<sup>[2]</sup> It has been used as a traditional medicine for the treatment of dyspepsia, cardiovascular disease, and hyperlipidemia.<sup>[3]</sup> Recent reports have disclosed hypolipidemic and antioxidant effects of the traditional medicine.<sup>[4,5]</sup> Besides, our previous investigation had indicated that these effects of hawthorn fruit were probably caused by the existence of phenols.<sup>[6]</sup>

Dietary phenols appear to possess antioxidant property, which scavenge reactive oxygen and nitrogen species, thereby potentially contributing against the pathogenesis of cardiovascular disease.<sup>[7]</sup> Phenols in hawthorn fruit are responsible for free radical quenching activity and considered to be the best antilipoperoxidants.<sup>[8]</sup> Hawthorn fruit is

abundant in phenolic compounds, including isoquercetin, hyperoside, protocatechuic acid, and chlorogenic acid,<sup>[9]</sup> which usually have hypolipidemic and antioxidant effects.<sup>[10]</sup> For the above reasons, we hypothesized that phenol-rich fraction in hawthorn fruit probably has hypolipidemic and antioxidant effects in hyperlipidemic rats.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

**Cite this article as:** Shao F, Gu L, Chen H, Liu R, Huang H, Chen L, *et al.* Evaluation of hypolipidemic and antioxidant effects in phenol-rich fraction of *Crataegus pinnatifida* fruit in hyperlipidemia rats and identification of chemical composition by ultra-performance liquid chromatography coupled with quadropole time-of-flight mass spectrometry. Phcog Mag 2017;13:725-31.

Although previous investigation has shown that high-performance liquid chromatography (HPLC)-electrospray ionization (ESI) mass spectrometry (MS) has been used in the characterization of 42 phenolic compounds in 80% ethanol extract of hawthorn fruit,<sup>[11,12]</sup> it is still unknown information concerning bioactive fraction and its compounds in hypolipidemic and antioxidant effects. At present, ultra performance liquid chromatography coupled with quadropole time-of-flight MS (UPLC-Q-TOF-MS/MS) demonstrated to be a more powerful tool for analyzing natural products, mainly because of its high resolution as well as accuracy in weight measurements.<sup>[13]</sup> It helps us to get more information concerning phenolic compounds and others in bioactive fraction of hawthorn fruit.

Hence, this study was designed to determine total phenol content of ethanol extract and four fractionated extracts by the Folin– Ciocalteu method to evaluate hypolipidemic and antioxidant effects in hyperlipidemic rats and to identify phenolic compounds and others in bioactivity fraction by UPLC-Q-TOF-MS/MS.

# **MATERIALS AND METHODS**

#### Materials and chemicals

Hawthorn (*C. pinnatifida* Bge. var. *major* N. E. Br) fruit was collected from Pingyi County (Shandong Province, China) and authenticated by Professor Kezhong Deng in School of Pharmacy, Jiangxi University of Traditional Chinese Medicine. A voucher specimen (No. 20121123) has been deposited in the Key Laboratory of Modern Preparation of TCM, Ministry of Education, Jiangxi University of Traditional Chinese Medicine.

Total cholesterol (TC) assay kit (Lot No. ZG3001), triglyceride (TG) assay kit (Lot No. ZG3001), low-density lipoprotein-cholesterol (LDL-C) assay kit (Lot No. ZG9003), and high-density lipoprotein-cholesterol (HDL-C) assay kit (Lot No. ZG3001) were offered by Sysmex Co, Japan. Glutathione peroxidase (GSH-Px) assay kit (Lot No. 20130831), superoxide dismutase (SOD) kit (Lot No. 20130831), malondialdehyde (MDA) assay kit (Lot No. 20130831), catalase (CAT) assay kit (Lot No. 20130903), nitric oxide (NO) assay kit (Lot No. 20130827), and NO synthase (NOS) assay kit (Lot No. 20130903) were obtained from Nanjing Jiancheng Bio-Engineering Company, China. No. 3 bile salt (Lot No. 20130531-00) was purchased from Hangzhou Hongbo Biological Engineering Co., Ltd, China. Propylthiouracil (Lot No. 20130523) and cholesterol were obtained from Wuhan Sheng Tianyu Biological Technology Co., Ltd, China. Lard was purchased from Henan Zhumadian Dingsheng Food Co., Ltd. Zhumadian, China.

Hyperoside (99%), quercetin (99%), vitexin (98%), and isovitexin (98%) were purchased from Must Bio-technology Co., Ltd. (Chengdu, China). Acetonitrile and formic acid for UPLC were obtained from Dikma Technologies Inc. (Lake Forest, USA) and HPLC-grade formic acid was purchased from Aladdin, China. Folin–Ciocalteu phenol reagent was

# Extraction and fractionation of hawthorn fruit

obtained from Sigma-Aldrich, USA.

Hawthorn fruit (45 kg) was extracted with 70% ethanol and the ratio of material to solvent was 1:3, under reflux successively (each 2 h, 4 times) and filtered. After concentration under vacuum condition, the obtained extract (Fr.1) was suspended in water and then partitioned with petroleum ether to obtain Fr.2. The pH of remaining aqueous fraction was adjusted to  $2.0 \pm 0.5$  with HCl, and then partitioned with water-saturated butanol. Ethanol was added to the aqueous fraction to precipitate polysaccharide labeled as Fr.3, and the water-saturated *n*-butanol fraction was extracted with 1% NaHCO<sub>3</sub>. The *n*-butanol fraction was adjusted to  $2.0 \pm 0.5$  with HCl and

extracted with water-saturated butanol again. The supernatant was evaporated to obtain Fr.5.

# Determination of total phenol content

Folin–Ciocalteu method was used to determine the content of total phenols using ultraviolet (UV) and visible spectrophotometer (Shimadzu, Japan) with gallic acid (6.25–100  $\mu$ g/ml) served as reference.<sup>[13]</sup> All samples were analyzed in three replications.

#### Animals

Male Sprague-Dawley rats  $(230 \pm 20 \text{ g}, \text{ age } 7-8 \text{ weeks})$  were supplied by Hunan Lake King of Laboratory Animal Co., Ltd. (Hunan, China). Rats were kept in room temperature  $(22-25^{\circ}\text{C}, 55 \pm 10\% \text{ humidity}, \text{ and} 12/12 \text{ h light/darkness cycle})$  with commercial rat normal standard chow (Hunan SJA Laboratory Animal Co., Ltd., Hunan, China) and water *ad libitum*. The surgical procedures and experimental protocol were approved by the Animal Ethics Committee of Jiangxi University of Traditional Chinese Medicine.

After allowing 7 days for adaptation, all rats were randomly assigned to 7 groups (n = 8). Group 1 rats (control) were orally treated with distilled water (10 ml/kg body weight) through gavage. Groups 2, 3, 4, 5, 6, and 7 rats were intragastrically administered with high-fat emulsion (cholesterol 10 g, propylthiouracil 1 g, lard oil 25 g, tween-80 25 ml, propylene glycol 20 ml, and No. 3 bile salt 2 g) (10 ml/kg body weight) once a day.<sup>[6]</sup> After 6 h, Group 1 (control) and Group 2 rats (model) were given distilled water (10 ml/kg body weight). Groups 3, 4, 5, 6, and 7 rats received the corresponding five fractions (Fr.1–Fr.5) at low-, medium-, and high-dose (equivalent to about 75, 150, and 300 mg/kg body weight), respectively.

After 4 weeks of administration by gastric gavage, the rats were fasted for 12 h and euthanized by decapitation. Blood was collected, left at room temperature for 15 min, and then centrifuged at 3000 rpm (4°C, 10 min). The serum obtained was stored at  $-80^{\circ}$ C until biochemical analysis. Livers were dissected, washed with saline, and homogenized (weighed 0.5 g, added 4.5 ml normal saline). The samples were centrifuged at 3500 rpm (4°C, 10 min). The supernatants were obtained and stored at  $-80^{\circ}$ C immediately until enzyme activity analysis.

#### Serum lipids and antioxidant enzyme activities

The serum lipid levels (TC, TG, LDL-C, and HDL-C) were measured by an automatic biochemical analyzer (Serial No. CHEMIX-180, Sysmex, Japan). The antioxidant enzyme activities (SOD, CAT, and GSH-Px) and the levels of MDA, NO, and NOS in serum and liver were, respectively, determined using commercial analysis kits by Microplate Reade (Serial No. SpectraMax 190, Molecular Devices, USA).

# Ultra-performance liquid chromatography coupled with quadropole time-of-flight mass spectrometry analysis

A Shimadzu UPLC system combined with an AB SCIEX triple TOF<sup>TM</sup> 5600<sup>+</sup> mass spectrometer system equipped with an DuoSprayTM source was used to acquire mass spectra of bioactivity fraction. The separations were performed on a 2.1 mm × 100 mm ACUIITY UPLC<sup>°</sup> HSS T3 column (1.8  $\mu$ m, USA) at 30°C, and the UV absorbance was monitored at 270 nm. The solvent system composed of acetonitrile (A) and 0.1% formic acid in water (B) using an optimized gradient program as follows: 0–5.0 min, 5%–5% of A; 5.0–65.0 min, 5%–34% of A.

Optimum parameters of MS in negative ESI modes were set as follows: ion spray voltage, -4500 V; collision energy, -40 V; and declustering potential, -100 V. The nebulizing gas (Gas 1) was 50 psi, heater gas (Gas

2) was 60 psi, and the curtain gas was 30 psi. Mass scan was over the m/z 100–1600, and turbo spray temperature was 600°C. Data were analyzed by Peak View Software<sup>TM</sup> 1.2 (AB SCIEX, Canada).

#### Statistical analysis

Values were presented as mean  $\pm$  standard deviation. A paired *t*-test was employed to evaluate statistical significance between the two groups with SPSS software (version 22 for Windows, Chicago, IL, USA). Differences are considered to be statistically significant when P < 0.05.

# **RESULTS AND DISCUSSION**

#### Determination of total phenol content

Phenolic compounds are considered the most critical bioactive compounds in *C. pinnatifida* BUNGE.<sup>[9]</sup> These chemical constituents possess potential health effects, such as lowering blood lipids and attenuating oxidative stress.<sup>[6,13]</sup> In this study, according to acidity and polarity principle, their ethanol extract (Fr.1) was fractioned into four fractionated extracts (Fr.2, Fr.3, Fr.4, and Fr.5) [Figure 1]. Compared with four others, Fr.4 was rich in phenols by possessing more than 2 times of total phenol content, as shown in Table 1. Hence, Fr.4 was probably attributed to the hypolipidemic and antioxidant activities with its abundant phenols.

#### Evaluation of hypolipidemic effect

TC, TG, LDL-C, and HDL-C are regarded as the key evaluation indexes of hyperlipidemic model. Serum lipid profiles can be attenuated by ethanol extract of hawthorn fruit.<sup>[14]</sup> In this study, compared to normal rats, TC, TG, and LDL-C levels in serum were markedly raised (P < 0.01) by administrating rats with a high-fat emulsion for 4 weeks, as shown in Figure 2. However, there was no significant difference between control group and model group for HDL-C that might be associated with the compensatory mechanism of rats themselves.<sup>[6,15]</sup>

On the basis of established model, we estimated hypolipidemic capacity of ethanol extract (Fr.1) and four fractionated extracts (Fr.2, Fr.3, Fr.4, and Fr.5) in hyperlipidemic rats. As a result, compared with the model, Fr.4 is the only one that at all doses lowered the TC, TG, and LDL-C contents significantly in hyperlipidemic rats (P < 0.05). Simultaneously, Fr.4 at medium and high doses and Fr.1 at medium dose elevated the level of HDL-C, but no significant difference was observed (P > 0.05), consistent with several previously published reports.<sup>[16-18]</sup> It is noteworthy that

compared with other fractions, Fr.4 at low dose decreased the levels of TC, TG, and LDL-C more obviously in hyperlipidemic rats (P < 0.05), as shown in Figure 2. Therefore, Fr.4 showed more significant ameliorative action than others in the serum lipid levels of hyperlipidemic rats.

#### Determination of antioxidant effect

Oxidative stress is a causative factor, which links hyperlipidemia with the pathogenesis of atherosclerosis,<sup>[19]</sup> and is induced by reactive oxygen species, for example, superoxide anions (•OOH) and hydroxyl radicals (•OH).<sup>[20]</sup> The activities of GSH-Px, SOD, and CAT directly reflect the ability of scavenging

Table 1: Total phenols of the fractions from Crataegus pinnatifida fruit (mg/g)

Fractions	Total phenol content
Fr. 1	83.6±0.67
Fr. 2	ND
Fr. 3	100.2±1.39
Fr. 4	678.2±2.20
Fr. 5	327.4±1.56

Values are represented as mean±SD of three different experiments. ND: Not detected; SD: Standard deviation



Figure 1: Preparation procedure of the fractions of hawthorn fruit



**Figure 2:** Effects of *Crataegus pinnatifida* fruit fractions on total cholesterol, triglyceride, high-density lipoprotein cholesterol, and low-density lipoprotein cholesterol in rats fed on normal diet or high-fat emulsion diet. The low-, medium-, and high-doses were equivalent to about 75, 150, and 300 mg/kg body weight, respectively. Values are shown as mean  $\pm$  standard deviation (n = 8). \*P < 0.05 and \*\*P < 0.01 compared with the control group. \*P < 0.05 and \*\*P < 0.01 compared with the model group

oxygen-free radicals.<sup>[21]</sup> In addition, MDA, the end product of lipid peroxidation, is caused by free radical chain reaction. In this study, after 4 weeks of treatment, compared to control group, decreased activities of SOD, CAT, and GSH-Px and increased content of MDA in serum and liver were observed in hyperlipidemic rats, respectively, as shown in Figures 3 and 4.

Phenolic compounds reduce the risk of hyperlipidemia and oxidative injury through increasing antioxidant enzyme activity and reducing

free radical formation.<sup>[22]</sup> In the similar report, seven phenolic compounds, namely, hyperoside, isoquercitrin, epicatechin, quercetin, rutin, chlorogenic acid, and protocatechuic acid in hawthorn fruit are extremely effective in protecting LDL from oxidation.<sup>[23]</sup> As there is phenol-rich fraction in hawthorn fruit, Fr.4 at high dose is the most remarkable one to increase the lowered SOD, CAT, and GSH-Px activities (P < 0.01) and to decrease the elevated MDA level (P < 0.01) in serum



**Figure 3:** Effects of *Crataegus pinnatifida* fruit on malondialdehyde, nitric oxide content, and the activities of catalase, superoxide dismutase, glutathione peroxidase, and nitric oxide synthase of serum in rats fed on high-fat emulsion or normal diet. The low-, medium-, and high-doses were equivalent to about 75, 150, and 300 mg/kg body weight, respectively. Values are shown as mean  $\pm$  standard deviation (n = 8). \*P < 0.05 and \*\*P < 0.01 compared with the control group. \*P < 0.05 and \*\*P < 0.01 compared with the model group



**Figure 4:** Effects of *Crataegus pinnatifida* fruit on malondialdehyde content and the activities of superoxide dismutase, catalase, and glutathione peroxidase of liver in rats fed on normal diet or high-fat emulsion diet. The low-, medium-, and high-doses were equivalent to about 75, 150, and 300 mg/kg body weight, respectively. Values are shown as mean  $\pm$  standard deviation (n = 8). \*P < 0.05 and \*\*P < 0.01 compared with the control group. \*P < 0.05 and \*\*P < 0.01 compared with the model group

and liver of hyperlipidemic rats. Furthermore, antioxidant capacity of Fr.4 at medium dose was approximately equivalent or stronger than that of other groups at high dose in serum of hyperlipidemic rats, as shown in Figure 3. A similar situation also existed for the SOD, CAT, and GSH-Px analyses in liver of hyperlipidemic rats, as shown in Figure 4.

NO, which is produced by endothelial NOS, is the principal factor that inhibits vessel platelet aggregation and dilate vessels so as to prevent vascular atherosis and thrombus formation.<sup>[24-26]</sup> Isolated artery experiments have revealed that phenolic compounds could cause NO-mediated endothelium-dependent relaxations and increase the endothelial formation of NO.<sup>[27]</sup> In this study, the NOS and NO levels of Fr.4 were higher than those of other administration groups at corresponding dose in serum of hyperlipidemic rats, as shown in Figure 3. These results indicated that phenol-rich fraction in hawthorn fruit exhibited more significant antioxidant capacity than others in hyperlipidemic rats.

#### Identification of compounds in Fr.4

Based on the above results, compound identification of Fr.4 was carried out by UPLC-Q-TOF-MS/MS. The negative ionization mode was more sensitive under the conditions, and was therefore selected for further use. As a result, 15 phenolic compounds (including six phenolic acids and nine flavonoids) and four organic acids have been identified or tentatively identified according to their retention times and fragment ions, as shown in Figure 5 and Table 2.

#### **Phenolic acids**

Six phenolic acids, such as protocatechuic acid (5), Chlorogenic acid isomers (6, 7), vanillic acid (8), and ferulic acid (18), 5-O-caffeoyl quinic acid butyl ester (19) were tentatively identified. Phenolic acids have strong *in vitro* and *in vivo* antioxidant activities associated with their ability to scavenge free radicals, break radical chain reactions, and chelate metals.<sup>[28]</sup>

Chlorogenic acid has attracted continuous attention among phenolic acids with its alleged biological effects.<sup>[29]</sup> Classically, chlorogenic acids are a family of esters formed between certain cinnamic acids and quinic acid. Peaks 6 and 7 were tentatively characterized as chlorogenic acid isomers for which they displayed the similar parent molecule ion at *m*/*z* 353.0878 and 353.0889. They presented the similar fragmentation pattern at *m*/*z* 191.0565 and 191.0560, corresponding to [quinic acid–H]<sup>-</sup> ion. The deprotonated caffeic acid fragment at *m*/*z* 179.0348 was also found at compound 7, revealing that the quinic acid was substituted at 3-position with the hydroxyl group, while compound 6 was not 3-OH replacement for the absence of *m*/*z* 179 fragmentation.<sup>[30]</sup>

Peak 19 showed a pseudomolecular ion at m/z 409.1528. Its MS/MS spectra gave m/z 191.0561 [quinic acid-H]<sup>-</sup>, 179.0560 [caffeic acid-H]<sup>-</sup>,



Figure 5: Total ion chromatogram of Fr.4 analyzed by ultra-performance liquid chromatography coupled with quadropole time-of-flight mass spectrometry

n	Compound	Formula	lon	t <sub>e</sub>	Calculated	Experimental	Product ion <i>m/z</i>	Erro
			mode		m/z	m/z		ppm
1	Quinic acid	C <sub>7</sub> H <sub>12</sub> O <sub>6</sub>	[M-H] <sup>-</sup>	1.62	191.0561	191.0559	191.0564, 127.0410, 111.0095, 93.0358	-1
2	Malic acid	C <sub>4</sub> H <sub>6</sub> O <sub>5</sub>	[M-H]⁻	1.80	133.0143	133.0152	133.0133, 115.0035	7.2
3	Citric acid	C <sub>6</sub> H <sub>8</sub> O <sub>7</sub>	[M-H] <sup>-</sup>	2.28	191.0197	191.0204	111.0096	3.7
4	Succinic acid	C <sub>4</sub> H <sub>6</sub> O <sub>4</sub>	[M-H] <sup>-</sup>	2.67	117.0193	117.0203	116.9290, 99.9265, 83.9314	8.6
5	Protocatechuic acid	C <sub>7</sub> H <sub>6</sub> O <sub>4</sub>	[M-H] <sup>-</sup>	7.56	153.0193	153.0198	109.0303	3.1
6	Chlorogenic acid isomers	C <sub>16</sub> H <sub>18</sub> O <sub>9</sub>	[M-H] <sup>-</sup>	16.58	353.0878	353.0894	191.0565	4.5
7	Chlorogenic acid isomers	C <sub>16</sub> H <sub>18</sub> O <sub>9</sub>	[M-H] <sup>-</sup>	18.19	353.0878	353.0889	353.0892, 191.0560, 179.0348, 173.0452, 135.0453	3.1
8	Vanillic acid	C <sub>8</sub> H <sub>8</sub> O <sub>4</sub>	[M-H]⁻	23.55	167.0350	167.0357	167.0361, 152.0116, 109.0260, 108.0229, 91.0205	4.3
9	Violanthin	C <sub>27</sub> H <sub>30</sub> O <sub>14</sub>	[M-H] <sup>-</sup>	29.45	577.1563	577.1607	577.1621, 487.1267, 457.1192, 367.0836, 337.0729	7.6
10	Rutin	C <sub>27</sub> H <sub>30</sub> O <sub>16</sub>	[M-H] <sup>-</sup>	30.15	609.1461	609.1516	609.1522, 301.0374, 300.0296, 271.0264	9
11	Schaftoside	C <sub>27</sub> H <sub>30</sub> O <sub>15</sub>	[M-H] <sup>-</sup>	31.16	593.1512	593.1570	593.1580, 473.1101, 413.0911, 311.0593, 293.0460	9.9
12	Vitexin	$C_{21}H_{20}O_{10}$	[M-H] <sup>-</sup>	31.93	431.0984	431.1006	431.1025, 341.0696, 311.0578, 283.0626	5.2
13	Vitexin rhamnoside	C <sub>27</sub> H <sub>30</sub> O <sub>14</sub>	[M-H] <sup>-</sup>	32.09	577.1563	577.1615	577.1635, 457.1189, 413.0915, 311.0579, 293.0475	9
14	Isovitexin	$C_{21}H_{20}O_{10}$	[M-H] <sup>-</sup>	32.73	431.0984	431.1008	431.1026, 341.0673, 311.0585, 283.0627	5.7
15	Hyperoside	$C_{21}H_{20}O_{12}$	[M-H] <sup>-</sup>	32.95	463.0955	463.0923	463.0911, 301.0367, 300.0285, 271.0260, 255.0313	-6.9
16	Isoquercitrin	$C_{21}H_{20}O_{12}$	[M-H] <sup>-</sup>	33.81	463.0955	463.0921	463.0970, 301.0361, 300.0280, 271.0254, 255.0307	-7.3
17	Astragalin	C <sub>21</sub> H <sub>20</sub> O <sub>11</sub>	[M-H] <sup>-</sup>	36.31	447.0933	447.0957	447.0964, 285.0412, 284.0399, 255.0306, 227.0353	5.3
18	Ferulic acid	$C_{10}H_{10}O_{4}$	[M-H] <sup>-</sup>	37.15	193.0506	193.0513	193.500, 178.0267, 161.0244, 133.0298	3.2
19	5-O-caffeoyl quinic acid	C <sub>20</sub> H <sub>26</sub> O <sub>9</sub>	[M-H] <sup>-</sup>	50.50	409.1528	409.1540	409.1528, 191.0561, 179.0560, 161.0244, 135.0456	0.8
	butyl ester	20 20 9						

 Table 2: Mass data of compounds detected in Fr. 4 of Crataegus pinnatifida fruit



161.0244 [caffeic acid–H<sub>2</sub>O]<sup>-</sup>, and 135.0456 [caffeic acid–CO<sub>2</sub>–H]<sup>-</sup>. It has previously reported that 5-O-caffeoyl quinic acid gave the ion at m/z 191 in its MS/MS fragmentation, while 3-O- and 4-O-caffeoyl quinic acids showed different behaviors at m/z 163 and 173, respectively.<sup>[30]</sup> Accordingly, peak 19 was deduced as 5-O-caffeoyl quinic acid butyl ester, which was first reported in hawthorn fruit. The detailed fragmentation pathway of peak 19 is shown in Figure 6.

Peak 5 was characterized as protocatechuic acid with the  $[M-H]^-$  at m/z 153.0198. Furthermore, MS/MS data showed a fragment at m/z 109.0303 (loss of CO<sub>2</sub>).<sup>[31]</sup>

Peaks 8 and 18 were assigned as vanillic acid and ferulic acid, respectively. The loss of a CH<sub>3</sub> moiety from deprotonated ion of peak 8 ([M–H]<sup>-</sup> at m/z 167.0350) and peak 18 ([M–H]<sup>-</sup> at m/z 193.0513) resulted in the MS<sup>2</sup> ion at m/z 152.0116 and 178.0267, respectively. Peak 8 also showed fragment ions at 108.0229 ([M–H–CH<sub>3</sub>–CO<sub>2</sub>]<sup>-</sup>), 91.0205 ([M–H–CH<sub>3</sub>–CO<sub>2</sub>–OH]<sup>-</sup>). Peak 18 showed fragment ions at 161.0244 ([M–H–CH<sub>3</sub>–OH]<sup>-</sup>), 133.0298 ([M–H–CH<sub>3</sub>–OH–CO]<sup>-</sup>) as well.

#### Flavonoids

As a large group of plant phenol secondary metabolites, flavonoids act through scavenging free radicals, promoting anti-oxidase or inhibiting oxidative enzymes while regulating the blood flow and keeping the heart healthy. Nine compounds were unambiguously identified as flavonoid glycosides.

Peaks 15, 16, and 10 all originated from quercetin, they all displayed a characteristic fragment ion at m/z 300, representing the quercetin ion moiety. In this case, the ion at m/z 300, proposed as diagnostic ion for quercetin glycoside, is higher than ion at m/z 301, and could be due to formation of the quinone anion, obtained after hemolytic cleavage of the *O*-glycosidic bond.<sup>[32]</sup> Peaks 15 and 16 that shared  $[M-H]^-$  at m/z 463 were deduced as hyperoside and isoquercitrin, respectively, based on spectra and comparison with standards.<sup>[12]</sup> Peak 10 had a  $[M-H]^-$  at m/z 609.1516 and characterized as rutin, with a chief MS/MS pattern at m/z 300.0296 (loss of rutinose).

Peaks 12 and 14 gave the deprotonated ions at m/z 431, and MS<sup>2</sup> characteristic fragments at m/z 341 [M–H–90]<sup>-</sup>([M–H–C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>]<sup>-</sup>), at m/z 311 [M–H–120]<sup>-</sup> ([M–H–C<sub>4</sub>H<sub>8</sub>O<sub>4</sub>]<sup>-</sup>), which were consistent with the characteristic ions of a *C*-glycosidic flavonoid.<sup>[33]</sup> Accordingly, peaks 12 and 14 were deduced to be vitexin and isovitexin, respectively, in accordance with the MS and MS<sup>2</sup> data of reference compounds.

Peaks 13 and 9 showed negative molecular ions at m/z 577 and were characterized as vitexin rhamnoside and violanthin, respectively, for their fragmentation patterns differed. In the MS<sup>2</sup> spectrum of peak

13, an ion at m/z 413.0915 was noted to lose its terminal rhamnose unit, followed by loss of a  $C_4H_8O_4$  group to give a fragment at m/z 293.0475. Peak 9 displayed characteristic ions at m/z 487.1267  $[M-H-90]^-$ , 457.1192  $[M-H-120]^-$ , 367.0836  $[M-H-120-90]^-$ , and 337.0729  $[M-H-120-120]^-$  of a *C*-glycosidic flavonoid with two glycosyls, which were in accordance with the fragmentation pathway of violanthin.

Peak 11, with deprotonated ion at m/z 593.1580, was tentatively identified as schaftoside, which is a *C*-glycosidic flavonoid with MS/MS fragments at m/z 473.1101 [M–H–120]<sup>-</sup>, 413.0911 [M–H–120–120]<sup>-</sup>, 311.0593 [M–H–120–1203]<sup>-</sup>, and 293.0460 [M–H–120–120]<sup>-</sup>.

Peak 17 gave a higher signal at m/z 447 ([M–H]<sup>-</sup>) which was characterized as astragalin. The fragment pattern gave a main fragment ion at m/z285 (kaempferol aglycone moiety) for the neutral loss of the glucose unit. A higher signal at m/z 284 was also observed, which accorded with previously reported rule that [kaempferol–H]<sup>-</sup> ion is sometimes higher than kaempferol ion in 3–OH position-substituted glycosidic flavonols.<sup>[31,34]</sup>

#### **Organic acids**

Peaks 1, 2, 3, and 4 were assigned as organic acids. Peak 1 showed a typical fragmentation behavior at m/z 191.0564, m/z 127.0410 ([M–H–CO<sub>2</sub>–H<sub>2</sub>O]<sup>-</sup>), m/z 111.0095 ([M–H–CO<sub>2</sub>–2H<sub>2</sub>O]<sup>-</sup>), and m/z93.0358 ([M–H–CO<sub>2</sub>–3H<sub>2</sub>O]<sup>-</sup>), corresponding to quinic acid. Peak 2, characterized as malic acid, showed a negative molecular ion at m/z133.0133, with MS/MS pattern at m/z 115.0035 for the loss of water. Peak 3 had a deprotonated ion at m/z 191.0204 and was deduced as citric acid because the characteristic fragment of citric acid appeared at m/z111.0096 ([M–H–CO<sub>2</sub>–2H<sub>2</sub>O]<sup>-</sup>). The [M–H]<sup>-</sup> fragment of peak 3 was different from peak 1 for the number variance after the decimal point. Peak 4, with a [M–H]<sup>-</sup> at m/z 117.0203, was identified as succinic acid and its fragments at m/z 99.9265, corresponding to elimination of water, and m/z 83.9314 corresponding to lost of another water

# CONCLUSION

In our study, compared with other fractions, phenolic-rich fraction in hawthorn fruit exhibited more significant ameliorative action in lipid profile levels and higher antioxidant contribution in hyperlipidemic rats. Moreover, 15 phenolic compounds and 4 organic acids in this fraction were identified based on MS data and MS/MS fragmentation pattern by UPLC-Q-TOF-MS/MS. Given these findings, we suggested that polyphenolic compounds of hawthorn fruit probably play the key role

in hypolipidemic and antioxidant effects, and phenolic-rich fraction in hawthorn fruit would be used in the further development of functional food.

# Acknowledgement

Our special thanks are due to Ms. Huiming Hu for proofreading on the manuscript.

#### Financial support and sponsorship

This project was supported by grants from the National Natural Science Foundation of China (No. 81260638), the Nature Science Foundation of Jiangxi Province, China (No. 20161BAB205220 and No. 20132BAB205083), the Health and Family Planning Commission Fund of Jiangxi province, China (No. 2013A152 and No. 2014A041), the Innovation Project for Postgraduate Research Fund of Jiangxi province, China (No. YC2016-B079), and the Innovation Project for Doctoral Research Fund of Jiangxi University of Traditional Chinese Medicine, China (No. JZYC16B02).

# **Conflicts of interest**

There are no conflicts of interest.

#### REFERENCES

- Zhao H, Feng B. Cultivation history. China Fruit-plant Monograph-Hawthorn Flora. Ch. II. Beijing, China: Zhongguo Lin Ye Press; 1996.
- Qian C, Chen H, Qin R, Lin R, Wu Z, Cui H, et al. Flora of China. Beijing, China: Science Press; 1974.
- The Pharmacopoeia Commission of PRC. Pharmacopoeia of the People's Republic of China. Beijing, China: Chemical Industry Press; 2015.
- Ling J, Wei B, Lv G, Ji H, Li S. Anti-hyperlipidaemic and antioxidant effects of turmeric oil in hyperlipidaemic rats. Food Chem 2012;130:229-35.
- Belguith-Hadriche O, Bouaziz M, Jamoussi K, Simmonds MS, El Feki A, Makni-Ayedi F. Comparative study on hypocholesterolemic and antioxidant activities of various extracts of fenugreek seeds. Food Chem 2013;138:1448-53.
- Shao F, Gu L, Chen H, Liu R, Huang H, Ren G. Comparation of hypolipidemic and antioxidant effects of aqueous and ethanol extracts of *Crataegus pinnatifida* fruit in high-fat emulsioninduced hyperlipidemia rats. Pharmacogn Mag 2016;12:64-9.
- Chan C, Gan R, Corke H. The phenolic composition and antioxidant capacity of soluble and bound extracts in selected dietary spices and medicinal herbs. Int J Food Sci Technol 2016;51:565-73.
- Chu CY, Lee MJ, Liao CL, Lin WL, Yin YF, Tseng TH. Inhibitory effect of hot-water extract from dried fruit of *Crataegus pinnatifida* on low-density lipoprotein (LDL) oxidation in cell and cellfree systems. J Agric Food Chem 2003;51:7583-8.
- Jurikova T, Sochor J, Rop O, Mlcek J, Balla S, Szekeres L, *et al.* Polyphenolic profile and biological activity of Chinese hawthorn (*Crataegus pinnatifida* BUNGE) fruits. Molecules 2012;17:14490-509.
- Gao H, Long Y, Jiang X, Liu Z, Wang D, Zhao Y, et al. Beneficial effects of Yerba Mate tea (*llex paraguariensis*) on hyperlipidemia in high-fat-fed hamsters. Exp Gerontol 2013;48:572-8.
- Zhang W, Xu M, Yu C, Zhang G, Tang X. Simultaneous determination of vitexin-4"-Oglucoside, vitexin-2"-O-rhamnoside, rutin and vitexin from hawthorn leaves flavonoids in rat plasma by UPLC-ESI-MS/MS. J Chromatogr B 2010;878:1837-44.
- Liu P, Yang B, Kallio H. Characterization of phenolic compounds in Chinese hawthorn (*Crataegus pinnatifida* Bge. var. *major*) fruit by high performance liquid chromatographyelectrospray ionization mass spectrometry. Food Chem 2010;121:1188-97.

- Wen L, Guo X, Liu RH, You L, Abbasi AM, Fu X. Phenolic contents and cellular antioxidant activity of Chinese hawthorn "Crataegus pinnatifida". Food Chem 2015;186:54-62.
- Kwok C, Li C, Cheng H, Ng Y, Chan T, Kwan Y, et al. Cholesterol lowering and vascular protective effects of ethanolic extract of dried fruit of *Crataegus pinnatifida*, hawthorn (Shan Zha), in diet-induced hypercholesterolaemic rat model. J Funct Foods 2013;5:1326-35.
- de Grooth GJ, Klerkx AH, Stroes ES, Stalenhoef AF, Kastelein JJ, Kuivenhoven JA. A review of CETP and its relation to atherosclerosis. J Lipid Res 2004;45:1967-74.
- Zhang Z, Ho WK, Huang Y, James AE, Lam LW, Chen ZY. Hawthorn fruit is hypolipidemic in rabbits fed a high cholesterol diet. J Nutr 2002;132:5-10.
- Chen JD, Wu YZ, Tao ZL, Chen ZM, Liu XP. Hawthorn (Shan Zha) drink and its lowering effect on blood lipid levels in humans and rats. World Rev Nutr Diet 1995;77:147-54.
- Ho WK, Chang HM, Lee CM. Method and Compositions for Lowering Blood Lipids. U.S. Patent No. 5665359; 1997.
- Slim RM, Toborek M, Watkins BA, Boissonneault GA, Hennig B. Susceptibility to hepatic oxidative stress in rabbits fed different animal and plant fats. J Am Coll Nutr 1996;15:289-94.
- Wu P, Ma G, Li N, Deng Q, Yin Y, Huang R. Investigation of *in vitro* and *in vivo* antioxidant activities of flavonoids rich extract from the berries of *Rhodomyrtus tomentosa* (Ait.) Hassk. Food Chem 2015;173:194-202.
- 21. Olsvik PA, Kristensen T, Waagbø R, Rosseland BO, Tollefsen KE, Baeverfjord G, *et al.* mRNA Expression of antioxidant enzymes (SOD, CAT and GSH-Px) and lipid peroxidative stress in liver of atlantic salmon (*Salmo salar*) exposed to hyperoxic water during smoltification. Comp Biochem Physiol C 2005;141:314-23.
- Afonso MS, de O Silva AM, Carvalho EB, Rivelli DP, Barros SB, Rogero MM, et al. Phenolic compounds from Rosemary (*Rosmarinus officinalis* L.) attenuate oxidative stress and reduce blood cholesterol concentrations in diet-induced hypercholesterolemic rats. Nutr Metab (Lond) 2013;10:19.
- Zhang Z, Chang Q, Zhu M, Huang Y, Ho WK, Chen Z. Characterization of antioxidants present in hawthorn fruits. J Nutr Biochem 2001;12:144-52.
- Schmitt CA, Dirsch VM. Modulation of endothelial nitric oxide by plant-derived products. Nitric Oxide 2009;21:77-91.
- Napoli C, de Nigris F, Williams-Ignarro S, Pignalosa O, Sica V, Ignarro LJ. Nitric oxide and atherosclerosis: An update. Nitric Oxide 2006;15:265-79.
- 26. Wennmalm A. Endothelial nitric oxide and cardiovascular disease. J Intern Med 1994;235:317-27.
- Schini-Kerth VB, Auger C, Kim JH, Etienne-Selloum N, Chataigneau T. Nutritional improvement of the endothelial control of vascular tone by polyphenols: Role of NO and EDHF. Pflugers Arch 2010;459:853-62.
- Shahidi F, Naczk M, Shahidi F, Naczk M. Phenolics in Food and Nutraceuticals. Florida, USA: CRC Press; 2003.
- Feng Y, Sun C, Yuan Y, Zhu Y, Wan J, Firempong CK, et al. Enhanced oral bioavailability and *in vivo* antioxidant activity of chlorogenic acid via liposomal formulation. Int J Pharm 2016;501:342-9.
- Clifford MN, Knight S, Kuhnert N. Discriminating between the six isomers of dicaffeoylquinic acid by LC-MS(n). J Agric Food Chem 2005;53:3821-32.
- Zhang L, Tu Z, Wang H, Fu Z, Wen Q, Chang H, et al. Comparison of different methods for extracting polyphenols from *Ipomoea batatas* leaves, and identification of antioxidant constituents by HPLC-QTOF-MS<sup>2</sup>. Food Res Int 2015;70:101-9.
- Constant HL, Slowing K, Graham JG, Pezzuto JM, Cordell GA, Beecher CW. A general method for the dereplication of flavonoid glycosides utilizing high performance liquid chromatography/mass spectrometric analysis. Phytochem Anal 1997;8:176-80.
- 33. Li X, Xiong Z, Ying X, Cui L, Zhu W, Li F. A rapid ultra-performance liquid chromatographyelectrospray ionization tandem mass spectrometric method for the qualitative and quantitative analysis of the constituents of the flower of *Trollius ledibouri* Reichb. Anal Chim Acta 2006;580:170-80.
- 34. Fabre N, Rustan I, de Hoffmann E, Quetin-Leclercq J. Determination of flavone, flavonol, and flavanone aglycones by negative ion liquid chromatography electrospray ion trap mass spectrometry. J Am Soc Mass Spectrom 2001;12:707-15.