

REFERENCES

- Kelly JP, Kaufman DW, Kelley K, Rosenberg L, Anderson TE, Mitchell AA, *et al.* Recent trends in use of herbal and other natural products. *Arch Intern Med* 2005;165:281-6.
- Hur J. Donguibogam. Seoul, Korea: Namsandang; 2004. p. 98.
- Kim HJ, Choi JH, Lim SW. The defensive effect of Keuibi-tang on the gastric mucous membrane of mouse injured by stress and ethanol. *J Korean Med* 2003;24:155-68.
- Busta I, Xie H, Kim MS. The use of Gui-pi-tang in small animals with immune-mediated blood disorders. *J Vet Clin* 2009;26:181-4.
- Mrozikiewicz PM, Bogacz A, Karasiewicz M, Mikolajczak PL, Ozarowski M, Seremak-Mrozikiewicz A, *et al.* The effect of standardized *Echinacea purpurea* extract on rat cytochrome P450 expression level. *Phytomedicine* 2010;17:830-3.
- Venkataramanan R, Komoroski B, Strom S. *In vitro* and *in vivo* assessment of herb drug interactions. *Life Sci* 2006;78:2105-15.
- Flockhart DA, Oesterheld JR. Cytochrome P450-mediated drug interactions. *Child Adolesc Psychiatr Clin N Am* 2000;9:43-76.
- Izzo AA, Ernst E. Interactions between herbal medicines and prescribed drugs: A systematic review. *Drugs* 2001;61:2163-75.
- Kobayashi K, Urashima K, Shimada N, Chiba K. Selectivities of human cytochrome P450 inhibitors toward rat P450 isoforms: Study with cDNA-expressed systems of the rat. *Drug Metab Dispos* 2003;31:833-6.
- Furge LL, Guengerich FP. Cytochrome P450 enzymes in drug metabolism and chemical toxicology: An introduction. *Biochem Mol Biol Educ* 2006;34:66-74.
- Ahn MJ, Lee MK, Kim YC, Sung SH. The simultaneous determination of coumarins in *Angelica gigas* root by high performance liquid chromatography-diode array detector coupled with electrospray ionization/mass spectrometry. *J Pharm Biomed Anal* 2008;46:258-66.
- Sudjaroen Y, Hull WE, Erben G, Würtele G, Changbumrung S, Ulrich CM, *et al.* Isolation and characterization of ellagitannins as the major polyphenolic components of longan (*Dimocarpus longan* Lour) seeds. *Phytochemistry* 2012;77:226-37.
- Niu C, Zhang J. Quantitative analysis and chromatographic fingerprinting of the Semen Zizyphi Spinosae by ultra-high-performance liquid chromatography coupled with diode-array detector. *J Sep Sci* 2011;34:2989-96.
- Kawashima K, Miyako D, Ishino Y, Makino T, Saito K, Kano Y, *et al.* Anti-stress effects of 3,4,5-trimethoxycinnamic acid, an active constituent of roots of *Polygala tenuifolia* (Onji). *Biol Pharm Bull* 2004;27:1317-9.
- Shan SM, Luo JG, Huang F, Kong LY. Chemical characteristics combined with bioactivity for comprehensive evaluation of *Panax ginseng* C.A. Meyer in different ages and seasons based on HPLC-DAD and chemometric methods. *J Pharm Biomed Anal* 2014;89:76-82.
- Song JZ, Yiu HH, Qiao CF, Han QB, Xu HX. Chemical comparison and classification of radix astragali by determination of isoflavonoids and astragalosides. *J Pharm Biomed Anal* 2008;47:399-406.
- Tsai CJ, Liang JW, Lin HR. Sesquiterpenoids from *Atractylodes macrocephala* act as farnesoid X receptor and progesterone receptor modulators. *Bioorg Med Chem Lett* 2012;22:2326-9.
- Li G, Xu ML, Lee CS, Woo MH, Chang HW, Son JK, *et al.* Cytotoxicity and DNA topoisomerases inhibitory activity of constituents from the sclerotium of *Poria cocos*. *Arch Pharm Res* 2004;27:829-33.
- Li A, Sun A, Liu R. Preparative isolation and purification of costunolide and dehydrocostuslactone from *Aucklandia lappa* Decne by high-speed counter-current chromatography. *J Chromatogr A* 2005;1076:193-7.
- Zhang Q, Ye M. Chemical analysis of the Chinese herbal medicine Gan-Cao (Licorice). *J Chromatogr A* 2009;1216:1954-69.
- Zick SM, Ruffin MT, Djuric Z, Normolle D, Brenner DE. Quantitation of 6-, 8- and 10-gingerols and 6-shogaol in human plasma by high-performance liquid chromatography with electrochemical detection. *Int J Biomed Sci* 2010;6:233-40.
- Pao LH, Hu OY, Fan HY, Lin CC, Liu LC, Huang PW, *et al.* Herb-drug interaction of 50 Chinese herbal medicines on CYP3A4 activity *in vitro* and *in vivo*. *Am J Chin Med* 2012;40:57-73.
- Dong HY, Shao JW, Chen JF, Wang T, Lin FP, Guo YH, *et al.* Transcriptional regulation of cytochrome P450 3A4 by four kinds of traditional Chinese medicines. *Zhongguo Zhong Yao Za Zhi* 2008;33:1014-7, 1089.
- Tai T, Huang X, Su Y, Ji J, Su Y, Jiang Z, *et al.* Glycyrrhizin accelerates the metabolism of triptolide through induction of CYP3A in rats. *J Ethnopharmacol* 2014;152:358-63.
- Martignoni M, Groothuis GM, de Kanter R. Species differences between mouse, rat, dog, monkey and human CYP-mediated drug metabolism, inhibition and induction. *Expert Opin Drug Metab Toxicol* 2006;2:875-94.
- Bogaards JJ, Bertrand M, Jackson P, Oudshoorn MJ, Weaver RJ, van Bladeren PJ, *et al.* Determining the best animal model for human cytochrome P450 activities: A comparison of mouse, rat, rabbit, dog, micropig, monkey and man. *Xenobiotica* 2000;30:1131-52.
- Wu R, Cui X, Dong W, Zhou M, Simms HH, Wang P, *et al.* Suppression of hepatocyte CYP1A2 expression by Kupffer cells via AhR pathway: The central role of proinflammatory cytokines. *Int J Mol Med* 2006;18:339-46.
- Chen J, Murray M, Liddle C, Jiang XM, Farrell GC. Downregulation of male-specific cytochrome P450s 2C11 and 3A2 in bile duct-ligated male rats: Importance to reduced hepatic content of cytochrome P450 in cholestasis. *Hepatology* 1995;22:580-7.
- Cheung C, Gonzalez FJ. Humanized mouse lines and their application for prediction of human drug metabolism and toxicological risk assessment. *J Pharmacol Exp Ther* 2008;327:288-99.
- Badyal DK, Dadhich AP. Cytochrome P450 and drug interactions. *Indian J Pharmacol* 2001;33:248-59.
- Gonzalez FJ. The molecular biology of cytochrome P450s. *Pharmacol Rev* 1988;40:243-88.