





*p*-nitrophenol in NaOH color reagent. The reaction was initiated by treating incubation of the medium containing the pretreated HLM in 0.02 mL of 30 mM uridine diphosphate-glucuronic acid for 15 min at 37°C. The total volume of the reaction mixture was 0.2 mL. The reaction was terminated by adding 0.08 ml of 20% w/v trichloroacetic acid to precipitate the protein. The reaction mixture was centrifuged at 2000 × *g* for 10 min. Then, 0.2 mL supernatant was mixed with 0.8 ml of 5 M NaOH to develop the light yellowish color of the mixture. The absorbance of *p*-nitrophenol consumed was measured at 405 nm. The amount of *p*-nitrophenol consumed through *O*-glucuronidation process in HLM was calculated from the standard curve of *p*-nitrophenol. The calibration solution concentrations of 0, 12.5 μM, 62.5 μM, 125 μM, 312.5 μM, 625 μM, 1250 μM, and 2500 μM were prepared from a 5 mM stock solution. Then, *p*-nitrophenol was transferred into test tubes containing the complete incubation mixture but with heat-inactivated enzymes. The blank was 1 ml of distilled water. The absorbance was measured at 405 nm. Each measurement was made in triplicate. Diclofenac (5 mM) was used as a positive control. The UGT activity was then expressed in nmol *p*-nitrophenol consumed/min/mg protein.

The absorbance of the amount of product formed, or probe consumed in case of UGT activity

(U) was calculated using the equation below:

$$U = (\text{Experimental group} - \text{Blank group}) - (\text{Control group} - \text{Blank group})$$

The positive control, diclofenac (5 mM), was included to assess the test validity in UGT assay. All test groups were pretreated with 0.008 mL Triton-X100 (0.75% v/v) instead of 0.01 mL Triton X-100 (0.375% v/v) as per other tests for 3 min to offset the membrane latency prior to the experiment.

## Attenuated total reflectance Fourier transform infrared fingerprints

Attenuated total reflectance Fourier transform infrared (ATR-FTIR) spectra of the aqueous extract and the juice were obtained with Nicolet™ iS5 FTIR spectrometer controlled by OMNIC software (Thermo Fisher Scientific, Waltham, Massachusetts, USA) for spectra collection and TQ Analyst software for data processing (Thermo Fisher Scientific, Waltham, Massachusetts, USA). The instrument is equipped with iD5 ATR accessory featuring a top plate diamond crystal with a fixed angle of incidence of 42°. The qualitative FTIR spectroscopy analysis was conducted in the mid-IR range of 4000–400/cm at a resolution of 4/cm with 14 scans. The background spectrum was recorded before obtaining the spectra of the samples.

## Data analysis

Data were presented as mean ± standard error of the mean. The data for the present study were analyzed using GraphPad Prism 5 (GraphPad Software, San Diego, California, USA) using analysis of variance following Dunnett's test to examine the significant differences of each treatment group (test and positive control group) compared to the negative control group. With a confidence level of 95% ( $P < 0.05$ ), the difference would be considered as statistically significant when the treatment group was compared to the control group.

## RESULTS AND DISCUSSION

### Extraction yield and total flavonoid contents

The percentage yields and TFCs of the fresh leaves juice and dried leaves aqueous extracts of *C. nutans* are shown in Table 1. The percentage yield of the aqueous extract was greater than juice. The high yield of *C. nutans* in water extracts was probably due to the high

solubility of major components of *C. nutans* in the polar solvent. The yields of fresh juices of 11 herbs were reported to be lower than those of ethanol and water extracts.<sup>[16]</sup> Thus, mechanical forces in the form of pressing and squeezing fresh leaves in preparing leaves preparation may be inferior to the chemical solvent extraction method. Although mechanical forces allowed leaves to breakdown and release the liquid soluble secondary metabolites, chemical extraction methods have higher efficiency by using aqueous and methanol solvent to improve the yield of extraction.<sup>[17,18]</sup>


The flavonoid contents of the two preparations were significantly different [Table 1]. TFC was higher in the leaf juice than the cold aqueous extract. Despite the higher yield of the aqueous extract, its TFC was lower than that of the juice. Total flavonoids content of *C. nutans* herbal tea prepared by using microwave oven dried method was reported to be lower than that of freeze-dried method suggesting that drying process can contribute to the loss in flavonoids.<sup>[19]</sup> Flavonoids have the potential to modulate CYP450 activities which can lead to the production of environmental carcinogens. Thus, the presence of flavonoids in *C. nutans* preparations may contribute to CYP3A4 and CYP2E1 and UGT enzymes activity. Therefore, the *in vitro* UGT and CYP 450 enzymes activities in HLMs were investigated.

### *In vitro* cytochrome P3A4 activity

The change in CYP3A4 hepatic drug metabolizing enzyme activity of aqueous leaf extract and leaf juice of *C. nutans* was evaluated. Preliminary experiment to optimize assay condition for CYP3A4 showed linearity in HLM activity. The assay for CYP3A4 conducted showed positive results in positive control. Figure 1 shows the inhibitory effect of *C. nutans* on CYP3A4 through aminopyrine *N*-demethylation in HLMs at the concentrations tested. Fresh leaf juice and aqueous leaves extract of *C. nutans* at 10 ng/mL and 1000 ng/mL showed statistically significant ( $P < 0.05$ ) inhibitory effect on CYP3A4 in HLMs as compared to the control. Differences in the

**Table 1:** The percentage yield and total flavonoid content of *Clinacanthus nutans* leaves preparations

|   | Cold aqueous extract | Fresh leaves juice |
|---|----------------------|--------------------|
| Weight of leaves (g)                                  | 75.00                | 305.98             |
| Percentage yield of extraction (% w/w)                | 6.87                 | 0.55               |
| Total flavonoid content (mg catechin Eq/g dry weight) | 245.48               | 283.33             |



**Figure 1:** *In vitro* effect of leaf juice and aqueous extract of *Clinacanthus nutans* on the activity of cytochrome P3A4 in human liver microsomes.  $N = 3$ ; data values were presented as mean ± standard error of the mean; analyzed using Dunnett's test; \* $P < 0.05$ , a significant difference compared to the negative control

flavonoid content, the number of the hydroxyl group and types of interaction with CYP3A4 could contribute to higher inhibition effect of the cold aqueous extract on CYP3A4 as compared to the juice.

The inhibitory effect could be ascribed to its flavonoid constituents.<sup>[18,20,21]</sup> Inhibition of CYP3A4 in human had been well documented for flavonoids.<sup>[11,22]</sup> Despite the very large and flexible active site for CYP3A4, substrate binding was mainly based on hydrophobicity with some steric interactions.<sup>[20]</sup> Flavones and flavonols inhibit enzyme activity by directly binding to CYP3A4 isoforms that involved in xenobiotics metabolisms.<sup>[18]</sup> Flavones which bound to CYP3A4 undergo hydroxylation or demethylation to inhibit the enzyme activity. Inhibition of CYP3A4 will lead to xenobiotics that metabolized by the same enzyme to slow down in its metabolism. Thus, concomitant use of a drug metabolized through CYP3A4 and *C. nutans* preparation will cause the xenobiotics to remain in the body system for a longer period. This may potentially cause toxicity in those drugs with major side effects at a low dose. From data on flavonoid-CYP interactions, flavonoids that possess hydroxyl groups inhibit CYP activity, whereas those lacking hydroxyl groups can stimulate enzyme activity.<sup>[20,21]</sup>

### In vitro cytochrome P2E1 activity

The CYP2E1 hepatic drug metabolizing enzyme (carcinogens activating enzyme) activity of the aqueous extract, and the juice was investigated by measuring the formation of *p*-nitro catechol (product) from hydroxylation of *p*-nitrophenol (substrate) through *p*-nitrophenol hydroxylation of HLMs. The preliminary experiment to optimize assay condition for CYP2E1 showed linearity in the human liver microsomal activity. The assays for CYP2E1 conducted showed positive results for the positive control. The results of CYP2E1 activity is shown in Figure 2. Only the leaf juice at 1000 ng/mL showed a statistically significant ( $P < 0.01$ ) inhibitory effect on CYP2E1 in HLMs as compared to the control group. The aqueous extract concentration at 1000 ng/mL showed a decrease in activity; however, the reduction in activity was not statistically significant ( $P > 0.05$ ) when compared to the control value. Similarly, at the concentration of 10 ng/mL juice and aqueous extract demonstrated a reduction in CYP2E1 activity, which was not significant ( $P > 0.05$ ) compared to control value. Only the juice

at high concentration inhibited significant effect on CYP2E1 activity. The inhibition of CYP2E1 activity may have an inhibitory effect on carcinogenesis initiation.<sup>[23,24]</sup>


### In vitro UDP-glucuronosyltransferase activity

The UGT activity of aqueous extract and juice of *C. nutans* leaves were measured from the consumption of *p*-nitrophenol (substrate) through *O*-glucuronidation process in HLMs which represented by *p*-nitrocatechol consumed/min/mg. Preliminary experiment to optimize assay condition for UGT showed linearity in HLM activity. The assay showed positive results in positive control. *C. nutans* preparations showed significant activation activity on UGT (Phase II hepatic drug metabolizing enzyme) in HLM [Figure 3]. The aqueous extract of *C. nutans* showed significant ( $P < 0.05$ ) activation on UGT activity at the concentration of 1000 ng/mL as compared to the negative control. At the concentrations of 10 ng/mL, the juice and the aqueous extract showed activation of UGT activity, but the activation was not significant ( $P > 0.05$ ) as compared to the control value. The activation of UGT activity indicates that simultaneous use of a drug metabolized through UGT and *C. nutans* preparations may result in the faster detoxification process and reduction in its potency. These extracts were found to contain flavonoids which have been reported to be associated with UGT induction in human hepatic cell lines and glucuronidation in UGT1A1 through a nonaryl hydrocarbon receptor-mediated mechanism.<sup>[18,20,25]</sup>


### Attenuated total reflectance fourier transform infrared fingerprints

The ATR-FTIR fingerprints are useful for the identification of functional groups of compounds present in the samples based on the characteristic IR vibrations frequencies of the functional groups. ATR-FTIR spectra of the aqueous extract and the juice are shown in Figure 4. The ATR-FTIR fingerprints of the two samples were similar.


The spectra showed an intense hydroxyl (-OH) band at 3200–3600/cm, C-H stretch at a frequency range of 2900–3000/cm, Carbonyl (C = O) stretch at a frequency of 1605–1630/cm, and C-O stretch at a frequency of 1080–1050/cm. The frequency range of 3200–3600/cm was the stretching vibration for hydroxyl group for alcohol, including phenol. Besides that, the frequency of 1628/cm was assigned to the carbonyl group for carbonyl compounds. The presence of hydroxyl



**Figure 2:** In vitro effect of leaf juice and aqueous extract of *Clinacanthus nutans* on the activity of cytochrome P2E1 in human liver microsomes.  $N = 3$ ; data values were presented as mean  $\pm$  standard error of the mean; analyzed using Dunnett's test; \* $P < 0.05$ , \*\* $P < 0.01$  significant difference compared to the control and  $\Phi$  no significant difference compared to the negative control



**Figure 3:** In vitro effect of leaf juice and aqueous extract of *Clinacanthus nutans* on the activity of UDP-glucuronosyltransferase in human liver microsomes.  $N = 3$ ; data values were presented as mean  $\pm$  standard error of the mean; analysed using Dunnett's test; \* $P < 0.05$ , \*\* $P < 0.01$  significant difference compared to the negative control and  $\Phi$  no significant difference compared to the negative control



**Figure 4:** Fourier transform infrared spectra *Clinacanthus nutans* leaf. (a) aqueous extract; (b) juice

band and carbonyl stretch suggested compounds in the extracts have a hydroxyl functional group and carbonyl functional group. Phenolic and flavonoids compounds are compounds containing hydroxyl and carbonyl group as functional groups of chemical constituents in the extract. C-glycosyl flavones in the extract have a hydroxyl functional group.

## CONCLUSION

*C. nutans* leaf preparations may interact with CYP3A4 and CYP2E1 enzymes, and it should be used with caution with drugs that metabolized by these enzymes to avoid potential adverse effects. The leaf preparation also activated UGT catalyzed metabolism which may result in a reduction of the potency of the drug metabolized by UGT pathway. Despite the findings of this study, the herbs could be safe for consumption since metabolism occurs through various enzymes than a single enzyme pathway. This *in vitro* study using HLM provides important drug interaction screening and research information by providing a fundamental understanding of safety during the interaction between the herb and drug. This *in vitro* study excludes drug transporter effect; nevertheless, the information can be beneficial in preclinical discovery stages in drug development. Data regarding the CYP450 inhibitory activities of *C. nutans* have shown the additional information needed to be considered in designing animal studies to predict the effect that might be anticipated in humans. *In vivo* effect of leaf extract and juice of *C. nutans* leaves on drug-metabolizing enzymes activity for quantitative measurement in quantitative prediction of human biotransformation are in progress.

## Financial support and sponsorship

The authors are grateful for the financial supports from CERVIE, UCSI, University research funding (Proj-In-FPS-004) and Universiti Sains Malaysia Research Grant Scheme (304/PFARMASI/6331040) to carry out the work successfully.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Bardia A, Nisly NL, Zimmerman MB, Gryzlak BM, Wallace RB. Use of herbs among adults based on evidence-based indications: Findings from the national health interview survey. *Mayo Clin Proc* 2007;82:561-6.
- Chin JH, Hussin AH. Effect of methanol leaf extract of *Orthosiphon stamineus* Benth. on hepatic drug metabolising enzymes in Sprague Dawley (SD) rats. *J Biosci* 2008;19:21-31.
- Mohamed I, Shuid A, Borhanuddin B, Fozi N. The application of phytomedicine in modern drug development. *Int J Herb Plant Med* 2012;1:1-9.
- Ekor M. The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol* 2014;4:177.
- Chow HH, Hakim IA, Vining DR, Crowell JA, Cordova CA, Chew WM, *et al.* Effects of repeated green tea catechin administration on human cytochrome P450 activity. *Cancer Epidemiol Biomarkers Prev* 2006;15:2473-6.
- Levéque D, Jehl F. Molecular pharmacokinetics of catharanthus (vinca) alkaloids. *J Clin Pharmacol* 2007;47:579-88.
- Brantley SJ, Graf TN, Oberlies NH, Paine MF. A systematic approach to evaluate herb-drug interaction mechanisms: Investigation of milk thistle extracts and eight isolated constituents as CYP3A inhibitors. *Drug Metab Dispos* 2013;41:1662-70.
- Tsai HH, Lin HW, Simon Pickard A, Tsai HY, Mahady GB. Evaluation of documented drug interactions and contraindications associated with herbs and dietary supplements: A systematic literature review. *Int J Clin Pract* 2012;66:1056-78.
- Quah SY, Chin JH, Akowuah GA, Khalivulla SI, Yeong SW, Sabu MC. Cytotoxicity and cytochrome P450 inhibitory activities of *Clinacanthus nutans*. *Drug Metab Pers Ther* 2017;32:59-65.
- Chan LW, Cheah EL, Saw CL, Weng W, Heng PW. Antimicrobial and antioxidant activities of cortex magnoliae officinalis and some other medicinal plants commonly used in South-East Asia. *Chin Med* 2008;3:15.
- Nicolas JM, Collart P, Gerin B, Mather G, Trager W, Levy R, *et al.* *In vitro* evaluation of potential drug interactions with levetiracetam, a new antiepileptic agent. *Drug Metab Dispos* 1999;27:250-4.
- Gould KS, Lister C. Flavonoids: Chemistry, biochemistry, and applications. In: Anderson OM, Markham KR, editors. *Flavonoid Functions in Plants*. New York: CRC Press; 2005. p. 1212-6.
- Cheng Q, Guengerich FP. Identification of endogenous substrates of orphan cytochrome P450 enzymes through the use of untargeted metabolomics approaches. In: Phillips IR, Shephard EA, de Montellano PO, editors. *Cytochrome P450 Protocols. Methods in Molecular Biology (Methods and Protocols)*. Vol. 987. New Jersey: Humana Press; 2013. p. 71-7.
- Chao P, Hsiu S, Hou Y. Flavonoids in herbs: Biological fates and potential interactions with xenobiotics. *J Food Drug Anal* 2002;10:219-28.
- Hodek P, Trefil P, Stiborová M. Flavonoids-potent and versatile biologically active compounds interacting with cytochromes P450. *Chem Biol Interact* 2002;139:1-21.
- Shimada T, Tanaka K, Takenaka S, Murayama N, Martin MV, Foroozesh MK, *et al.* Structure-function relationships of inhibition of human cytochromes P450 1A1, 1A2, 1B1, 2C9, and 3A4 by 33 flavonoid derivatives. *Chem Res Toxicol* 2010;23:1921-35.
- Yao HT, Lin JH, Chiang MT, Chiang W, Luo MN, Lii CK. Suppressive effect of the ethanolic extract of adlay bran on cytochrome P-450 enzymes in rat liver and lungs. *J Agric Food Chem* 2011;59:4306-14.
- Ye Q, Lian F, Chavez PR, Chung J, Ling W, Qin H, *et al.* Cytochrome P450 2E1 inhibition prevents hepatic carcinogenesis induced by diethylnitrosamine in alcohol-fed rats. *Hepatobiliary Surg Nutr* 2012;1:5-18.
- Bilecová-Rabajdová M, Birková A, Urban P, Gregová K, Durovcová E, Mareková M, *et al.* Naturally occurring substances and their role in chemo-protective effects. *Cent Eur J Public Health* 2013;21:213-9.

20. Walle UK, Walle T. Induction of human UDP-glucuronosyltransferase UGT1A1 by flavonoids-structural requirements. *Drug Metab Dispos* 2002;30:564-9.
21. Pekal A, Pyrzynska K. Evaluation of aluminium complexation reaction for flavonoid content assay. *Food Anal Methods* 2014;7:1776-82.
22. Van Vleet TR, Bombick DW, Coulombe RA Jr. Inhibition of human cytochrome P450 2E1 by nicotine, cotinine, and aqueous cigarette tar extract *in vitro*. *Toxicol Sci* 2001;64:185-91.
23. Cederbaum AI. Methodology to assay CYP2E1 mixed function oxidase catalytic activity and its induction. *Redox Biol* 2014;2:1048-54.
24. Oliveira EJ, Watson DG. *In vitro* glucuronidation of kaempferol and quercetin by human UGT-1A9 microsomes. *FEBS Lett* 2000;471:1-6.
25. Desai AA, Innocenti F, Ratain MJ. UGT pharmacogenomics: Implications for cancer risk and cancer therapeutics. *Pharmacogenetics* 2003;13:517-23.