

# Preventive Effect of Crude Polysaccharide Extract from Chinese Wolfberry against Hyperglycemia-Induced Oxidative Stress and Inflammation in Streptozotocin-Induced Diabetic Rats

Junjie Li<sup>1</sup>, Yong Zhang<sup>1</sup>, Linshan Jiao<sup>1</sup>, Opeyemi Joshua Olatunji<sup>2</sup>, Bing He<sup>1</sup>

<sup>1</sup>Department of Nephrology, Zhengzhou Central Hospital Affiliated to Zhengzhou University, Zhengzhou City, Henan Province, China, <sup>2</sup>Faculty of Traditional Thai Medicine, Prince of Songkla University, Hat Yai, Thailand

Submitted: 11-04-2019

Revised: 28-05-2019

Published: 19-09-2019


## ABSTRACT

**Background:** The protective effects of the crude polysaccharide extract Polysaccharide *Lycium chinense* (PLC) from the leaves of *Lycium chinense* (Chinese wolfberry) were evaluated in streptozotocin (STZ)-induced diabetic rats. **Materials and Methods:** Diabetes was induced in rats by administering STZ (60 mg/kg, i.p), and diabetic rats were orally treated with 100 or 400 mg/kg of PLC extract for 4 weeks. **Results:** Diabetic rats showed high fasting blood glucose levels, altered serum lipid profile; triglycerides, total cholesterol, low-density lipoprotein cholesterol, high-density lipoprotein cholesterol, and liver function enzymes; and alanine aminotransferase, aspartate aminotransferase, and alkaline phosphatase. In addition, oxidative stress and pro-inflammatory cytokines were elevated in the kidney and liver tissues of diabetic rats. After treatment with PLC, it was observed that hyperglycemia, hyperlipidemia, and liver and kidney functions were restored to almost normal while enzymatic antioxidant levels of glutathione peroxidase, superoxide dismutase, and catalase were significantly increased, in addition to remarkable reduction in malondialdehyde and pro-inflammatory cytokines. **Conclusion:** These results demonstrated the protective effect of PLC may be mediated by its antioxidant and anti-inflammatory effects and may be employed as a therapy for preventing diabetes and its complications.

**Key words:** Hyperglycemia, inflammation, *Lycium chinense*, oxidative stress, polysaccharide

## SUMMARY

- Wolfberry polysaccharide extract at 100 and 400 mg/kg showed protective effect against streptozotocin-induced diabetes
- Antidiabetic effect is mediated by alleviation of oxidative stress and inflammation.



**Abbreviations used:** PLC: Polysaccharide *Lycium chinense*; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; ALP: Alkaline phosphatase; TG: Triglyceride; TC: Total cholesterol; LDLC: Low-density lipoprotein cholesterol; HDLC: High-density lipoprotein cholesterol; BUN: Blood urea nitrogen; SOD: Superoxide dismutase; CAT: Catalase; GSH-Px: Glutathione peroxidase; MDA: Malondialdehyde; TNF- $\alpha$ : Tumor necrosis factor alpha; IL-1 $\beta$ : Interleukin 1 beta; IL-6: Interleukin 6.

## Correspondence:

Dr. Bing He,  
Department of Nephrology,  
Zhengzhou Central Hospital Affiliated to  
Zhengzhou University, Zhengzhou City,  
Henan Province 450007, China.  
E-mail: hebingzhongxin@sina.com  
DOI: 10.4103/pm.pm\_164\_19

## Access this article online

Website: [www.phcog.com](http://www.phcog.com)

## Quick Response Code:



## INTRODUCTION

In recent years, metabolic diseases have become a major public health problem, with a worldwide prevalence of 20%–30% of the population.<sup>[1]</sup> Metabolic diseases depict several abnormalities including hyperglycemia, dyslipidemia, high blood pressure, and increased risk of cardiovascular diseases.<sup>[2]</sup> One of the most dangerous and life-threatening metabolic disorders is diabetes mellitus, a disease that affects all age groups, and it is clinically characterized by persistent high blood glucose due to deficiency in the action of insulin (insulin resistance or insufficient insulin production). Diabetes is often associated with life-threatening damages to vital organs in the body, such as the eye, kidney, liver, and nerves. The damages accrued to these organs in diabetic situations significantly contribute to the severity of the disease.<sup>[3-5]</sup>

Hyperglycemia-induced oxidative stress and inflammation which are instigated by the overproduction of reactive oxygen species (ROS)

and reduced antioxidant defense system are vital contributors to the progression of diabetes mellitus and diabetic complications.<sup>[4,6]</sup> Accordingly, substances with antioxidants and anti-inflammatory effects could be of potential therapeutic benefit against diabetes mellitus.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprints@medknow.com](mailto:reprints@medknow.com)

**Cite this article as:** Li J, Zhang Y, Jiao L, Olatunji OJ, He B. Preventive effect of crude polysaccharide extract from Chinese wolfberry against hyperglycemia-induced oxidative stress and inflammation in streptozotocin-induced diabetic rats. *Phcog Mag* 2019;15:638-44.

The use of natural products from medicinal plants as food and medicine is becoming significantly important in recent years. A significant number of people in both rural and urban areas of the world use some form of medicine from natural sources as part of their primary health care for the prevention or treatment of various diseases. The reason behind this has been attributed to the accessibility, cost, efficacy, and lesser undesirable effect as compared to most of the synthetic drugs.<sup>[7,8]</sup> Polysaccharides are one of the important naturally occurring bioactive constituents found in plants, algae, micro-organisms, and animals. They display a wide range of structural and biological characteristics, which make them a center of attraction in recent years. Polysaccharides are a group of biological macromolecules which are made up of ten or more monosaccharides.<sup>[9,10]</sup> Polysaccharides have gained prominent attention as components of functional food and nutraceuticals due to their diverse pharmacological properties, non-toxic nature, and effectiveness.<sup>[9,11,12]</sup> A growing number of studies have portrayed efficacy of purified and crude form of polysaccharides in diabetic situations. The antidiabetic effects of these reported polysaccharides have been attributed to various mechanisms including improving glucose metabolism, lipid, and metabolism, alleviating pancreatic  $\beta$ -cell dysfunction, and inhibiting  $\alpha$ -amylase and  $\alpha$ -glucosidase. In addition, some polysaccharide has been reported to inhibit oxidative and inflammatory pathways responsible for complications associated with diabetes.<sup>[10,13-16]</sup>

The genus *Lycium* belongs to the family *Solanaceae*, comprising mainly edible perennial shrubs with seven identified species. Of these, two varieties (*Lycium chinense* and *Lycium barbarum*) are extensively cultivated and used in China. *L. chinense*, popularly known as Chinese boxthorn or Chinese wolfberry, is an important traditional Chinese medicinal plant due to its application as food and medicine. The leaves of the plant have been pharmacologically explored as a neuroprotective, antimicrobial, antioxidant, anti-angiotensin I-converting enzyme, antiulcer, and antidiabetic agent.<sup>[8,17,18]</sup> The leaf of the plant is a very rich source of a variety of bioactive active compounds, such as polyphenols, diterpene glycosides, with anolides, aliphatic acids, and polysaccharides.<sup>[18-20]</sup> Numerous studies have portrayed the significant contributions of polysaccharides from medicinal plants and edible fungi as an antidiabetic;<sup>[7,16,21]</sup> however, polysaccharides from *L. chinense* have not been extensively explored for their pharmacological benefits. Therefore, the crude polysaccharide extract from the leaves of *L. chinense* was extracted and evaluated for this study. Thus, this present study was aimed at revealing the protective effect of *L. chinense* polysaccharide in streptozotocin (STZ)-induced diabetic rats.

## MATERIALS AND METHODS

### Plant material

The leaves of the plant were collected from Henan Province, China, in 2016. The plant was identified at the School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou, Henan. A voucher specimen (No. 2016010 LC) was stored at the herbarium of the university.

### Extraction of polysaccharide

The leaves of *L. chinense* was powdered, was depigmented using acetone, and was subjected to distilled water extraction using Soxhlet apparatus at 90°C for 3 h. The water extract was centrifuged at 5000 rpm for 15 min, and the resulting supernatant was concentrated under reduced pressure and further precipitated with 95% ethanol. The precipitate obtained was further deproteinized using Savage method,<sup>[22]</sup> dialyzed, and lyophilized to obtain the crude polysaccharide (PLC) which was stored at 4°C until further use.

## Animals and induction of diabetic rat model

Male Wistar rats (150–200 g) were kept in a room maintained at a temperature of 24°C  $\pm$  2°C, relative humidity of 55%  $\pm$  5%, and a 12-h light/dark cycle. The rats were allowed 7 days of acclimatization and were fed with a standard rat diet during the experimental period. The animals also had free access to water *ad libitum*. The experimental protocol was approved by the Animals Ethics Committee of Zhengzhou Central Hospital (Ethics number: zxyy20181101). After 7 days of acclimatization, the rats were fasted for 12 h and administered with a single dose of freshly prepared STZ (60 mg/kg, i.p) in sodium citrate buffer solution (pH 4.5) to induce diabetes. Fasting blood glucose (FBG) levels of the rats were measured after 72 h of STZ administration, and rats with Fasting blood glucose (FBG) levels  $\geq$ 15.0 mmol/L were considered diabetic and included in further experiments.

## Experimental design

Rats were randomly allotted into five groups (6 rats per group) as indicated below:

- Normal control group (NCG) received normal saline
- Diabetic control group (DCG) received normal saline
- Glibenclamide-treated group treated with glibenclamide (20 mg/kg)
- PLC 100-treated group (PLC100) treated with PLC (100 mg/kg)
- PLC 400-treated group (PLC400) treated with PLC (400 mg/kg).

The rats were orally treated with PLC and glibenclamide once daily for 4 weeks. Glibenclamide, the standard medication for treating diabetic patients, was used as the positive control drug. The rats were overnight fasted on the last day of the experiment; the body weight and FBG were measured using Accu-Chek active glucometer (Roche Diagnostics, Mannheim, Germany). At the end of the experiments, the animals were anesthetized and sacrificed by cervical dislocation. Blood, kidney, and liver samples were obtained. The blood collected was centrifuged at 5000 rpm for 15 min to obtain the serum. The kidney and liver samples were homogenized in phosphate buffer (0.1 M) and centrifuged at 5000 rpm for 15 min. The supernatant obtained after centrifuging was used for the determination of biochemical parameters.

## Biochemical analysis

The serum was used for the determination of liver function enzymes namely alanine aminotransferase (ALT), aspartate aminotransferase (AST), and alkaline phosphatase (ALP); serum lipids namely triglycerides (TG), total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and high-density lipoprotein cholesterol (HDL-C); and blood urea nitrogen (BUN), urea, and creatinine with the aid of the corresponding commercial ELISA assay kits. The homogenates from the kidney and liver were used for assessing the antioxidant enzyme levels (superoxide dismutase [SOD], glutathione peroxidase [GSH-Px] and catalase [CAT]), lipid peroxidation (malondialdehyde [MDA]) levels, and pro-inflammatory cytokines (tumor necrosis factor alpha [TNF- $\alpha$ ], interleukin 1 beta [IL-1 $\beta$ ], and interleukin-6 [IL-6]) using commercially available kits.

## Statistical analysis

Graphpad Prism Software (Version 7.0, GraphPad Software Inc., San Diego, California, USA) was employed for statistical analysis. All the data are presented as mean  $\pm$  standard deviation. Statistical analyses were performed by one-way ANOVA followed by Tukey *post hoc* test.  $P < 0.05$  was considered statistically significant.

**RESULTS**

The effect of PLC on the body weight, FBG, and insulin levels in STZ-induced diabetic rats is displayed in Table 1. Comparison of the body weight, insulin, and FBG levels of the DCG with the NCG clearly indicated significant decrease in body weight and insulin levels and a marked increase in the FBG level of the DCG of rats. On treatment with 100 and 400 mg/kg of PLC, the FBG level was apparently decreased, while the body weight and insulin levels were higher than those of the rats in the DCG.

After 4 weeks of treatment with PLC, the serum lipid levels namely LDL-C, TG, and TC were observed to be obviously decreased in PLC100 and PLC400 groups. In addition, HDL-C level was observed to be markedly reduced in the DCG as compared to NCG. After the treatment of diabetic rats with PLC (100 and 400 mg/kg), the serum levels of HDL-C were observed to be evidently higher than those of the untreated DCG [Figure 1]. The effect of PLC on serum level of liver function indexes in the diabetic rats is presented in Table 2. The serum ALP, ALT, and AST levels were significantly higher in the untreated diabetic rats (DCG) when compared to the NCG. However, in the PLC-treated rats, a significant and concentration-dependent decrease in the levels of ALP, ALT, and AST was observed.

The results obtained from this study indicated that serum levels of creatinine, urea, and BUN were remarkably increased in the untreated diabetic rats when compared with the normal control (NC) rats. After 4 weeks of treatment with PLC, the levels of creatinine, urea, and BUN in the serum were significantly decreased in all treated diabetic rat groups [Table 3].

The untreated diabetic rats displayed marked decrease in kidney and liver antioxidant enzyme activities of SOD, GSH-Px, and CAT, while the lipid peroxidation product MDA was apparently increased when compared to the NC rats [Table 4]. Treatment of the diabetic rats with 100 and 400 mg/kg PLC markedly increased the activities of the antioxidant enzymes as well as decreased the level of MDA in the kidney and liver tissues [Table 4].

The anti-inflammatory effect of PLC was further evaluated in the kidney tissues of diabetic rats. As clearly shown in Figure 2, the levels of

pro-inflammatory cytokines TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 were significantly increased in the kidney of diabetic rats when compared with the NCG. After treatment of the diabetic rats with PLC (100 and 400 mg/kg), the levels of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 in the kidney were significantly reduced [Figure 2].

**Table 1:** Effect of PLC on body weight, fasting blood glucose and insulin level in diabetic rats


Groups	Body weight (g)	FBG (mmol/L)	Insulin ( $\mu$ U/mL)
NCG	280.20 $\pm$ 6.11	6.90 $\pm$ 0.30	21.20 $\pm$ 1.42
DCG	170.19 $\pm$ 3.45 <sup>#</sup>	21.40 $\pm$ 1.04 <sup>#</sup>	6.80 $\pm$ 0.78 <sup>#</sup>
DLG	243.57 $\pm$ 7.01**	9.20 $\pm$ 0.69**	17.80 $\pm$ 1.15**
PLC 100	212.05 $\pm$ 5.44**	14.60 $\pm$ 1.16**	11.30 $\pm$ 1.22**
PLC 400	235.70 $\pm$ 5.82**	11.50 $\pm$ 0.86**	14.10 $\pm$ 0.92**

Values are expressed as mean $\pm$ SD (n=6). <sup>#</sup>P<0.05 versus normal control group; \*\*P<0.05 versus diabetic control group. FBG: Fasting blood glucose; NCG: Normal control group; DCG: Diabetic control group; DLG: Diabetic rats treated with glibenclamide; SD: Standard deviation; PLC 100: Diabetic rats treated with *Lycium chinense* polysaccharide (100 mg/kg); PLC 400: Diabetic rats treated with *Lycium chinense* polysaccharide (400 mg/kg)

**Table 2:** Effect of PLC on aspartate aminotransferase, alkaline phosphatase, and alanine aminotransferase level in diabetic rats

Groups	AST (U/L)	ALP (U/L)	ALT (U/L)
NCG	122.73 $\pm$ 7.04	11.26 $\pm$ 1.69	37.09 $\pm$ 2.01
DCG	258.20 $\pm$ 11.22 <sup>#</sup>	35.59 $\pm$ 3.71 <sup>#</sup>	95.04 $\pm$ 2.66 <sup>#</sup>
DLG	148.40 $\pm$ 8.26**	19.15 $\pm$ 2.04**	48.37 $\pm$ 3.68**
PLC 100	190.81 $\pm$ 10.61**	28.05 $\pm$ 2.84**	70.49 $\pm$ 5.08**
PLC 400	159.38 $\pm$ 9.19**	23.05 $\pm$ 3.03**	56.70 $\pm$ 4.40**

Values are expressed as mean $\pm$ SD (n=6). <sup>#</sup>P<0.05 versus normal control group; \*\*P<0.05 versus diabetic control group. AST: Aspartate aminotransferase; ALP: Alkaline phosphatase; ALT: Alanine aminotransferase; NCG: Normal control group; DCG: Diabetic control group; DLG: Diabetic rats treated with glibenclamide; SD: Standard deviation; PLC 100: Diabetic rats treated with *Lycium chinense* polysaccharide (100 mg/kg); PLC 400: Diabetic rats treated with *Lycium chinense* polysaccharide (400 mg/kg)



**Figure 1:** Effect of PLC on serum lipids; (a) TG, (b) TC, (c) LDL-C, and (d) HDL-C in STZ-induced diabetic rats. Values are expressed as mean  $\pm$  SD (n = 6). <sup>#</sup>P < 0.05 versus normal control group; \*\*P < 0.05 versus diabetic control group. NCG: Normal control group; DCG: Diabetic control group; DLG: Diabetic rats treated with glibenclamide; PLC 100: Diabetic rats treated with *Lycium chinense* polysaccharide (100 mg/kg); PLC 400: Diabetic rats treated with *Lycium chinense* polysaccharide (400 mg/kg); TG: Triglyceride; TC: Total cholesterol; LDL-C: Low-density lipoprotein cholesterol; HDL-C: High-density lipoprotein cholesterol; STZ: Streptozotocin; SD: Standard deviation



## DISCUSSION

Bioactive substances from natural origin have been long explored as potential sources of antidiabetic agents. These bioactive natural substances have the ability to function in a multifacet dimension by interacting with various molecular pathways that suppress or attenuate diseases, thus making them prolific alternative to the synthetic drugs, which often act by single mechanism of action and also have undesirable side effects.<sup>[23]</sup> This study assessed the effects of *L. chinense* polysaccharide extract on blood glucose, lipids, and antioxidant status in STZ-induced diabetic rats. As an alkylating compound, STZ destroys pancreatic  $\beta$ -cells by activating DNA strand breakage as well as instigating the excessive production of ROS, especially superoxide radicals. The resultant toxic effect of STZ leads to hyperglycemia and hypoinsulinemia.<sup>[24,25]</sup> Hyperglycemia, which is the chief hallmark of diabetes, is mainly responsible for the chronic organ damages experienced by diabetic patients. As such, an effective glycemic control is paramount to mitigate or impede diabetic complications.<sup>[26,27]</sup> It was observed in this present study that the administration of STZ to the rats significantly increased the blood glucose level and caused a decline in body weight of the rats, which were in consensus with previous reports.<sup>[7,28-30]</sup> Diabetic rats treated with PLC extract had their blood glucose and declined insulin levels ameliorated. Furthermore, the decline in the body weight of diabetic rats was restored by PLC which indicated improved glycemia, thus suggesting that PLC had excellent antihyperglycemic properties.

One of the key abnormalities experienced in diabetes is dyslipidemia and dyslipoproteinemia, a condition that occurs as a result of impairment in lipid metabolism. Dyslipidemia and dyslipoproteinemia

are major risk factors for atherosclerosis and cardiovascular diseases, which are one of the secondary complications of diabetes.<sup>[31-33]</sup> Hyperglycemic-induced dyslipidemia and dyslipoproteinemia are characterized by elevated levels of TC, TG, and LDL-C and decreased levels of HDL-C. We observed high levels of serum lipids in the untreated STZ-induced diabetic rats in this study, which was similar to results obtained in previous studies.<sup>[34-36]</sup> However, PLC-treated rats exhibited abated levels of TG, TC, and LDL-C. These results are illustrative of the beneficial actions of PLC as an antihyperlipidemic agent.

Accumulating evidence has continued to implicate ROS and oxidative stress as critical factors in the development and progression of diabetes and its associated complications. The excessive generation of ROS and the incapability of the enzymatic and nonenzymatic antioxidant systems in the cells to scavenge the reactive species generated lead to an imbalance in the oxidant/antioxidant system, resulting in oxidative stress. Persistent hyperglycemia can induce the overproduction of ROS, and STZ-induced diabetes can lead to impairment of the antioxidant defense system through the production of excess intracellular ROS as well as the imposition of oxidative stress.<sup>[37,38]</sup> Cellular antioxidant enzymes such as GSH-Px, CAT, and SOD primarily act as the foremost defense mechanism against the harmful effect of ROS, thereby affording protection against cellular damages.<sup>[7,39,40]</sup> On the other hand, lipid peroxidation end product MDA is considered as a suitable index for determining oxidative stress/oxidative damage, as high level of MDA is a reflection of loss or decrease in antioxidant enzymes capacity.<sup>[41,42]</sup> The results from this study indicated a decrease in the antioxidant enzyme activities (SOD, CAT, and GSH-Px), with a corresponding increase in MDA levels in the kidney and liver tissues of diabetic rats. The data are in agreement with previous report on increased oxidative stress markers in experimental diabetic animals.<sup>[8,43]</sup> In the diabetic rats that were treated with PLC, stimulation of the antioxidant defense system was notably observed as shown by the increase in the activities of SOD, GSH-Px, and CAT and decrease in the MDA level in the kidney and liver.

Recently, accumulating evidence has indicated the pro-inflammatory mediators such as TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 are also critically involved in the onset and progression of diabetes and its complication.<sup>[44,45]</sup> It is an undeniable fact that oxidative stress and inflammation have been proposed to work synergistically in diabetes. Hyperglycemia-induced oxidative stress can activate a number of stress-induced kinases as well as the formation of advanced glycation end products, which promotes increased expression of pro-inflammatory cytokines. The

**Table 3:** Effect of PLC on serum creatinine, urea, and blood urea nitrogen in diabetic rats


Groups	Creatinine (mg/L)	Urea (mg/L)	BUN (mg/L)
NCG	0.46±0.03	24.68±0.84	12.65±1.06
DCG	1.81±0.10 <sup>†</sup>	61.03±1.49 <sup>†</sup>	24.96±1.30 <sup>†</sup>
DLG	0.71±0.05**	33.01±0.99**	18.37±0.85**
PLC 100	0.87±0.02**	39.20±0.71**	18.80±0.80**
PLC 400	0.69±0.04**	29.71±0.88**	17.70±1.01**

Values are expressed as mean±SD (n=6). <sup>†</sup>P<0.05 versus normal control group; \*\*P<0.05 versus diabetic control group. BUN: Blood urea nitrogen; NCG: Normal control group; DCG: Diabetic control group; DLG: Diabetic rats treated with glibenclamide; SD: Standard deviation; PLC 100: Diabetic rats treated with *Lycium chinense* polysaccharide (100 mg/kg); PLC 400: Diabetic rats treated with *Lycium chinense* polysaccharide (400 mg/kg)

**Table 4:** Effect of PLC on oxidative stress markers in the kidney and liver of diabetic rats

Groups	GSH-Px (U/mg prot)	SOD (U/mg prot)	CAT (U/mg prot)	MDA (nmol/mg prot)
Kidney				
NCG	396.27±12.04	135.70±9.16	34.64±4.08	4.83±0.36
DCG	158.59±12.51 <sup>†</sup>	58.47±11.04 <sup>†</sup>	8.81±3.46 <sup>†</sup>	12.77±0.41 <sup>†</sup>
DLG	281.73±17.13**	83.61±12.20**	15.02±3.07**	8.37±0.24**
PLC 100	270.30±14.08**	90.22±10.07**	17.00±3.50**	8.92±0.17**
PLC 400	315.02±15.00**	117.60±12.88**	22.18±3.90**	6.70±0.20**
Liver				
NCG	582.90±16.81	192.41±8.04	62.18±4.78	5.77±0.51
DCG	304.21±18.51 <sup>†</sup>	67.05±7.39 <sup>†</sup>	20.40±2.50 <sup>†</sup>	17.19±1.04 <sup>†</sup>
DLG	408.73±14.77**	99.10±12.88**	38.01±3.03**	8.92±0.30**
PLC 100	394.05±22.36**	106.63±5.72**	36.91±3.40**	9.04±0.28**
PLC 400	507.78±25.40**	136.87±12.88**	46.09±2.17**	7.03±0.19**

Values are expressed as mean±SD (n=6). <sup>†</sup>P<0.05 versus normal control group; \*\*P<0.05 versus diabetic control group. GSH-Px: Glutathione Peroxidase; SOD: Superoxide Dismutase; CAT: Catalase; MDA: Malondialdehyde; NCG: Normal control group; DCG: Diabetic control group; DLG: Diabetic rats treated with glibenclamide; SD: Standard deviation; PLC 100: Diabetic rats treated with *Lycium chinense* polysaccharide (100 mg/kg); PLC 400: Diabetic rats treated with *Lycium chinense* polysaccharide (400 mg/kg)



**Figure 2:** Effect of PLC on proinflammatory cytokines; (a) TNF- $\alpha$ , (b) IL-6, and (c) IL-1  $\beta$  in the kidney tissues of STZ-induced diabetic rats. Values are expressed as mean  $\pm$  SD ( $n = 6$ ). \* $P < 0.05$  versus normal control group; \*\* $P < 0.05$  versus diabetic control group. NCG: Normal control group; DCG: Diabetic control group; DLG: Diabetic rats treated with glibenclamide; PLC 100: Diabetic rats treated with *Lycium chinense* polysaccharide (100 mg/kg); PLC 400: Diabetic rats treated with *Lycium chinense* polysaccharide (400 mg/kg); TNF- $\alpha$ : Tumor necrosis factor alpha; IL-1 $\beta$ : Interleukin 1 beta; IL-6: Interleukin 6; STZ: Streptozotocin; SD: Standard deviation

pro-inflammatory cytokines produced further instigate the generation of ROS production. In addition, inflammation and oxidative stress play a significant role in the onset and progression of insulin resistance as well as in the macro- and micro-vascular diabetic complications.<sup>[46-48]</sup> A number of recent studies have reported increased levels and expression of TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and nuclear factor-kappaB in the kidney of diabetic rats.<sup>[49-52]</sup> In this study, TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 were observed to be increased in the kidney of diabetic rats; interestingly, diabetic rats administered with PLC displayed significantly lowered levels of pro-inflammatory cytokines.

The antidiabetic effect of a number of purified and crude polysaccharide extracts from the genus *Lycium* has been reported, especially from *L. barbarum* and *Lycium ruthenicum*. Luo *et al.* reported the antidiabetic effect of a crude and purified polysaccharide in alloxan-induced diabetic rabbits; the crude polysaccharide extract showed excellent antioxidant, hypoglycemic, and hyperlipidemic effects in the treated animals.<sup>[53]</sup> Zhou *et al.* also reported that *L. barbarum* polysaccharide (LBP) ameliorated insulin resistance, increased GLUT-4 surface level, as well as improved intracellular insulin signaling in STZ-induced diabetic rats. Furthermore, Al-Fartosy reported that a polysaccharide (LBP3b) obtained from *L. barbarum* significantly reduced serum glucose level, ameliorated decrease in body weight and lipid metabolism, as well as prevented diabetic complications.<sup>[54,55]</sup> The crude *L. ruthenicum* polysaccharide extract was reported to have hypoglycemic properties in alloxan-induced diabetic mice by significantly reducing blood glucose level and enhancing serum and liver antioxidant enzyme status of diabetic mice.<sup>[56]</sup> The results obtained in our study is clearly in agreement and correlates well with these previous studies. PLC

effectively ameliorated hyperglycemia, hyperlipidemia, oxidative stress, and inflammation in the treated diabetic rats, in addition to improving insulin secretion.

Most of the polysaccharides isolated or characterized from the genus *Lycium* and several other plant species have been shown to comprise majorly of neutral and acidic sugars, such as arabinose, glucose, galactose, mannose, rhamnose, xylose, glucuronic acid, and galacturonic acid.<sup>[13,14,56-58]</sup> These components play a major role in the bioactivity of these polysaccharides. We envisaged that the antidiabetic properties displayed by PLC may be attributed to some of these constituents, owing to their role as bioactive agents in crude or purified polysaccharides in the previous report. Further studies on the identification and characterization of the substances in the PLC polysaccharide extract are ongoing.

## CONCLUSION

The results obtained from this study suggested that *L. chinense* polysaccharide attenuated hyperglycemia-induced oxidative stress, inflammation, and abnormalities in kidney and liver functions indexes in STZ-induced diabetic rats. These findings suggest that *L. chinense* polysaccharide can be useful as a functional food or as a nutraceutical for treating diabetes and diabetic complications.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Castro-Barquero S, Ruiz-León AM, Sadurní M, Estruch R, Casas R. Dietary strategies for metabolic syndrome: A review. *J Obest Ther* 2017;1:2.
- Kaur J. A comprehensive review on metabolic syndrome. *Cardiol Res Pract* 2014;2014:943162.
- Kaur G, Padiya R, Adela R, Putcha UK, Reddy GS, Reddy BR, *et al.* Garlic and resveratrol attenuate diabetic complications, loss of  $\beta$ -cells, pancreatic and hepatic oxidative stress in streptozotocin-induced diabetic rats. *Front Pharmacol* 2016;7:360.
- Khamchan A, Paseephol T, Hanchang W. Protective effect of wax apple (*Syzygium samarangense* (Blume) Merr. & L.M. Perry) against streptozotocin-induced pancreatic  $\beta$ -cell damage in diabetic rats. *Biomed Pharmacother* 2018;108:634-45.
- Liu M, Song X, Zhang J, Zhang C, Gao Z, Li S, *et al.* Protective effects on liver, kidney and pancreas of enzymatic-and acidic-hydrolysis of polysaccharides by spent mushroom compost (*Hypsizigum marmoreus*). *Sci Rep* 2017;7:43212.
- Fridlyand LE, Philipson LH. Oxidative reactive species in cell injury: Mechanisms in diabetes mellitus and therapeutic approaches. *Ann NY Acad Sci* 2005;1066:136-51.
- Zhang Q, Olatunji OJ, Chen H, Tola AJ, Oluwaniyi OO. Evaluation of the anti-diabetic activity of polysaccharide from *Cordyceps cicadae* in experimental diabetic rats. *Chem Biodivers* 2018;15:e1800219.
- Olatunji OJ, Chen H, Zhou Y. *Lycium chinense* leaves extract ameliorates diabetic nephropathy by suppressing hyperglycemia mediated renal oxidative stress and inflammation. *Biomed Pharmacother* 2018;102:1145-51.
- Xie JH, Jin ML, Morris GA, Zha XQ, Chen HQ, Yi Y, *et al.* Advances on bioactive polysaccharides from medicinal plants. *Crit Rev Food Sci Nutr* 2016;56 Suppl 1:S60-84.
- Yu Y, Shen M, Song Q, Xie J. Biological activities and pharmaceutical applications of polysaccharide from natural resources: A review. *Carbohydr Polym* 2018;183:91-101.
- Huang W, Deng H, Jin S, Ma X, Zha K, Xie M. The isolation, structural characterization and anti-osteosarcoma activity of a water soluble polysaccharide from *Agrimonia pilosa*. *Carbohydr Polym* 2018;187:19-25.
- Liu J, Willför S, Xu C. A review of bioactive plant polysaccharides: Biological activities, functionalization and biomedical applications. *Bioact Carbohydr Diet Fibre* 2018;5:31-61.
- Zheng Y, Bai L, Zhou Y, Tong R, Zeng M, Li X, *et al.* Polysaccharides from Chinese herbal medicine for anti-diabetes recent advances. *Int J Biol Macromol* 2019;121:1240-53.
- Masci A, Carradori S, Casadei MA, Paolicelli P, Petralito S, Ragno R, *et al.* *Lycium barbarum* polysaccharides: Extraction, purification, structural characterisation and evidence about hypoglycaemic and hypolipidaemic effects. A review. *Food Chem* 2018;254:377-89.
- Wu J, Shi S, Wang H, Wang S. Mechanisms underlying the effect of polysaccharides in the treatment of type 2 diabetes: A review. *Carbohydr Polym* 2016;144:474-94.
- Hu JL, Nie SP, Xie MY. Antidiabetic mechanism of dietary polysaccharides based on their gastrointestinal functions. *J Agric Food Chem* 2018;66:4781-6.
- Mocan A, Vlase L, Vodnar DC, Bischin C, Hanganu D, Gheldiu AM, *et al.* Polyphenolic content, antioxidant and antimicrobial activities of *Lycium barbarum* L. And *Lycium chinense* Mill. Leaves. *Molecules* 2014;19:10056-73.
- Yao R, Heinrich M, Weckerle CS. The genus *Lycium* as food and medicine: A botanical, ethnobotanical and historical review. *J Ethnopharmacol* 2018;212:50-66.
- Yao X, Peng Y, Xu LJ, Li L, Wu QL, Xiao PG. Phytochemical and biological studies of *Lycium* medicinal plants. *Chem Biodivers* 2011;8:976-1010.
- Potterat O. Goji (*Lycium barbarum* and *L. chinense*): Phytochemistry, pharmacology and safety in the perspective of traditional uses and recent popularity. *Planta Med* 2010;76:7-19.
- Ru Y, Chen X, Xu J, Huang L, Jiang M, Guo L, *et al.* Hypoglycemic effects of a polysaccharide from *Tetragium hemsleyanum* Diels & Gilg in alloxan-induced diabetic mice. *Chem Biodivers* 2018;15:e1800070.
- Li X, Zhao R, Zhou HL, Wu DH. Deproteinization of polysaccharide from the *Stigma maydis* by Sevag method. *Adv Mater Res* 2011;340:416-20.
- Arulselvan P, Fard MT, Tan WS, Gothai S, Fakurazi S, Norhaizan ME, *et al.* Role of antioxidants and natural products in inflammation. *Oxid Med Cell Longev* 2016;2016:5276130.
- Lenzen S. The mechanisms of alloxan-and streptozotocin-induced diabetes. *Diabetologia* 2008;51:216-26.
- Maritim AC, Sanders RA, Watkins JB 3<sup>rd</sup>. Diabetes, oxidative stress, and antioxidants: A review. *J Biochem Mol Toxicol* 2003;17:24-38.
- Cade WT. Diabetes-related microvascular and macrovascular diseases in the physical therapy setting. *Phys Ther* 2008;88:1322-35.
- Nordwall M, Arngqvist HJ, Bojestig M, Ludvigsson J. Good glycemic control remains crucial in prevention of late diabetic complications – The Linköping diabetes complications study. *Pediatr Diabetes* 2009;10:168-76.
- Anyanwu GO, Iqbal J, Khan SU, Zaib S, Rauf K, Onyeneke CE, *et al.* Antidiabetic activities of chloroform fraction of *Anthocleista vogelii* planch root bark in rats with diet-and alloxan-induced obesity-diabetes. *J Ethnopharmacol* 2019;229:293-302.
- Owolabi MA, Jaja SI, Olatunji OJ, Oyekanmi OO, Adepoju S. Attenuation of oxidative damage in alloxan induced diabetic rabbits following administration of the extract of the leaves of *Vernonia amygdalina*. *Free Radic Antioxid* 2011;1:94-101.
- Zhang L, Liu Y, Ke Y, Liu Y, Luo X, Li C, *et al.* Antidiabetic activity of polysaccharides from *Suililellus luridus* in streptozotocin-induced diabetic mice. *Int J Biol Macromol* 2018;119:134-40.
- Ktari N, Mnafigui K, Nasri R, Hamden K, Bkhairia I, Ben Hadj A, *et al.* Hypoglycemic and hypolipidemic effects of protein hydrolysates from zebra blenny (*Salaria basilisca*) in alloxan-induced diabetic rats. *Food Funct* 2013;4:1691-9.
- Chait A, Bornfeldt KE. Diabetes and atherosclerosis: Is there a role for hyperglycemia? *J Lipid Res* 2009;50 Suppl:S335-9.
- Chen X, Bai X, Liu Y, Tian L, Zhou J, Zhou Q, *et al.* Anti-diabetic effects of water extract and crude polysaccharides from tuberous root of *Liriope spicata* var. *Prolifera* in mice. *J Ethnopharmacol* 2009;122:205-9.
- Ahmad W, Khan I, Khan MA, Ahmad M, Subhan F, Karim N. Evaluation of antidiabetic and antihyperlipidemic activity of *Artemisia indica* linn (aerial parts) in streptozotocin induced diabetic rats. *J Ethnopharmacol* 2014;151:618-23.
- Baek HJ, Jeong YJ, Kwon JE, Ra JS, Lee SR, Kang SC. Antihyperglycemic and antilipidemic effects of the ethanol extract mixture of *Ligularia fischeri* and *Momordica charantia* in type II diabetes-mimicking mice. *Evid Based Complement Alternat Med* 2018;2018:3468040.
- Gao H, Wen JJ, Hu JL, Nie QX, Chen HH, Xiong T, *et al.* Polysaccharide from fermented *Momordica charantia* L. With *Lactobacillus plantarum* NCU116 ameliorates type 2 diabetes in rats. *Carbohydr Polym* 2018;201:624-33.
- Pan Y, Wang C, Chen Z, Li W, Yuan G, Chen H, *et al.* Physicochemical properties and antidiabetic effects of a polysaccharide from corn silk in high-fat diet and streptozotocin-induced diabetic mice. *Carbohydr Polym* 2017;164:370-8.
- Matsunami T, Sato Y, Hasegawa Y, Ariga S, Kashimura H, Sato T, *et al.* Enhancement of reactive oxygen species and induction of apoptosis in streptozotocin-induced diabetic rats under hyperbaric oxygen exposure. *Int J Clin Exp Pathol* 2011;4:255-66.
- Phull AR, Majid M, Haq IU, Khan MR, Kim SJ. *In vitro* and *in vivo* evaluation of anti-arthritis, antioxidant efficacy of fucoidan from *Undaria pinnatifida* (Harvey) Suringar. *Int J Biol Macromol* 2017;97:468-80.
- Zhang J, Fan S, Mao Y, Ji Y, Jin L, Lu J, *et al.* Cardiovascular protective effect of polysaccharide from *Ophiopogon japonicus* in diabetic rats. *Int J Biol Macromol* 2016;82:505-13.
- Abuja PM, Albertini R. Methods for monitoring oxidative stress, lipid peroxidation and oxidation resistance of lipoproteins. *Clin Chim Acta* 2001;306:1-7.
- Guo L, Ma R, Sun H, Raza A, Tang J, Li Z. Anti-inflammatory activities and related mechanism of polysaccharides isolated from *Sargentodoxa cuneata*. *Chem Biodivers* 2018;15:e1800343.
- Tang X, Olatunji OJ, Zhou Y, Hou X. *Allium tuberosum*: Antidiabetic and hepatoprotective activities. *Food Res Int* 2017;102:681-9.
- Gothai S, Ganesan P, Park SY, Fakurazi S, Choi DK, Arulselvan P. Natural phyto-bioactive compounds for the treatment of type 2 diabetes: Inflammation as a target. *Nutrients* 2016;8: pii: E461.
- Spranger J, Kroke A, Möhlig M, Hoffmann K, Bergmann MM, Ristow M, *et al.* Inflammatory cytokines and the risk to develop type 2 diabetes: Results of the prospective population-based European prospective investigation into cancer and nutrition (EPIC)-Potsdam study. *Diabetes* 2003;52:812-7.
- Vlassara H, Uribarri J. Advanced glycation end products (AGE) and diabetes: Cause, effect, or both? *Curr Diab Rep* 2014;14:453.
- Jung UJ, Choi MS. Obesity and its metabolic complications: The role of adipokines and the relationship between obesity, inflammation, insulin resistance, dyslipidemia and nonalcoholic fatty liver disease. *Int J Mol Sci* 2014;15:6184-223.
- Chen L, Chen R, Wang H, Liang F. Mechanisms linking inflammation to insulin resistance. *Int J Endocrinol* 2015;2015:508409.
- Baker RG, Hayden MS, Ghosh S. NF- $\kappa$ B, inflammation, and metabolic disease. *Cell Metab* 2011;13:11-22.
- Tavakoly R, Maracy MR, Karimifar M, Entezari MH. Does fenugreek (*Trigonella foenum-graecum*) seed improve inflammation and oxidative stress in patients with type 2

- diabetes mellitus? A parallel group randomized clinical trial. *Eur J Integr Med* 2018;18:13-7.
51. Shoelson SE, Lee J, Goldfine AB. Inflammation and insulin resistance. *J Clin Invest* 2006;116:1793-801.
52. Al Hroob AM, Abukhalil MH, Alghonmeen RD, Mahmoud AM. Ginger alleviates hyperglycemia-induced oxidative stress, inflammation and apoptosis and protects rats against diabetic nephropathy. *Biomed Pharmacother* 2018;106:381-9.
53. Luo Q, Cai Y, Yan J, Sun M, Corke H. Hypoglycemic and hypolipidemic effects and antioxidant activity of fruit extracts from *Lycium barbarum*. *Life Sci* 2004;76:137-49.
54. Zhao R, Li Q, Xiao B. Effect of *Lycium barbarum* polysaccharide on the improvement of insulin resistance in NIDDM rats. *Yakugaku Zasshi* 2005;125:981-8.
55. Al-Fartosy AJ. Protective effect of galactomannan extracted from Iraqi *Lycium barbarum* L. fruits against alloxan-induced diabetes in rats. *Am J Biochem Biotechnol* 2015;11:73-83.
56. Wang JH, Chen XQ, Zhang WJ. Study on hypoglycemic function of polysaccharides from *Lycium ruthenicum* Murr. Fruit and its mechanism. *Chin J Food Sci* 2009;30:244-8.
57. Peng Q, Lv X, Xu Q, Li Y, Huang L, Du Y. Isolation and structural characterization of the polysaccharide LRGP1 from *Lycium ruthenicum*. *Carbohydr Polym* 2012;90:95-101.
58. Qian D, Zhao Y, Yang G, Huang L. Systematic review of chemical constituents in the genus *Lycium* (*Solanaceae*). *Molecules* 2017;22. pii: E911.