

# Anticancer Effects of Piperine-Free *Piper nigrum* Extract on Cholangiocarcinoma Cell Lines

Aman Tedasen<sup>1,2</sup>, Araya Khoka<sup>3</sup>, Siribhorn Madla<sup>1</sup>, Somchai Sriwiriyan<sup>3</sup>, Potchanapond Graidist<sup>1,2</sup>

<sup>1</sup>Department of Biomedical Sciences, Faculty of Medicine, Prince of Songkla University, <sup>2</sup>The Excellent Research Laboratory of Cancer Molecular Biology, Prince of Songkla University, <sup>3</sup>Department of Pharmacology, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla, Thailand

Submitted: 01-08-2019

Revised: 30-09-2019

Published: 31-03-2020

## ABSTRACT


**Background:** Black pepper (*Piper nigrum* L.) is widely used as a traditional medicine, including usage for pain relief, fevers, as well as an anticancer agent. Previously, we reported that piperine-free *P. nigrum* extract (PFPE) inhibited breast cancer *in vitro* and *in vivo*.

**Objective:** In this present study, we explored the anticancer effects of PFPE on cholangiocarcinoma (CCA). **Materials and Methods:** 3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyltetrazolium bromide assay was performed to analyze cytotoxic potential of PFPE whereas deoxyribonucleic acid (DNA) fragmentation followed by Western blot analysis were used. **Results:** PFPE composed of alkaloid, flavonoid, amide, lignans, opioid, and steroid. This crude extract represented cytotoxic effect against CCA cells which stronger than dichloromethane *P. nigrum* crude extract and piperine, especially on KKU-M213 (median inhibition concentration [IC<sub>50</sub>] at 13.70 µg/ml) and TFK-1 (IC<sub>50</sub> at 15.30 µg/ml). Interestingly, PFPE showed lower cytotoxicity against normal human cholangiocyte MMNK-1 cells (IC<sub>50</sub> at 19.65 µg/ml) than KKU-M213 and TFK-1 cells. Then, the molecular mechanisms of PFPE were firstly evaluated by DNA fragmentation followed by Western blot analysis. The degradation of DNA was observed on KKU-M213 and TFK-1 cells after treatment with PFPE at day 2. Then, proliferation proteins including topoisomerase II, AKT8 virus oncogene cellular homolog, avian myelocytomatosis virus oncogene cellular homolog, cyclin D1, signal transducer and activator of transcription 3, cyclooxygenase-2, and nuclear factor kappa-light-chain-enhancer of activated B cells (NF-κB) were decreased and p21 was increased. Furthermore, apoptotic proteins, such as tumor protein p53, Bcl-2-associated X protein, and p53 upregulated modulator of apoptosis were upregulated. Meanwhile, antiapoptotic protein B-cell lymphoma 2 was down-regulated. **Conclusion:** These results indicated that PFPE inhibited CCA through the down-regulation of cell proliferation and induction of apoptosis pathway.

**Key words:** Anticancer, apoptosis, cell proliferation, cholangiocarcinoma, *Piper nigrum*

## SUMMARY

- piperine free *Piper nigrum* extract (PFPE) inhibited cholangiocarcinoma (CCA) cell lines
- PFPE induces CCA cells to undergo apoptosis and cell cycle arrest via the inhibition of topoisomerase II
- PFPE inhibit cell growth through the inhibition of nuclear factor kappa-light-chain-enhancer of activated B cells.



**Abbreviations used:** PFPE: Piperine free *Piper nigrum* extract; CCA: Cholangiocarcinoma; DPCE: dichloromethane *P. nigrum* crude extract; NMU: N-nitrosomethylurea; ER: Estrogen receptor; MMP-9: Matrix metalloproteinase-9; MMP-2: Matrix metalloproteinase-2; VEGF: Vascular endothelial growth factor; GC-MS: Gas chromatograph-mass spectrometer; MTT: 3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyltetrazolium bromide; DMSO: Dimethylsulfoxide; IC<sub>50</sub>: Median inhibition concentration; MCLE: Methanol crude extract of *Curcuma longa*; DNA: Deoxyribonucleic acid; STAT3: Signal transducer and activator of transcription 3; COX-2: Cyclooxygenase-2; NF-κB: Nuclear factor kappa-light-chain-enhancer of activated B cells; c-Myc: Avian myelocytomatosis virus oncogene cellular homolog; Akt: AKT8 virus oncogene cellular homolog; Bcl-2: B-cell lymphoma 2; p53: Tumor protein p53; Bax: Bcl-2-associated X protein; PUMA: p53 upregulated modulator of apoptosis.

## Correspondence:

Dr. Potchanapond Graidist,  
Department of Biomedical Sciences, Faculty of  
Medicine, Prince of Songkla University, Hat Yai,  
Songkhla, Thailand.  
E-mail: gpotchan@medicine.psu.ac.th  
DOI: 10.4103/pm.pm\_288\_19

## Access this article online

Website: www.phcog.com

## Quick Response Code:



## INTRODUCTION

Cholangiocarcinoma (CCA) is an epithelial cancer originating from the bile ducts with features of cholangiocyte differentiation.<sup>[1]</sup> There are 2 types of CCA (based on its location) including intrahepatic and extrahepatic.<sup>[2]</sup> For over the past four decades, incidence of CCA has been increased in United States of America,<sup>[3]</sup> Australia, England,<sup>[4]</sup> and Northeastern Thailand.<sup>[5]</sup> There are several risk factors for CCA, including primary sclerosing cholangitis, liver fluke infections (*Clonorchis sinensis* and *Opisthorchis viverrini*), choledochal cysts, Caroli's disease, hepatitis B and C infection, obesity, cirrhosis and hepatolithiasis.<sup>[5,6]</sup> The therapeutic for CCA are limited and no

current effective treatment because the majority of patients present with advanced stage disease.<sup>[7]</sup> Even treatments with advances in surgical

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

**Cite this article as:** Tedasen A, Khoka A, Madla S, Sriwiriyan S, Graidist P. Anticancer effects of piperine-free *Piper nigrum* extract on cholangiocarcinoma cell lines. Phcog Mag 2020;16:S28-38.

techniques, chemotherapy and radiotherapy, the 5-year survival rate of patients after diagnosis still remain about 10%.<sup>[8]</sup> Although surgical resection has improved in the survival of most patients, the recurrent disease was found within 2 years after tumor resection.<sup>[9]</sup> Chemotherapy and radiation therapy are ineffective and show various side effects such as harmful to normal cells and bone marrow suppression.<sup>[10]</sup> Therefore, effective therapeutic and alternative treatments with no serious side effect for CCA are urgently needed.

*P. nigrum* L. belongs to family Piperaceae and can be used as antiapoptotic, antibacterial, anticolon toxin, antidepressant, antifungal, anti-diarrhoeal, anti-inflammatory, antimutagenic, antimetastatic, antioxidative, antipyretic, antispasmodic, antispermato-genic, antitumor, antithyroid, ciprofloxacin potentiator, cold extremities, gastric ailments, hepatoprotective, insecticidal, intermittent fever, and larvicidal activities.<sup>[11]</sup> The chemical constituents of *P. nigrum* are aromatic essential oils, alkaloids, amides, propenylphenols, lignans, terpenes, flavones, and steroids.<sup>[12]</sup> Ethanolic crude extract of *P. nigrum* consists of high total phenol content shows antioxidant and anti-inflammation as well as cytotoxic property against colorectal carcinoma cell lines.<sup>[13]</sup> Using ethanol and high pressure (200 bar), *P. nigrum* crude extracts exhibits cytotoxicity against MCF-7 with median inhibition concentration (IC<sub>50</sub>) of 14.40 ± 3.30 µg/ml and represents tumor inhibitory effect in mammary adenocarcinoma mouse.<sup>[14]</sup> Previously, we reported that piperine-free *P. nigrum* extract (PFPE) strongly inhibited breast cancer MCF-7 cells with IC<sub>50</sub> value of 7.45 µg/ml. Moreover, PFPE inhibited tumor growth in *N*-nitrosomethylurea-induced mammary tumorigenesis rats without liver and kidney toxicity.<sup>[15]</sup> Interestingly, PFPE upregulated tumor protein p53 (p53) and downregulated estrogen receptor, E-cadherin, matrix metalloproteinase-9 (MMP-9), MMP-2, avian myelocytomatosis virus oncogene cellular homolog (c-Myc) and vascular endothelial growth factor (VEGF) *in vitro* and *in vivo*.<sup>[16]</sup> In this present research, we further explored the phytochemical component, investigated cytotoxicity and molecular mechanisms of PFPE on CCA cell lines.

## MATERIALS AND METHODS

### Preparation of piperine free *Piper nigrum* extract

Seeds of *P. nigrum* L. were collected from Songkhla province in Thailand. The plant specimen (voucher specimen number SKP 146161401) was identified by Asst. Prof. Dr. Supreeya Yuenyongsawad and deposited in the herbarium at the Southern Centre of Thai Traditional Medicine, Department of Pharmacognosy and Pharmaceutical Botany, Prince of Songkla University, Thailand. PFPE was prepared as previously described. Briefly, grounded 250 g of dried seeds of *P. nigrum* L. were soaked in 300 mL of dichloromethane and incubated at 35°C for 3 h in a shaking incubator. After filtration with Whatman filter paper No. 1 and concentration using rotary evaporator, the dark brown oil residue of extracts was obtained and then recrystallized with cold diethyl ether in an ice bath to get rich of yellow crystals (piperine) and obtain brown oil residue (PFPE).<sup>[15]</sup> PFPE was kept in a desiccator until used.

### Phytochemical analysis and identification of bioactive constituents by gas chromatograph-mass spectrometer

The analysis of the phytochemical screening and composition of PFPE extracts were carried out using a Gas Chromatography-Agilent 7890B combination with an Agilent 5977A triple quadrupole mass spectrometer (Agilent Technologies Inc, USA). Gas chromatograph-mass spectrometer (GC-MS) analysis is a common confirmation test, which used to make an effective chemical analysis. The PFPE samples were evaluated phytochemicals such as a flavonoids, tannins, alkaloids,

steroids, phenols, glycosides, lignans, and terpenoids. An inlet temperature of 280°C with the split ratio 7:1 was employed and the helium was used as the carried gas at the constant flow rate of 7 ml/min. The oven temperature was initially maintained at 60°C for 5 min and increase at a rate of 5°C/min to 315°C for 15 min. For MS detection, an electron ionization mode was used with an ionization energy of 70 eV, ion source temperature of 230°C, and scan mass range *m/z* 35–500. The components were identified based on a correlation of the recorded fragmentation patterns of mass spectra that provided in the GC-MS system software version Wiley10 and NIST14. All procedures were performed at Scientific Equipment Center, Prince of Songkla University, Songkhla, Thailand.

### Measuring total phenolic, tannin, flavonoid content and radical scavenging activity

The total phenolic content was determined based on Folin–ciocalteu method. Gallic acid was used as the standard and total phenolics were expressed as mg gallic acid equivalent/mg extract (mg GAE/mg extract). Total condensed tannin was measured based on HCL-vanillin method and catechin was used as the standard. The total tannin was reported as mg catechin equivalent/mg extract (mg CE/mg extract). The total flavonoid content was determined by aluminum chloride solution (AlCl<sub>3</sub>) colorimetric method. Quercetin was employed as the standard and expressed the total flavonoids as mg quercetin equivalent/mg extract (mg QE/mg extract). 2,2-diphenyl-1-picryl-hydrazyl-hydrate (DPPH) radical scavenging activity was performed according to the DPPH trolox assay and reported as mg trolox equivalent antioxidant capacity/mg extract (mg TEAC/mg extract). All procedures were performed at Center of Excellence in Natural Products Innovation, Mae Fah Luang University, Chiang Rai, Thailand.

### Cell lines and culture conditions

Three CCA (KKU-100, KKU-M213 and KKU-M055) and one cholangiocyte (MMNK-1) cells were kindly donated by Dr. Mutita Junking (Faculty of Medicine, Mahidol University, Bangkok, Thailand). TFK-1 cells were obtained from RIKEN BioResource Center and HuCC-T1 cells were obtained from the Japanese Collection of Research Bioresources Cell Bank. Mouse fibroblast, L-929 cells, were kindly donated by Associate Professor Dr. Jasadee Kaewsichan (Department of Pharmaceutical Chemistry, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Songkhla, Thailand).

KKU-100, KKU-M213, KKU-M055, MMNK-1 and L-929 cells were grown in DMEM medium (Invitrogen), which contained 10% of fetal bovine serum (Invitrogen), 2 mmol/L of L-glutamine (Invitrogen), and an antibiotic mixture of 100 units/mL of penicillin and 100 µg/mL of streptomycin (Invitrogen). TFK-1 and HuCC-T1 cells were grown in RPMI 1640 (Invitrogen) supplemented with the same supplement as for DMEM. All cells were maintained by incubating in a 5% CO<sub>2</sub> atmosphere, at 37°C and 96% relative humidity.

### *In vitro* cytotoxicity

The cytotoxicity assay was performed in 96-well plate. KKU-100, KKU-M055, and MMNK-1 cells were seeded at a density of 5 × 10<sup>3</sup> cells/well. KKU-M213, TFK-1, and HuCC-T1 cells were seeded at a density of 7.5 × 10<sup>3</sup> cells/well and L-929 cells were seeded at a density of 8 × 10<sup>3</sup> cells/well. After incubation for 24 h, cells were treated with PFPE at various concentration for 48 h. The cells were then washed with 1X PBS and incubated in 100 µl of 0.5 mg/ml of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) solution at 37°C for 30 min. Under light protection, the purple crystals of formazan or MTT metabolites were dissolved with 100 µl of dimethyl

sulfoxide and incubate at 37°C for 30 min. The absorbance was measured at 570 and 650 nm using a microplate reader spectrophotometer (Spectra Max M5, Molecular Devices), and the IC<sub>50</sub> values were calculated.<sup>[17]</sup> According to USNCI plant screening program, a crude extract is generally considered to have *in vitro* cytotoxic activity with IC<sub>50</sub> value ≤20 µg/ml.<sup>[18]</sup>

## Deoxyribonucleic acid fragmentation analysis

KKU-M213 and TFK-1 cells in their exponential growth phase were seeded into 6 cm culture plate at a density of  $2.5 \times 10^5$  cells/plate for 24 h and then treated with PFPE at 3 folds of IC<sub>50</sub> values. After treatment for 96 h, cells were harvested by trypsinization. Cell pellets were lysed using the extraction buffer (containing 0.7 M NaCl, 17 mM SDS, 10 mM Tris-HCl (pH 8.0) and 2 mM EDTA (pH 8.0)) and fragmented deoxyribonucleic acid (DNA) in the supernatant was extracted once with an equal volume of phenol: chloroform: isoamyl alcohol (25:24:1) and once with chloroform: isoamyl alcohol (24:1). The DNA was precipitated with a two-thirds volume of cold isopropanol followed by centrifugation at 8,000 ×g and washed once in 70% ethanol. Finally, DNA pellet was resuspended in deionized water and analyzed by 1.5% agarose gel electrophoresis.<sup>[19]</sup>

## Western blot analysis

KKU-M213 and TFK-1 cells were seeded into 6 cm culture plate at a density of  $2.5 \times 10^5$  cells/plate for 24 h and then treated with PFPE at IC<sub>50</sub> values. After treatment, cells were harvested every day for 4 days. Then, cell pellets were lysed using the RIPA buffer (containing 150 mM NaCl, 50 mM Tris, pH 7.4, 1% (v/v) NP-40, 0.25% (w/v) sodium deoxycholate and 1 mM EDTA). Total protein samples (150 mg) were loaded on 12% of SDS-polyacrylamide gel electrophoresis and transferred onto a 0.45 mm nitrocellulose membrane (Bio-Rad, 162-0115). Membrane was blocked at room temperature for 1 h with 5% non-fat milk in 1X TBS-T and then washed with 1% non-fat milk in 1X TBS-T. Membrane was incubated with primary antibodies against topoisomerase II, Bcl-2-associated X protein (Bax), B-cell lymphoma 2 (Bcl-2), p53 upregulated modulator of apoptosis (PUMA), p21, AKT8 virus oncogene cellular homolog (Akt), cyclooxygenase-2 (COX-2), Nuclear factor kappa-light-chain-enhancer of activated B cells (NF-κB), signal transducer and activator of transcription 3 (STAT-3), cyclin D1 and p53 proteins. The membrane was then incubated with secondary horseradish peroxidase-conjugated antibodies. Bound antibodies were developed by a chemiluminescence detection kit using the SuperSignal™ West Dura Extended Duration Substrate (Thermo Scientific) and detected using a Fusion FX vilber lourmat, CCD camera (Fisher Biotechnology). GAPDH was used to normalize protein loading. Protein levels were expressed as a relative ratio to GPADH.

## Statistical analysis

The median inhibition concentration (IC<sub>50</sub>) data was acquired by SoftMax 1 Pro 5 program (MDS Analytical Technologies Inc., California, USA). Student's *t*-test was used to analyze intergroup differences. A *P* < 0.05 was considered to be statistically significant. All results were represented as the mean ± standard deviation (SD). The values were obtained from at least three independent experiments.

# RESULTS

## Total phenolic, tannin, and flavonoid contents

Phenolics, flavonoids, and tannins are one class of secondary plant metabolites which represented anticancer activity of plant. As present in Table 1, PFPE contained phenolic, tannin and flavonoid lower than methanol crude extract of *Curcuma longa* (MCLE). However, the cytotoxicity of PFPE against breast cancer MCF-7 cells (IC<sub>50</sub> value

**Table 1:** Total phenolic, tannin and flavonoid contents in piperine free *Piper nigrum* crude extract

Crude	Phenolics (mg GAE/g extract) <sup>a</sup>	Flavonoids (mg QE/mg extract) <sup>b</sup>	Tannins (mg CE/mg extract) <sup>c</sup>
PFPE	402.46±7.49	40.69±5.99	201.82±17.78
MCLE	2090.63±15.81	148.94±33.64	2373.75±92.77

<sup>a</sup>Mg of gallic acid equivalence by mg of extract; <sup>b</sup>Mg of quercetin equivalence by mg of extract; <sup>c</sup>Mg of catechin equivalence by mg of extract; *P. nigrum*: *Piper nigrum*; PFPE: Piperine free *P. nigrum* extract; *C. longa*: *Curcuma longa*; MCLE: Metanolic *C. longa* extract; GAE: Gallic acid equivalent; QE: Quercetin equivalent; CE: Catechin equivalent

at  $7.45 \pm 0.6$  µg/ml) not significantly lower than MCLE (IC<sub>50</sub> value at  $5.74 \pm 1.48$  µg/ml). Therefore, we performed GC-MS in next experiment to identify the chemical compounds in PFPE.

## Phytochemical screening

In this study, the phytochemical analysis using GC-MS was carried out. The chromatogram and predicted constituents are shown in Figure 1 and Table 2. Results showed that PFPE contained five chemical groups including alkaloids, terpenes, amides, lignans, opioid and steroid with 17, 13, 7, 3, 1, and 1 compounds, respectively. The highest percentage of peak area of each group were piperidine (21.66%, alkaloid), caryophyllene (13.28%, terpene), acrivastine (2.34%, amide), kusunokinin (1.28%, lignan), methyldihydromorphine (1.18%, opioid), and beta-stigmasterol (1.74%, steroid) which showed the anticancer activity.

## Effect of piperine free *Piper nigrum* extract on the viability of cholangiocarcinoma, cholangiocyte and normal fibroblast cell lines

The cell viability of CCA and normal cell lines was measured using the MTT assay. All cell lines were incubated with extracts for 48 h. The IC<sub>50</sub> values represented the mean ± SD of three different experiments. Among these cell lines, PFPE showed the highest cytotoxicity against KKU-M213 cells with IC<sub>50</sub> value of  $13.70 \pm 1.14$  µg/ml. Moreover, PFPE demonstrated cytotoxic effect stronger than dichloromethane *P. nigrum* crude extract (DPCE) (IC<sub>50</sub> at  $22.22 \pm 0.26$  µg/ml) and piperine (IC<sub>50</sub> at  $27.01 \pm 0.36$  µg/ml). The positive reference drug (doxorubicin) showed a very strong cytotoxic activity on normal and almost cancer cells. Surprisingly, doxorubicin showed same cytotoxic activity with PFPE against TFK-1 cells [Table 3].


## Piperine free *Piper nigrum* extract induces deoxyribonucleic acid fragmentation on KKU-M213 and TFK-1 cells

A DNA fragmentation assay was used to determine whether the action of PFPE was associated with apoptosis or not. Apoptosis can be visualized as a ladder pattern of 180-200 base pairs due to DNA cleavage by the activation of a nuclear endonuclease enzyme. Since, PFPE demonstrated a strong cytotoxic effective on KKU-M213 and TFK-1 cells, both cell lines were used to determined DNA fragmentation. As shown in Figure 2, the DNA ladder pattern was observed at day 2 after exposure with 3 folds of IC<sub>50</sub> concentration of PFPE.

## Piperine free *Piper nigrum* extract inhibited proteins associated with inflammation that induces bile duct cancer

In this experiment, we determined proteins associated with inflammation that induced bile duct cancer including STAT-3, COX-2 and NF-κB using Western blot analysis. KKU-M213 cells were treated with 13.69 µg/ml of





**Table 2:** Chemical constituents in piperine free *Piper nigrum* extract

Identified compounds	Formula	Nature of compound	Molecular massb (g/mol)	Retention time	Area (%)	Biological activity
3-Carene	$C_{10}H_{16}$	Terpenes	136.24	9.0896	0.28	Antioxidant, antihyperuricemic and anti-inflammatory <sup>[33]</sup>
D-Limonene	$C_{10}H_{16}$	Terpenes	136.24	9.7228	0.39	Enhanced the antitumor effect of docetaxel against prostate cancer cells <sup>[34]</sup>
Clohexane, 4-ethenyl-4-methyl-3-(1-methylethenyl)-1-(1-methylethyl)-, (3R-trans) 2,4-diisopropenyl-1-methyl-1-vinylcyclohexane (or beta-Elementene)	$C_{15}H_{24}$	Terpenes	204.36	19.2545	2.20	Cytotoxic effect on K562 (leukemic) cells by the induction of apoptosis <sup>[35]</sup>
Copaene	$C_{15}H_{24}$	Terpenes	204.36	20.2929	1.26	Antimicrobial activity against an anaerobic microorganism <i>Prevotella nigrescens</i> <sup>[36]</sup>
2,4-diisopropenyl-1-methyl-1-vinylcyclohexane (beta-Elementene)	$C_{15}H_{24}$	Terpenes	204.36	20.7150	0.73	Cytotoxic effect on K562 (leukemic) cells by the induction of apoptosis <sup>[35]</sup>
Caryophyllene	$C_{15}H_{24}$	Terpenes	204.36	21.4893	13.28	Antioxidant, preventing lipidic oxidative damage and prevention of atherosclerosis <sup>[37]</sup> , antigenotoxic and santioxidant <sup>[38]</sup>
1,4,7,7-Cycloundecatriene, 1,5,9,9-tetra methyl-, Z, Z, Z-Naphthalene, decahydro-4a-methyl-1-methylene-7-(1-methylethenyl)-, [4aR-(4a.alpha.,7.alpha.,8a.beta.)]- (or beta-helmscapene, beta-Selinene)	$C_{15}H_{24}$ $C_{15}H_{24}$	Terpenes Terpenes	204.36 204.35	22.3144 23.1348	1.15 0.60	No activity reported Antioxidant and cytotoxic activity against HT29 (colon cancer) cells <sup>[39]</sup> , cytotoxicity against KB (oral cancer), MCF-7 (breast cancer) and NCI-H187 (small cell lung cancer) cells <sup>[40]</sup>
2-Isopropenyl-4a, 8-dimethyl-1,2,3,4,4a, 5,6,8a-octahydronaphthalene (or 7-Epi-alpha-Selinene) delta-Cadinene	$C_{15}H_{24}$	Terpenes	204.36	23.3522	0.54	Antimicrobial activity against <i>Bacillus subtilis</i> and <i>Candida albicans</i> <sup>[41]</sup>
Caryophyllene oxide	$C_{15}H_{24}O$	Terpenes	220.36	24.0207	0.61	Induction of apoptosis and cell cycle arrest on OVACR-3 (ovarian cancer) cells <sup>[42]</sup>
Isospathulenol	$C_{15}H_{24}O$	Terpenes	220.37	25.4618	0.42	Chemosensitizing agents for doxorubicin chemotherapy <sup>[43]</sup>
2,4-Decadienamide, N-isobutyl-, (E, E)- (or Pellitorine)	$C_{14}H_{25}NO$	Amides	223.36	26.4932	0.71	anticancer <sup>[44]</sup> increased the efficacy of DOX in MDA-MB-231 (breast cancer) cells <sup>[45]</sup> , inhibit STAT3 signaling pathway <sup>[46]</sup>
Piperidine, 1-(1-oxo-3-phenyl-2-prope nyl)- (or piperidine, 1-Cinnamoylpiperidine)	$C_{14}H_{17}NO$	Alkaloids	215.29	32.8537	2.28	Cytotoxic effects against <i>Aspergillus niger</i> , <i>Artemia salina</i> and <i>Caenorhabditis elegans</i> <sup>[47]</sup>
(2E,4E)-1-(Pyrrolidin-1-yl) deca-2,4-dien-1-one (or Iyeramide A, sarmentine)	$C_{14}H_{23}NO$	Alkaloids	221.34	36.1008	0.22	Antibacterial, anticancer and anti-inflammatory <sup>[48]</sup>
(2E,4E)-N-Isobutyldeca-2,4-dienamide (or Dodecatetraenoic acid isobutylamide)	$C_{16}H_{29}NO$	Amides	251.41	36.2247	0.37	No activity reported
N-Benzylidene-4-fluoroaniline	$C_{13}H_{10}FN$	Alkaloids	199.23	36.7524	0.48	Cytotoxicity against CCRF-CEM (acute lymphoblastic leukemia), HL-60 (acute promyelocytic leukemia), PC-3 (prostate carcinoma), and HA22T (hepatoma) cells <sup>[27]</sup> inhibit lipoxigenase (5-LOX) and cyclooxygenase-1 (COX-1) <sup>[49]</sup>
(E)-5-(Benzol[d][1,3]dioxol-5-yl)-1-(pi peridin-1-yl) pent-2-en-1-one (or piperanine)	$C_{17}H_{21}NO_3$	Alkaloids	287.359	44.1035	0.34	Inhibit allergic and inflammatory <sup>[50]</sup>
Piperlonguminine	$C_{16}H_{19}NO_3$	Alkaloids	273.33	44.5123	0.88	No activity reported
(E)-1-(Piperidin-1-yl) hexadec-2-en-1-one	$C_{22}H_{39}NO$	Alkaloids	321.54	44.8101	4.77	Hepatoprotective effect <sup>[51]</sup>
Piperine	$C_{17}H_{19}NO_3$	Alkaloids	285.34	45.3603	0.79	Anticancer against breast cancer cells <sup>[31]</sup>
(2E,4E,10E)-N-Isobutylhexadeca-2,4,10-trienamide	$C_{26}H_{43}NO$	Amides	305.50	46.3182	5.09	No activity reported
(2E,4E)-N-Isobutyltodeca-2,4-dienamide (or Pipericine)	$C_{22}H_{35}NO$	Amides	335.58	46.5162	0.48	Anticancer against Hep-G2 (hepatocellular carcinoma) <sup>[52]</sup> and Hela (cervical cancer) cells <sup>[53]</sup>
1-Benzyl-2-(1-ethoxycarbonyl-2-phenylethyl)-4,5-dihydroimidazole (Acrivastine)	$C_{22}H_{24}N_2O_2$	Amides	348.45	46.6004	0.85	No activity reported
(E)-7-(Benzol[d][1,3]dioxol-5-yl)-1-(pyrrolidin-1-yl) hept-6-en-1 one (or Methylidihydromorphine)	$C_{18}H_{23}NO_3$	Opioid		46.6023	2.34	Hepatoprotective effect <sup>[54]</sup>
				47.8646	1.18	No activity reported

Contid...

Table 2: Contd...

Identified compounds	Formula	Nature of compound	Molecular massb (g/mol)	Retention time	Area (%)	Biological activity
Pyrrolidine, 1-[5-(1,3-benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]-, (E, E)- (or Pyrrolidine, Trichostachine, Piperiline)	C <sub>16</sub> H <sub>17</sub> NO <sub>3</sub>	Alkaloids	271.32	47.9359	2.58	Antiproliferative effect, cycle arrest, induce apoptosis on MCF-7 cells and antitumor effect <i>in vivo</i> <sup>[55]</sup>
1H-Indene, 2-fluoro-2,3-dihydro-1-methoxy-, trans-(+)- (E)-1-(Piperidin-1-yl) octadec-2-en-1-one (or Piperitine)	C <sub>10</sub> H <sub>11</sub> FO C <sub>23</sub> H <sub>43</sub> NO C <sub>19</sub> H <sub>25</sub> NO <sub>3</sub>	Amides Alkaloids Alkaloids	349.60 315.41	48.1182 48.3679 48.5620	0.66 21.66 0.24	No activity reported Insecticidal activity <sup>[25]</sup> No activity reported
(E)-7-(Benzo[d][1,3]dioxol-5-yl)-1-(piperidin-1-yl) hept-6-en-1-one (or Piperolein A)	C <sub>19</sub> H <sub>23</sub> NO <sub>3</sub>	Alkaloids	313.39	49.1390	5.65	Cytotoxicity against CCRF-CEM (acute lymphoblastic leukemia), HL-60 (acute promyelocytic leukemia), PC-3 (prostate carcinoma), and HA22T (hepatoma) cells <sup>[27]</sup>
(2E,6E)-7-(Benzo[d][1,3]dioxol-5-yl)-1-(piperidin-1-yl) hepta-2,6-dien-1-one (or Pipersintenamide)						Cytoprotective activity on normal fibroblast L929 cells and hepatoprotective activity <sup>[54]</sup>
(2E,4E,14E)-N-Isobutylicos-2,4,1,4-trienamide (or 2,4,14-Eicosatrienamide)	C <sub>24</sub> H <sub>43</sub> NO	Amides	361.61	49.3379	0.59	Antiinflammatory, <sup>[56]</sup> anticancer <sup>[32]</sup>
2-Furanol, 3,4-bis (1,3-benzodioxol-5-ylmethyl) tetrahydro- (or 2-Furanol, Cubebin)	C <sub>20</sub> H <sub>20</sub> O <sub>6</sub>	Lignan	356.37	49.6489	0.28	
Retrofractamide-A	C <sub>20</sub> H <sub>25</sub> NO <sub>3</sub>	Alkaloids	327.42	50.3585	0.34	Larvicidal activity against <i>Culex pipiens</i> pallens, <i>Aedes aegypti</i> and <i>Aedes togoi</i> ; <sup>[57]</sup> hepatoprotective effect <sup>[54]</sup>
2 (3H)-Furanone, 3,4-bis (1,3-benzodioxol-5-ylmethyl) dihydro-, (3R-trans)- (or (+)-Hinokinin, Cubebinolide)	C <sub>20</sub> H <sub>18</sub> O <sub>6</sub>	Lignan	354.36	50.5191	1.13	Antiinflammatory, <sup>[58]</sup> antioxidant <sup>[59]</sup>
(E)-9-(Benzo[d][1,3]dioxol-5-yl)-1-(pyrrolidin-1-yl) non-8-en-1-one (or Pyrrolidine, Tricholeine)	C <sub>20</sub> H <sub>27</sub> NO <sub>3</sub>	Alkaloids	329.44	50.7269	0.42	Antiproliferative activity against various cancer cells <sup>[60]</sup>
(3R,4R)-3-(Benzo[d][1,3]dioxol-5-yl methyl)-4-(3,4-dimethoxybenzyl) dihydrofuran-2 (3H) one (or Kusunokinin)	C <sub>21</sub> H <sub>22</sub> O <sub>6</sub>	Lignan	370.40	51.0435	1.28	Anticancer; <sup>[31]</sup> insecticidal activity against <i>Virola sebifera</i> and fungicidal activity against <i>Leucoagaricus gongylophorus</i> <sup>[61]</sup>
(E)-9-(Benzo[d][1,3]dioxol-5-yl)-1-(piperidin-1-yl) non-8-en-1-one (or Piperolein B)	C <sub>21</sub> H <sub>29</sub> NO <sub>3</sub>	Alkaloids	343.47	51.3920	1.03	Inhibitor of acyl CoA: Diacylglycerol acyltransferase for potential therapy for the treatment of obesity and type 2 diabetes <sup>[62]</sup>
(2E,4E,12E)-13-(Benzo[d][1,3]dioxol-5-yl)-N-isobutytrideca-2,4,12-trienamide (or Guineensine)	C <sub>24</sub> H <sub>33</sub> NO <sub>3</sub>	Alkaloids	383.53	51.8600	10.17	Antiinflammatory <sup>[63]</sup>
(2E,4E,6E)-7-(Benzo[d][1,3]dioxol-5-yl)-1-(piperidin-1-yl) hepta-2,4,6-trien-1-one (or Piperitine)	C <sub>19</sub> H <sub>21</sub> NO <sub>3</sub>	Alkaloids	311.38	52.9692	0.31	Trypanocidal effects against epimastigotes and amastigotes of <i>Trypanosoma cruzi</i> <sup>[64]</sup>
(22E)-Stigmasta-5,22-dien-3-ol (or beta-Stigmasterol, Poriferasterol)	C <sub>29</sub> H <sub>48</sub> O	Steroid	412.70	53.0319	1.74	Induce DNA damage and cell death <sup>[65]</sup>
(2E,4E,8E)-9-(Benzo[d][1,3]dioxol-5-yl)-1-(piperidin-1-yl) nona-2,4,8-trien-1-one (or Dehydroperipernonaline)	C <sub>21</sub> H <sub>25</sub> NO <sub>3</sub>	Alkaloids	339.47	53.5356	2.32	Coronary vasodilating activity <sup>[66]</sup>
gamma-Sitosterol (or clonasterol)	C <sub>29</sub> H <sub>50</sub> O	Terpenes	414.72	53.7147	0.48	Cytotoxicity against P388 (murine lymphocytic leukaemia) and HL60 (leukemia) cells <sup>[67]</sup>
(2E,4E,12E)-13-(Benzo[d][1,3]dioxol-5-yl)-Nisobutytrideca-2,4,12-trienamide (or Guineensine)	C <sub>24</sub> H <sub>33</sub> NO <sub>3</sub>	Alkaloids	383.53	55.6810		Antiinflammatory <sup>[63]</sup>

at 5.09% [Table 2]. Similarly, CP2 (PFPE) exhibited  $IC_{50}$  values of  $7.45 \pm 1.59 \mu\text{g/ml}$  in MCF-7 cell lines, which was better than DPCE ( $IC_{50}$  at  $23.46 \pm 1.10 \mu\text{g/ml}$ ).<sup>[17]</sup> These results indicate that PFPE, less piperine, was a potential crude extract in anticancer.

*O. viverrini* excretory/secretory products and *O. viverrini* antigen induce the expression of TLR4, IL-6, IL-8, TLR2, NF- $\kappa$ B, iNOS and COX-2 causing damage to biliary epithelium.<sup>[68]</sup> In this current study, PFPE showed down regulation of NF- $\kappa$ B, STAT-3 and COX-2 proteins [Figure 2]. In cancer cells, NF- $\kappa$ B and STAT-3 are major transcription factors that regulate proliferation, inflammatory, angiogenesis, invasive and apoptosis resistance by induction of several proteins, such as cyclin D, cyclin E1, CDK2, CDK4, CDK6, c-myc, tumor necrosis factor alpha, interleukin-1 (IL-1), IL-6, IL-8, VEGF and MMP-9.<sup>[69]</sup> NF- $\kappa$ B and STAT-3 proteins are induced by IL-6 to stimulate COX-2 expression in the inflammation process and cell cycle,<sup>[70,71]</sup> which associate to CCA progression. Therefore, suppression of NF- $\kappa$ B, STAT-3 and COX-2 proteins cause cancer growth inhibition. Piperlongumine,

an alkaloid from *P. longum* reduces NF- $\kappa$ B and c-Myc protein levels and inhibits binding of NF- $\kappa$ B with DNA at promoters in lymphoma cancer cells.<sup>[72]</sup> Moreover, piperlongumine also reduced the phosphorylation of JAK-1, JAK-2 and STAT-3 in gastric cancer cells.<sup>[73]</sup> Matrine, an alkaloid from *Sophora flavescens* Ait., significantly inhibits the viability by reduction the phosphorylation levels of JAK-2 and STAT3 proteins in CCA cells.<sup>[74]</sup> Curcumin, a natural extracted polyphenol from *C. longa*, also suppresses proliferation in human biliary cancer cells through inhibition of NF- $\kappa$ B, STAT-3 and JAK1 proteins.<sup>[75]</sup>


There are many evidences on genes and proteins which relate to bile duct cancer growth and progression, such as p53 mutation, inactivation of p21 and activation of Ras and MAPKs proteins.<sup>[76]</sup> Here, we found that PFPE could inhibit CCA cancer proliferation by decreasing of topoisomerase II, Akt, c-Myc, cyclin D1, and increasing of p21 protein levels [Figure 4]. Topoisomerase II is an enzyme involved in the DNA replication process that controls cell cycle with peaking at G2/M phase.<sup>[77]</sup> Therefore, down regulation of topoisomerase II by PFPE could induced DNA damage, interrupted cell growth and caused cell death on KKKU-M213 and TFK-1 cells. Most of the clinically active agents, including etoposide (lignan) and doxorubicin (alkaloid) are topoisomerase inhibitors.<sup>[78]</sup> Previously andrographolide analogue 3A.1 from *Andrographis paniculata*, a diterpenoid lactone, induces cell cycle arrest by down-regulation of CDK6 and cyclin D1 in KKKU-M213 cell lines.<sup>[79]</sup> Surprisingly, PFPE also exerted a significant reduction of Akt protein leading to decreasing of c-Myc and cyclin D1 and increasing of p21 levels [Figure 6]. Akt and cyclin D1 stimulate the cell cycle progression from G1/S phase to G2/M phase.<sup>[80]</sup>  $\beta$ -caryophyllene oxide, a terpene compound from *P. nigrum*, shows down-regulation of downstream of AKT pathway, including cyclin D1, COX-2 and VEGF and also up-regulation of p53 and p21 proteins in human prostate and breast cancer cells.<sup>[81]</sup>

In this study, we founded that the PFPE induced cell death by causing DNA fragmentation, increasing apoptotic proteins (p53, Bax and PUMA) and decreasing Bcl-2 protein levels [Figure 5]. p53, a tumor suppressor and transcription factor, is initially induced when DNA

**Table 3:** Cytotoxicity of piperine free *Piper nigrum* extract against cholangiocarcinoma, cholangiocyte and normal mouse fibroblast cell lines


Cell lines	$IC_{50}$ value $\pm$ SD ( $\mu\text{g/ml}$ )			
	DPCE	Piperine	PFPE	Doxorubicin
CCA				
KKU-100	22.88 $\pm$ 0.43	46.53 $\pm$ 0.09	17.79 $\pm$ 0.88	0.78 $\pm$ 0.03
KKU-M213	22.22 $\pm$ 0.26	27.01 $\pm$ 0.36	13.70 $\pm$ 1.14	1.75 $\pm$ 0.02
KKU-M055	46.66 $\pm$ 0.48	55.32 $\pm$ 0.22	16.74 $\pm$ 0.61	0.69 $\pm$ 0.09
TFK-1	23.25 $\pm$ 0.45	29.38 $\pm$ 0.07	15.30 $\pm$ 0.18	15.19 $\pm$ 0.12
HuCC-T1	37.17 $\pm$ 0.03	35.02 $\pm$ 0.12	20.72 $\pm$ 0.75	2.53 $\pm$ 0.04
Normal cholangiocyte				
MMNK-1	33.25 $\pm$ 0.28	60.68 $\pm$ 0.72	19.65 $\pm$ 0.26	0.62 $\pm$ 0.05
Normal fibroblast				
L-929	No effect	No effect	45.53 $\pm$ 0.50	0.20 $\pm$ 0.01

*P. nigrum*: *Piper nigrum*; DPCE: Dichloromethane *P. nigrum* crude extract; PFPE: Piperine free *P. nigrum* extract; CCA: Cholangiocarcinoma; SD: Standard deviation




**Figure 3:** Expression of inflammation-related proteins in KKKU-M213 (a and c) and TFK-1 (b and d) cells treated with piperine free *Piper nigrum* extract at 24, 48, 72 and 96 h. The levels of signal transducer and activator of transcription 3, cyclooxygenase-2 and Nuclear factor kappa-light-chain-enhancer of activated B cells and GAPDH proteins were measured using the Western blot analysis. Densitometric analysis normalized to GAPDH. Data were represented as mean  $\pm$  standard deviation and three independent experiments were done. \* $P < 0.05$  compared with control group (0 h)





**Figure 4:** Effect of piperine free *Piper nigrum* extract on cell growth and cell cycle arrest. KKU-M213 (a and c) and TFK-1 (b and d) cells were treated with Median inhibition concentration concentration of piperine free *Piper nigrum* extract for 24, 48, 72 and 96 h. Then, the levels of topoisomerase II, AKT8 virus oncogene cellular homolog, avian myelocytomatosis virus oncogene cellular homolog, cyclin D1 and p21 proteins were investigated using Western blot analysis. Fold change of each protein was measured by densitometry quantitation using ImageJ software and normalized with GAPDH.  $P < 0.05$  of three independent experiments was considered to indicate a statistically significant differences compared to control group (0 h)




**Figure 5:** Effect of piperine free *Piper nigrum* extract on apoptosis. KKU-M213 (a and c) and TFK-1 (b and d) cells were treated with Median inhibition concentration concentration of piperine free *Piper nigrum* extract for 24, 48, 72 and 96 h. Then, the levels of tumor protein p53, B-cell lymphoma 2, Bcl-2-associated X protein and PUMA proteins were investigated using Western blot analysis. Fold change of each protein was measured by densitometry quantitation using ImageJ software and normalized with GAPDH.  $P < 0.05$  of three independent experiments was considered to indicate a statistically significant difference compared to control group (0 h)

damage and takes responsibility to activate several apoptotic genes, such as Bax, PUMA and NOXA.<sup>[82-84]</sup> Similarly, ethanolic extract of *P. nigrum* has antiproliferative effect on MCF-7 cells, antitumor effect *in vivo* and triggering apoptosis via p53 and Bax and decreasing of Bcl-2 proteins.<sup>[55]</sup> Curcumin effectively induces apoptosis in CCA (CCLP-1 and SG-231) cells by stimulation of Notch1, Hes-1 and survivin apoptotic proteins.<sup>[85]</sup> Andrographolide analog 3A.1 has cytotoxicity

with  $IC_{50}$  of 8.0  $\mu$ M on KKU-M213 cells at 24 h after treatment and induces apoptosis via induction of cleaved PARP-1, Bax, caspase-3, and p53.<sup>[79]</sup> Matrine stimulates apoptosis in CCA cells through induction of cytochrome c releasing from mitochondria and reduction of caspase-3 and-9 activity.<sup>[74]</sup> Taken together, PFPE can be a potential candidate for CCA treatment in future. However, study in CCA *in vivo* and clinical trial need to be carried out.





**Figure 6:** The anticancer mechanism of piperine free *Piper nigrum* extract in cholangiocarcinoma

## CONCLUSION

PFPE showed strong cytotoxicity against KKU-M213 and TFK-1 cell lines with  $IC_{50}$  values of  $13.70 \pm 1.14$  and  $15.30 \pm 0.18$   $\mu\text{g/ml}$ , respectively. PFPE suppressed inflammation through down-regulation of NF- $\kappa$ B, STAT-3 and COX-2. Moreover, PFPE inhibited CCA cells growth and proliferation by down-regulation of topoisomerase II, Akt, c-Myc and cyclin D and up-regulation of p21. Furthermore, PFPE triggered apoptosis through inhibition of Bcl-2 and induction of p53, Bax and PUMA levels as summarized in the Figure 5. In summary, PFPE can be served as a promising crude extract for CCA treatment.

## Acknowledgement

The authors thank Prince of Songkla University for the equipment funding (the Excellent Research Laboratory (ExLab004)).

## Financial support and sponsorship

This work was supported by the research fund (REC 60-279-04-2) Faculty of Medicine and the Postdoctoral Fellowship, Prince of Songkla University.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Blechacz B, Gores GJ. Cholangiocarcinoma: Advances in pathogenesis, diagnosis, and treatment. *Hepatology* 2008;48:308-21.
- Ghouri YA, Mian I, Blechacz B. Cancer review: Cholangiocarcinoma. *J Carcinog* 2015;14:1.
- Saha SK, Zhu AX, Fuchs CS, Brooks GA. Forty-year trends in cholangiocarcinoma incidence in the U.S.: Intrahepatic disease on the rise. *Oncologist* 2016;21:594-9.
- Khan SA, Taylor-Robinson SD, Toledano MB, Beck A, Elliott P, Thomas HC. Changing international trends in mortality rates for liver, biliary and pancreatic tumours. *J Hepatol* 2002;37:806-13.
- Sripa B, Pairojkul C. Cholangiocarcinoma: Lessons from Thailand. *Curr Opin Gastroenterol* 2008;24:349-56.
- Plentz RR, Malek NP. Clinical presentation, risk factors and staging systems of cholangiocarcinoma. *Best Pract Res Clin Gastroenterol* 2015;29:245-52.
- Gatto M, Alvaro D. New insights on cholangiocarcinoma. *World J Gastrointest Oncol* 2010;2:136-45.
- Squadroni M, Tondulli L, Gatta G, Mosconi S, Beretta G, Labianca R. Cholangiocarcinoma. *Crit Rev Oncol Hematol* 2017;116:11-31.

- van der Gaag NA, Kloek JJ, de Bakker JK, Musters B, Geskus RB, Busch OR, *et al.* Survival analysis and prognostic nomogram for patients undergoing resection of extrahepatic cholangiocarcinoma. *Ann Oncol* 2012;23:2642-9.
- Barreto JN, McCullough KB, Ice LL, Smith JA. Antineoplastic agents and the associated myelosuppressive effects: A review. *J Pharm Pract* 2014;27:440-6.
- Ahmad N, Fazal H, Abbasi BH, Farooq S, Ali M, Khan MA. Biological role of *Piper nigrum* L. (Black pepper): A review. *Asian Pac J Trop Biomed* 2012;2:S1945-53.
- Parmar VS, Jain SC, Bisht KS, Jain R, Taneja P, Jha A, *et al.* Phytochemistry of the genus *Piper*. *Phytochemistry* 1997;46:597-673.
- Prashant A, Rangaswamy C, Yadav AK, Reddy V, Sowmya MN, Madhunapantula S. Anticancer activity of ethanolic extracts of & against colorectal carcinoma cell lines. *Int J Appl Basic Med Res* 2017;7:67-72.
- Grinevicius VM, Andrade KS, Ourique F, Micke GA, Ferreira SR, Pedrosa RC. Antitumor activity of conventional and supercritical extracts from *Piper nigrum* L. cultivar Bragantina through cell cycle arrest and apoptosis induction. *J Supercrit Fluids* 2017;128:94-101.
- Sriwiriyan S, Tedasen A, Lailerd N, Boonyaphiphat P, Nitiruangjarat A, Deng Y, *et al.* Anticancer and cancer prevention effects of piperine-free *Piper nigrum* extract on n-nitrosomethylurea-induced mammary tumorigenesis in rats. *Cancer Prev Res (Phila)* 2016;9:74-82.
- Deng Y, Sriwiriyan S, Tedasen A, Hiransai P, Graidist P. Anti-cancer effects of *Piper nigrum* via inducing multiple molecular signaling *in vivo* and *in vitro*. *J Ethnopharmacol* 2016;188:87-95.
- Sriwiriyan S, Ninpesh T, Sukpondma Y, Nasomyon T, Graidist P. Cytotoxicity screening of plants of genus *Piper* in breast cancer cell lines. *Trop J Pharm Res* 2014;13:921-8.
- Geran RI, Greenberg NH, McDonald MM, Schumacher AM, Abbot BJ. Protocol for screening chemical agents and natural products against animal tumors and other biological systems. *Cancer Chemoth Rep* 1972;3:1-103.
- Graidist P, Martla M, Sukpondma Y. Cytotoxic activity of *Piper cubeba* extract in breast cancer cell lines. *Nutrients* 2015;7:2707-18.
- Plengsuriyakarn T, Thitapakorn V, Na-Bangchang K, Karbwang J. Thai medicinal plants: Potential sources of anticholangiocarcinoma drugs. *Int J Pharm Pharmacol* 2013;2:68-82.
- Leelawat S, Leelawat K. Molecular mechanisms of cholangiocarcinoma cell inhibition by medicinal plants. *Oncol Lett* 2017;13:961-6.
- Zainal Ariffin SH, Wan Omar WH, Zainal Ariffin Z, Safian MF, Senafi S, Megat Abdul Wahab R. Intrinsic anticarcinogenic effects of *Piper sarmentosum* ethanolic extract on a human hepatoma cell line. *Cancer Cell Int* 2009;9:6.
- OVADJE P, Ma D, Tremblay P, Roma A, Steckle M, Guerrero JA, *et al.* Evaluation of the efficacy biochemical mechanism of cell death induction by *Piper longum* extract selectively in *in vitro* and *in vivo* models of human cancer cells. *PLoS One* 2014;9:e113250.
- Mahavorasirikul W, Viyanant V, Chaijaroenkul W, Itharat A, Na-Bangchang K. Cytotoxic activity of Thai medicinal plants against human cholangiocarcinoma, laryngeal and hepatocarcinoma cells *in vitro*. *BMC Complement Altern Med* 2010;10:55.
- Geris R, Ribeiro PR, Brandão MD, Da Silva HH, Silva IG. Bioactive natural products as potential candidates to control *Aedes aegypti*, the vector of dengue. In: Rahman A, editor. *Studies in Natural Products Chemistry*. Vol. 37. Oxford: Elsevier; 2012. p. 227-376.
- Muharini R, Liu Z, Lin W, Proksch P. New amides from the fruits of *Piper retrofractum*. *Tetrahedron Lett* 2015;56:2521-5.
- Chen JJ, Huang YC, Chen YC, Huang YT, Wang SW, Peng CY, *et al.* Cytotoxic amides from *Piper sintonense*. *Planta Med* 2002;68:980-5.
- Chen JJ, Duh CY, Huang HY, Chen IS. Cytotoxic Constituents of *Piper sintonense*. *Helv Chim Acta* 2003;86:2058-64.
- Dahham SS, Tabana YM, Iqbal MA, Ahamed MB, Ezzat MO, Majid AS, *et al.* The Anticancer, antioxidant and antimicrobial properties of the sesquiterpene  $\beta$ -caryophyllene from the essential oil of *Aquilaria crassna*. *Molecules* 2015;20:11808-29.
- Shen T, Zhang L, Wang YY, Fan PH, Wang XN, Lin ZM, *et al.* Steroids from Commiphora mukul display antiproliferative effect against human prostate cancer PC3 cells via induction of apoptosis. *Bioorg Med Chem Lett* 2012;22:4801-6.
- Sriwiriyan S, Sukpondma Y, Srisawat T, Madla S, Graidist P. (-)-Kusunokinin and piperloguminine from *Piper nigrum*: An alternative option to treat breast cancer. *Biomed Pharmacother* 2017;92:732-43.
- Rajalekshmi DS, Kabeer FA, Madhusoodhanan AR, Bahulayan AK, Prathapan R, Prakasan N, *et al.* Anticancer activity studies of cubebin isolated from *Piper cubeba* and its synthetic derivatives. *Bioorg Med Chem Lett* 2016;26:1767-71.
- Kohoude MJ, Gbaguidi F, Agbani P, Ayedoun MA, Cazaux S, Bouajila J. Chemical composition

- and biological activities of extracts and essential oil of *Boswellia dalzielii* leaves. *Pharm Biol* 2017;55:33-42.
34. Rabi T, Bishayee A. d-Limonene sensitizes docetaxel-induced cytotoxicity in human prostate cancer cells: Generation of reactive oxygen species and induction of apoptosis. *J Carcinog* 2009;8:9.
35. Zou L, Liu W, Yu L. Beta-elemene induces apoptosis of K562 leukemia cells. *Zhonghua Zhong Liu Za Zhi* 2001;23:196-8.
36. Martins Cde M, do Nascimento EA, de Moraes SA, de Oliveira A, Chang R, Cunha LC, *et al.* Chemical constituents and evaluation of antimicrobial and cytotoxic activities of *Kielmeyera coriacea* Mart. & Zucc. essential oils. *Evid Based Complement Alternat Med* 2015;2015:1-9. doi.org/10.1155/2015/842047.
37. Baldissera MD, Souza CF, Grando TH, Stefani LM, Monteiro SG.  $\beta$ -caryophyllene reduces atherogenic index and coronary risk index in hypercholesterolemic rats: The involvement of cardiac oxidative damage. *Chem Biol Interact* 2017;270:9-14.
38. Alvarez-González I, Madrigal-Bujaidar E, Castro-García S. Antigenotoxic capacity of beta-caryophyllene in mouse, and evaluation of its antioxidant and GST induction activities. *J Toxicol Sci* 2014;39:849-59.
39. Ali NA, Wursterb M, Denkert A, Arnold N, Fadail I, Al-Didamony G, *et al.* Chemical composition, antimicrobial, antioxidant and cytotoxic activity of essential oils of *Plectranthus cyllindraceus* and *Meriandra benghalensis* from Yemen. *Nat Prod Commun* 2012;7:1099-102.
40. Keawsa-ard S, Liawruangrath B, Liawruangrath S, Teerawutgulrag A, Pyne SG. Chemical constituents and antioxidant and biological activities of the essential oil from leaves of *Solanum spirale*. *Nat Prod Commun* 2012;7:955-8.
41. Ali NAA, Chhetri BK, Dosoky NS, Shari K, Al-Fahad AJ, Wessjohann L, *et al.* Antimicrobial, antioxidant, and cytotoxic activities of *Ocimum forskolei* and *Teucrium yemense* (Lamiaceae) essential oils. *Medicines (Basel)* 2017;4:7.
42. Hui LM, Zhao GD, Zhao JJ.  $\delta$ -Cadinene inhibits the growth of ovarian cancer cells via caspase-dependent apoptosis and cell cycle arrest. *Int J Clin Exp Pathol* 2015;8:6046-56.
43. Di Giacomo S, Di Sotto A, Mazzanti G, Wink M. Chemosensitizing properties of  $\beta$ -Caryophyllene and  $\beta$ -caryophyllene oxide in combination with doxorubicin in human cancer cells. *Anticancer Res* 2017;37:1191-6.
44. Fidyk T, Fiedorowicz A, Strzadala L, Szumny A.  $\beta$ -caryophyllene and  $\beta$ -caryophyllene oxide-natural compounds of anticancer and analgesic properties. *Cancer Med* 2016;5:3007-17.
45. Hanušová V, Caltová K, Svobodová H, Ambroz M, Skarka A, Murínová N, *et al.* The effects of  $\beta$ -caryophyllene oxide and trans-nerolidol on the efficacy of doxorubicin in breast cancer cells and breast tumor-bearing mice. *Biomed Pharmacother* 2017;95:828-36.
46. Kim C, Cho SK, Kapoor S, Kumar A, Vali S, Abbasi T, *et al.*  $\beta$ -Caryophyllene oxide inhibits constitutive and inducible STAT3 signaling pathway through induction of the SHP-1 protein tyrosine phosphatase. *Mol Carcinog* 2014;53:793-806.
47. Dosoky NS, Satyal P, Gautam TP, Setzer WN. Composition and biological activities of & *Murraya paniculata* (L.) Jack essential oil from Nepal. *Medicines (Basel)* 2016;3:7.
48. Ku SK, Lee IC, Kim JA, Bae JS. Anti-septic effects of pellitorine in HMGB1-induced inflammatory responses *in vitro* and *in vivo*. *Inflammation* 2014;37:338-48.
49. Stöhr JR, Xiao PG, Bauer R. Constituents of Chinese *Piper* species and their inhibitory activity on prostaglandin and leukotriene biosynthesis *in vitro*. *J Ethnopharmacol* 2001;75:133-9.
50. Gulledge TV, Collette NM, Mackey E, Johnstone SE, Moazami Y, Todd DA, *et al.* Mast cell degranulation and calcium influx are inhibited by an *Echinacea purpurea* extract and the alkylamide dodeca-2E,4E-dienoic acid isobutylamide. *J Ethnopharmacol* 2018;212:166-74.
51. Matsuda H, Ninomiya K, Morikawa T, Yasuda D, Yamaguchi I, Yoshikawa M. Protective effects of amide constituents from the fruit of *Piper chaba* on D-galactosamine/TNF- $\alpha$ -induced cell death in mouse hepatocytes. *Bioorg Med Chem Lett* 2008;18:2038-42.
52. Gunasekaran V, Elangovan K, Niranjali Devaraj S. Targeting hepatocellular carcinoma with piperine by radical-mediated mitochondrial pathway of apoptosis: An *in vitro* and *in vivo* study. *Food Chem Toxicol* 2017;105:106-18.
53. Han SZ, Liu HX, Yang LQ, Cui LD, Xu Y. Piperine (PP) enhanced mitomycin-C (MMC) therapy of human cervical cancer through suppressing Bcl-2 signaling pathway via inactivating STAT3/NF- $\kappa$ B. *Biomed Pharmacother* 2017;96:1403-10.
54. Matsuda H, Ninomiya K, Morikawa T, Yasuda D, Yamaguchi I, Yoshikawa M. Hepatoprotective amide constituents from the fruit of *Piper chaba*: Structural requirements, mode of action, and new amides. *Bioorg Med Chem* 2009;17:7313-23.
55. de Souza Grinevicius VM, Kwiecinski MR, Santos Mota NS, Ourique F, Porfirio Will Castro LS, Andregueti RR, *et al.* *Piper nigrum* ethanolic extract rich in piperamides causes ROS overproduction, oxidative damage in DNA leading to cell cycle arrest and apoptosis in cancer cells. *J Ethnopharmacol* 2016;189:139-47.
56. Lima TC, Lucarini R, Volpe AC, de Andrade CQJ, Souza AM, Pauletti PM, *et al.* *In vivo* and *in silico* anti-inflammatory mechanism of action of the semisynthetic (-)-cubebin derivatives (-)-hinokinin and (-)-O-benzylcubebin. *Bioorg Med Chem Lett* 2017;27:176-9.
57. Park IK, Lee SG, Shin SC, Park JD, Ahn YJ. Larvicidal activity of isobutylamides identified in *Piper nigrum* fruits against three mosquito species. *J Agric Food Chem* 2002;50:1866-70.
58. Godoy de Lima R, Barros MT, da Silva Laurentiz R. Medicinal attributes of Lignans extracted from & *Piper cubeba*: Current developments. *ChemistryOpen* 2018;7:180-91.
59. Medola JF, Cintra VP, Pesqueira E Silva EP, de Andrade Royo V, da Silva R, Saraiva J, *et al.* (-)-Hinokinin causes antigenotoxicity but not genotoxicity in peripheral blood of Wistar rats. *Food Chem Toxicol* 2007;45:638-42.
60. Wang YH, Goto M, Wang LT, Hsieh KY, Morris-Natschke SL, Tang GH, *et al.* Multidrug resistance-selective antiproliferative activity of *Piper* amide alkaloids and synthetic analogues. *Bioorg Med Chem Lett* 2014;24:4818-21.
61. Bicalho KU, Terezan AP, Martins DC, Freitas TG, Fernandes JB, Silva MF, *et al.* Evaluation of the toxicity of *Virola sebifera* crude extracts, fractions and isolated compounds on the nest of leaf-cutting ants. *Psyche* 2012;2012:785424.
62. Lee SW, Rho MC, Park HR, Choi JH, Kang JY, Lee JW, *et al.* Inhibition of diacylglycerol acyltransferase by alkamides isolated from the fruits of *Piper longum* and *Piper nigrum*. *J Agric Food Chem* 2006;54:9759-63.
63. Reynoso-Moreno I, Najar-Guerrero I, Escareño N, Flores-Soto ME, Gertsch J, Viveros-Paredes JM. An endocannabinoid uptake inhibitor from black pepper exerts pronounced anti-inflammatory effects in mice. *J Agric Food Chem* 2017;65:9435-42.
64. Ribeiro TS, Freire-de-Lima L, Previato JO, Mendonça-Previato L, Heise N, de Lima ME. Toxic effects of natural piperine and its derivatives on epimastigotes and amastigotes of *Trypanosoma cruzi*. *Bioorg Med Chem Lett* 2004;14:3555-8.
65. Chaturvedula VS, Gao Z, Hecht SM, Jones SH, Kingston DG. A new acylated oleanane triterpenoid from *Couepia polyandra* that inhibits the lyase activity of DNA polymerase  $\beta$ . *J Nat Prod* 2003;66:1463-65.
66. Shoji N, Umeyama A, Saito N, Takemoto T, Kajiwara A, Ohizumi Y. Dehydropipernonaline, an amide possessing coronary vasodilating activity, isolated from *Piper longum* L. *J Pharm Sci* 1986;75:1188-9.
67. Manoharan KP, Yang D, Hsu A, Huat BT. Evaluation of *Polygonum bistorta* for anticancer potential using selected cancer cell lines. *Med Chem* 2007;3:121-6.
68. Prueksapanich P, Piyaichaturawat P, Aumpansub P, Ridditid W, Chaiterakij R, Rerknimit R. Liver fluke-associated biliary tract cancer. *Gut Liver* 2018;12:236-45.
69. Fan Y, Mao R, Yang J. NF- $\kappa$ B and STAT3 signaling pathways collaboratively link inflammation to cancer. *Protein Cell* 2013;4:176-85.
70. Basu A, Das AS, Sharma M, Pathak MP, Chattopadhyay P, Biswas K, *et al.* STAT3 and NF- $\kappa$ B are common targets for kaempferol-mediated attenuation of COX-2 expression in IL-6-induced macrophages and carrageenan-induced mouse paw edema. *Biochem Biophys Res* 2017;12:54-61.
71. Chen SC, Chang YL, Wang DL, Cheng JJ. Herbal remedy magnolol suppresses IL-6-induced STAT3 activation and gene expression in endothelial cells. *Br J Pharmacol* 2006;148:226-32.
72. Han SS, Son DJ, Yun H, Kamberos NL, Janz S. Piperlongumine inhibits proliferation and survival of Burkitt lymphoma *in vitro*. *Leuk Res* 2013;37:146-54.
73. Song B, Zhan H, Bian Q, Gu J. Piperlongumine inhibits gastric cancer cells via suppression of the JAK1,2/STAT3 signaling pathway. *Mol Med Rep* 2016;13:4475-80.
74. Yang N, Han F, Cui H, Huang J, Wang T, Zhou Y, *et al.* Matrine suppresses proliferation and induces apoptosis in human cholangiocarcinoma cells through suppression of JAK2/STAT3 signaling. *Pharmacol Rep* 2015;67:388-93.
75. Prakobwong S, Gupta SC, Kim JH, Sung B, Pinlaor P, Hiraku Y, *et al.* Curcumin suppresses proliferation and induces apoptosis in human biliary cancer cells through modulation of multiple cell signaling pathways. *Carcinogenesis* 2011;32:1372-80.
76. Fava G. Molecular mechanisms of cholangiocarcinoma. *World J Gastrointest Pathophysiol* 2010;1:12-22.
77. Larsen AK, Skladanowski A, Bojanowski K. The roles of DNA topoisomerase II during the cell cycle. *Prog Cell Cycle Res* 1996;2:229-39.
78. Nitiss JL. Targeting DNA topoisomerase II in cancer chemotherapy. *Nat Rev Cancer* 2009;9:338-50.
79. Nateewattana J, Dutta S, Reabroi S, Saeeng R, Kasemsook S, Chairoungdua A, *et al.* Induction of apoptosis in cholangiocarcinoma by an andrographolide analogue is mediated through topoisomerase II  $\alpha$  inhibition. *Eur J Pharmacol* 2014;723:148-55.

80. Xu N, Lao Y, Zhang Y, Gillespie DA. Akt: A double-edged sword in cell proliferation and genome stability. *J Oncol* 2012;2012:1-15. doi:10.1155/2012/951724.
81. Park KR, Nam D, Yun HM, Lee SG, Jang HJ, Sethi G, *et al.*  $\beta$ -Caryophyllene oxide inhibits growth and induces apoptosis through the suppression of PI3K/AKT/mTOR/S6K1 pathways and ROS-mediated MAPKs activation. *Cancer Lett* 2011;312:178-88.
82. Miyashita T, Reed JC. Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell* 1995;80:293-9.
83. Nakano K, Vousden KH. PUMA, a novel proapoptotic gene, is induced by p53. *Mol Cell* 2001;7:683-94.
84. Yu J, Zhang L, Hwang PM, Kinzler KW, Vogelstein B. PUMA induces the rapid apoptosis of colorectal cancer cells. *Mol Cell* 2001;7:673-82.
85. Koprowski S, Sokolowski K, Kunnimalaiyaan S, Gamblin TC, Kunnimalaiyaan M. Curcumin-mediated regulation of Notch1/hairy and enhancer of split-1/survivin: molecular targeting in cholangiocarcinoma. *J Surg Res* 2015;198:434-40.