















19. Ordóñez AA, Gómez JD, Vattuone MA. Antioxidant activities of *Sechium edule* (Jacq.) Swartz extracts. *Food Chem* 2006;97:452-8.
20. Oyaizu M. Studies on products of browning reactions: Antioxidative activities of products of browning reaction prepared from glucosamine. *Jpn J Nutr* 1986;44:307-15.
21. Blois MS. Antioxidant determinations by the use of a stable free radical. *Nature* 1958;181:1199-200.
22. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C, *et al.* Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999;26:1231-7.
23. Seal T, Chaudhuri K, Pillai B. Nutraceutical and antioxidant properties of *Cucumis hardwickii* Royle: A potent wild edible fruit collected from Uttarakhand, India. *J Pharmacogn Phytochem* 2017;6:1837-47.
24. Dundar A, Okumus V, Ozdemir S, Celik KS, Boğa M, Ozcagli E. Determination of cytotoxic, anticholinesterase, antioxidant and antimicrobial activities of some wild mushroom species. *Cogent Food Agric* 2016;2:1178060.
25. Zieliński H, Kozłowska H. Antioxidant activity and total phenolics in selected cereal grains and their different morphological fractions. *J Agric Food Chem* 2000;48:2008-16.
26. Bravo L. Polyphenols: Chemistry, dietary sources, metabolism and nutritional significance. *Nutr Rev* 1998;56:317-33.
27. Anwar F, Jamil A, Iqbal S, Sheikh MA. Antioxidant activity of various plant extracts under ambient and accelerated storage of sunflower oil. *Grasas Aceites* 2006;57:189-97.
28. Seal T, Chaudhuri K, Pillai B. Effect of solvent extraction system on the antioxidant activity of some selected wild edible fruits of Meghalaya state in India. *J Chem Pharm Res* 2013;5:276-82.
29. Seal T, Pillai B, Chaudhuri K. Effect of solvent extraction system on the antioxidant activity of some selected wild leafy vegetables of Meghalaya state in India. *Int J Pharm Sci Res* 2013;4:1046-51.
30. Siddhuraju P, Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *J Agric Food Chem* 2003;51:2144-55.
31. Sultana B, Anwar F, Przybylski R. Antioxidant activity of phenolic components present in barks of *Azadirachta indica*, *Terminalia arjuna*, *Acacia nilotica* and *Eugenia jambolana* Lam. trees. *Food Chem* 2007;104:1106-14.
32. Jimoh FO, Adedapo AA, Afolayan AJ. Comparison of the nutritive value, antioxidant and antibacterial activities of *Sonchus asper* and *Sonchus oleraceus*. *Rec Nat Prod* 2011;5:29-42.
33. Pourmorad F, Hosseinimehr SJ, Shahabimajid N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr J Biotechnol* 2006;5:1142-45.
34. Jamuna KS, Ramesh CK, Srinivasa TR, Raghu KL. *In vitro* antioxidant studies in some common fruits. *Int J Pharm Pharm Sci* 2011;3:60-3.
35. Koleva II, Van Beek TA, Linssen JP, Groot AD, Evstatieva LN. Screening of plant extracts for antioxidant activity: A comparative study on three testing methods. *Phytochem Anal* 2002;13:8-17.
36. Mattila P, Könkö K, Eurolo M, Pihlava JM, Astola J, Vahteristo L, *et al.* Contents of vitamins, mineral elements and some phenolic compounds in cultivated mushrooms. *J Agric Food Chem* 2001;49:2343-8.
37. Bayili RG, Abdoul-Latif F, Kone OH, Diao M, Bassole IH, Dicko MH. Phenolic compounds and antioxidant activities in some fruits and vegetables from Burkina Faso. *Afr J Biotechnol* 2011;10:13543-7.
38. Masella R, Santangelo C, D'Archivio M, Li Volti G, Giovannini C, Galvano F, *et al.* Protocatechuic acid and human disease prevention: Biological activities and molecular mechanisms. *Curr Med Chem* 2012;19:2901-17.
39. Uma DB, Ho CW, Wan Aida WM. Optimization of extraction parameters of total phenolic compounds from Henna (*Lawsonia inermis*) leaves. *Sains Malaysiana* 2010;39:119-28.
40. Periyannan V, Veerasamy V. Syringic acid may attenuate the oral mucosal carcinogenesis via improving cell surface glycoconjugation and modifying cytokeratin expression. *Toxicol Rep* 2018;5:1098-106.
41. Karthikeyan R, Devadasu C, Srinivasa Babu P. Isolation, characterization and RP-HPLC estimation of p-coumaric acid from methanolic extract of durva grass (*Cynodon dactylon* Linn.) (Pers.). *Int J Analyt Chem* 2015;2015:1-7.
42. Mussatto SI, Dragone G, Roberto IC. Ferulic and p-coumaric acids extraction by alkaline hydrolysis of brewer's spent grain. *Ind Crops Prod* 2007;25:231-7.
43. Agarwal S, Sairam RK, Srivastava GC, Tyagi A, Meena RC. Role of ABA, salicylic acid, calcium and hydrogen peroxide on antioxidant enzymes induction in wheat seedlings. *Plant Sci* 2005;169:559-70.
44. Srinivasan K, Kaul CL, Ramarao P. Partial protective effect of rutin on multiple low dose streptozotocin-induced diabetes in mice. *Indian J Pharmacol* 2005;37:327.
45. Lin JP, Yang JS, Lin JJ, Lai KC, Lu HF, Ma CY, *et al.* Rutin inhibits human leukemia tumor growth in a murine xenograft model *in vivo*. *Environ Toxicol* 2012;27:480-4.
46. D'Andrea G. Quercetin: A flavonol with multifaceted therapeutic applications? *Fiterapia* 2015;106:256-71.
47. Seal T, Pillai B, Chaudhuri K. Identification and quantification of phenolic acids by HPLC, in two wild edible plants viz. *Solanum gilo* and *Solanum kurzii* collected from North-Eastern region in India. *J Chem Biol Phys Sci* 2016;6:1108-21.
48. Ahmed OM, Hassan MA, Abdel-Twab SM, Abdel Azeem MN. Navel orange peel hydroethanolic extract, naringin and naringenin have anti-diabetic potentials in type 2 diabetic rats. *Biomed Pharmacother* 2017;94:197-205.
49. Asif M, Khodadadi E. Medicinal uses and chemistry of flavonoid contents of some common edible tropical plants. *J Paramed Sci* 2013;4:119-38.
50. Calderon-Montano JM, Burgos-Morón E, Pérez-Guerrero C, López-Lázaro M. A review on the dietary flavonoid kaempferol. *Mini Rev Med Chem* 2011;11:298-344.