Simultaneous determination of five marker constituents in Ssanghwa tang by HPLC/DAD

Jin Bae Won¹, Jin Yeul Ma³, Young Ran Um³, Choong Je Ma^{1,2}

¹Department of Biomaterials Engineering, School of Biotechnology and Bioengineering, Kangwon National University, Chuncheon 200-701, ²Research Institute of Biotechnology, Kangwon National University, Chuncheon 200-701, ³TKM Converging Research Division, Korea Institute of Oriental Medicine, 483 Exporo, Yuseong-gu, Daejeon 305-811, Korea

Submitted: 12-01-2010

Revised: 22-02-2010

Published: 05-05-2010

ABSTRACT

A HPLC-DAD method was established for the simultaneous evaluation of five bioactive compounds in Ssanghwa tang (SHT) including glycyrrhizin, paeoniflorin, cinnamic acid, decursin and 6-gingerol. These compounds were separated in less than 40 min using a Dionex C₁₈ column with a gradient elution system of water and methanol at a flow rate of 1 ml/min. Calibration curve of standard components presented excellent linear regression ($R^2 > 0.9903$) within the test range. Limit of detection and limit of quantification varied from 0.07 to 0.46 μ g/ml and 0.13 to 1.11 μ g/ml, respectively. The relative standard deviations (RSDs) of data of the intraday and interday experiments were less than 3.67 and 5.73%, respectively. The accuracy of recovery test ranged from 95.98 to 105.88% with RSD values 0.10– 4.82%.

Key words: Herbal medicine, HPLC- DAD, quality control, quantification, Ssanghwa tang

INTRODUCTION

Traditional herbal medicines are usually prepared from various herbs, and they exhibit various therapeutic effects with a complex of multiplicity components.^[1,2] And the quality of these herbs has been affected by many factors such as collection time, place, temperature, cultivation environment and manufacturing process.^[3-5]

This suggests the necessity of the establishment of systematic quality evaluation. In fact, quality control method of single herb has been reported earlier.^[6-8] However, the quality control of traditional herbal medicinal preparation or decoction, a mixture of herb combination, has not been a research interest for a long time. Only several marker compounds were analyzed by simple HPLC analysis method. This method was fruitless as it was expensive and time consuming, and also it failed to ensure efficient quality control and standardization of traditional herbal medicine. Therefore, it is critical to develop a method for the simultaneous determination of several marker compounds in traditional herbal medicine.

Address for correspondence:

Prof. Choong Je Ma, Department of Biomaterials Engineering, School of Bioscience and Biotechnology, Kangwon National University, Hyoja-2 Dong, Chuncheon 200-701, Koria. E-mail: cjma@kangwon.ac.kr

DOI: 10.4103/0973-1296.62896

Ssanghwa tang (SHT) is a traditional herbal medicinal prescription used in the treatment of infirmity, refreshment congestion, sweat, convalescence recovery, and consists of *Paeonia lactiflora, Glycyrrhiza glabra, Rehmannia glutinosa, Astragalus membranaceus, Angelica gigas, Zingiber officinale, Zizyphus jujuba, Cinnamomum cassia*, and *Cnidium officinale.*^[9] In Korea, SHT has been commercially produced as granules by several medicinal manufactures. To ensure efficacy and safety, a suitable assay method for quality control is required.

In this study, we employed a HPLC-DAD method for the simultaneous determination of five marker constituents in SHT, glycyrrhizin, paeoniflorin, cinnamic acid, decursin, and 6-gingerol. In addition, established analysis method was applied for the analysis of various SHT samples. The method was successfully validated and it was used to simultaneously determine five important SHT compounds [Figure 1].

EXPERIMENTAL

Reagents and Materials

All the five standard compounds, paeoniflorin, *trans*cinnamic acid, glycyrrhizin, decursin and 6-gingerol, were purchased from Natural Product Chemistry BioTech Inc. (Seoul, Korea). The purity of all five marker constituents was more than 98%. All the plant materials were purchased

Won, et al.: Determination of five marker constituents in Ssanghwa tang



Figure 1: Chemical structures of five marker components in Ssanghwa tang

from Kyungdong traditional herbal market (Seoul, Korea). Five commercial brands of SHT granules were purchased from local providers. HPLC grade solvents (water and methanol) and reagents were obtained from J.T. Baker (USA).

Chromatographic conditions

Analysis was performed on the Dionex Ultimate 3000 HPLC system (Dionex, Germany) equipped with a pump (LPG 3X00), auto sampler (ACC-3000), column oven and diode array UV/VIS detector (DAD-3000(RS)). The output signal of the detector was recorded using a Dionex ChromelonTM Chromatography Data System. The separation was executed on a Dionex C₁₈ column (5 μ m, 120 Å, 4.6 mm× 150 mm). The mobile phase was composed of methanol (A) and water (B) with gradient elution system (0-3.5 min, 30% A isocratic; 3.5-10 min, 30-50% A; 10-30 min, 50-70% A; 30-40 min, 50% A isocratic) at a flow rate of 1.0 ml/min. The injection volume was 20 μ l. The detection UV wavelength was set at 230, 254 and 280 nm. The column temperature was maintained at 25°C.

Preparation of standard solutions and sample

Each standard stock solution was prepared by dissolving each marker components in 60% methanol at a concentration of 1 mg/ml. Each five concentrations of working solutions diluted from stock solution were used for the establishment

of calibration curve. The stock solutions were stored at 4°C. For the preparation of sample, 20 mg of commercial SHT powder was accurately weighed and dissolved in 20 ml of 60% methanol. The sample solution was filtered through a 0.45 μ m filter before HPLC injection.

RESULTS AND DISCUSSION

The chromatographic condition was optimized to separate every peak of SHT compounds with a good resolution. We chose the Dionex C_{18} column among many reverse phase columns through the preliminary test, including a XTerra RP 18 column (250 × 4.6 mm, 5µm; Waters), LUNA C_{18} column (250 × 4.6 mm, 5µm; Phenomenex). For the simultaneous determination of the five marker constituents in SHT, glycyrrhizin, paeoniflorin, 6-gingerol, cinnamic acid and decursin, gradient solvent system of water and methanol was applied as a mobile phase. The wavelength of DAD detector was tested at 230, 254, 260, and 280 nm and set at 230 nm for peoniflorin, 6-gingerol and decursin, 254 nm for glycyrrhizin and 280 nm for cinnamic acid, where the marker compounds showed the maximum absorption as measured by a DAD detector. The presence of five marker compounds in this herbal medicine was confirmed by comparing each retention time and UV spectrum with those of each standard compound, and adding authentic standards. As a result, the optimal gradient mobile phase consisting of methanol-water was subsequently employed for the analysis of SHT, which gave good resolution and satisfactory peak shape at 230, 254, and 280 nm, respectively [Figure 2].

Specificity was confirmed by the purity of peaks detected by the diode array detector. The absorption spectrum of a single component remained little variable at each time point in one peak, which supported the specificity of each peak [Figure 2]. Our results clearly showed the specificity of each peak for five marker compounds. The linearity of five compounds was calculated based on the five concentrations of each compound. The regression equation and correlation coefficients (R^2) are listed in Table 1 and high correlation coefficient values ($R^2 > 0.9903$) showed good linearity at a relatively wide range of concentration. Limit of detection (LOD) and limit of quantification (LOQ) were determined based on the method recommended by ICH (LOD = $3.3 \times (SD / slope)$ and LOQ = $10 \times (SD / slope)$ slope), SD is the standard deviation of the response, slope is the slope of calibration curve). LOD and LOQ of five marker compounds were within a range of $0.07 - 0.46 \,\mu\text{g}/$ ml and $0.13 - 1.11 \,\mu g/ml$, respectively, which showed a high sensitivity at this chromatographic condition [Table 1]. The precision test was accomplished by the intraday and interday test for these compounds. The intraday test was analyzed at three concentrations on the same day and interday test was analyzed at three concentrations on three sequential days (1, 3, 5 days). The RSD values of intraday and interday were 0.08 - 3.67% and 1.24 - 5.73%, respectively. These results indicated that this method exerted good precision [Table 2].

The recovery test was validated by the method of spiked test. The accuracy of each compound was 95.98-105.88% with RSD values less than 4.82% (n = 3) [Table 3].

The established HPLC method was applied for the simultaneous determination of the five marker components in commercial SHT sample. Analysis result suggested that this method effectively separated marker components in SHT sample without interference of peak of other components [Table 4]. Therefore, this method was very useful to evaluate the quality control and standardization of SHT.

In this study, a HPLC method was developed for the simultaneous determination of five marker components, paeoniflorin, 6-gingerol, decursin, glycyrrhzin and cinnamic



Figure 2: The HPLC chromatogram of standard mixture (A) and Ssanghwa tang sample (B)

Table 1: The linearity, correlation coefficient (R^2), limit of detection (LOD) and limit of quantification (LOQ) of the compounds studied

Linear range (µg/ml)	Regression equation ^a	R ² (n=5)	LOD (µg/ml)	LOQ (µg/ml)
0.2 - 20	Y= 0.5074 x +0.1988	0.9952	0.41	0.83
0.2 - 20	Y=0.2015 x +0.0603	0.9903	0.23	0.56
0.8 - 80	Y=0.4009 x +0.1433	0.9984	0.46	1.11
0.5 - 50	Y=0.1811 x +0.0897	0.9946	0.12	0.18
0.1 - 10	Y=0.5770 x +0.0399	0.9976	0.07	0.13
	Linear range (µg/ml) 0.2 - 20 0.2 - 20 0.8 - 80 0.5 - 50 0.1 - 10	Linear range (μg/ml)Regression equationa0.2 - 20Y= 0.5074 x +0.19880.2 - 20Y=0.2015 x +0.06030.8 - 80Y=0.4009 x +0.14330.5 - 50Y=0.1811 x +0.08970.1 - 10Y=0.5770 x +0.0399	Linear range (μg/ml)Regression equationaR² (n=5)0.2 - 20Y= 0.5074 x +0.19880.99520.2 - 20Y=0.2015 x +0.06030.99030.8 - 80Y=0.4009 x +0.14330.99840.5 - 50Y=0.1811 x +0.08970.99460.1 - 10Y=0.5770 x +0.03990.9976	Linear range (μg/ml)Regression equationa R^2 (n=5)LOD (μg/ml)0.2 - 20Y= 0.5074 x + 0.19880.99520.410.2 - 20Y=0.2015 x + 0.06030.99030.230.8 - 80Y=0.4009 x + 0.14330.99840.460.5 - 50Y=0.1811 x + 0.08970.99460.120.1 - 10Y=0.5770 x + 0.03990.99760.07

^aY: peak area, *x*:concentration (µg/ml)

Table 2: Analytical results of intra and interday test

Components	Concentration	Intraday (n=5)		Interday (n=5)	
	(µg/ml)	Mean± SD (µg/ml)	RSD (%)	Mean±SD (µg/ml)	RSD (%)
Paeoniflorin	4.00	4.84 ± 0.004	0.08	4.64 ± 0.11	2.31
	2.00	1.68 ± 0.04	2.08	1.51 ± 0.06	4.16
	0.20	0.13 ± 0.004	3.01	0.11 ± 0.01	2.48
6-Gingerol	4.00	4.81 ± 0.09	1.89	4.70 ± 0.07	1.49
-	2.00	1.56 ± 0.02	1.25	2.10 ± 0.12	5.73
	0.20	0.20 ± 0.01	3.67	0.37 ± 0.01	3.92
Decursin	7.00	6.10 ± 0.05	0.74	5.80 ± 0.23	3.93
	3.50	3.29 ± 0.02	0.75	2.80 ± 0.10	3.41
	0.35	0.35 ± 0.01	2.23	0.25 ± 0.005	1.83
Glycyrrhizin	10.00	12.13 ± 0.25	2.03	11.64 ± 0.31	2.68
	5.00	4.05 ± 0.02	0.38	4.64 ± 0.08	1.67
	0.50	0.54 ± 0.01	1.16	0.64 ± 0.03	4.43
Cinnamic acid	2.00	2.31 ± 0.01	0.23	2.24 ± 0.06	2.52
	1.00	0.96 ± 0.01	0.63	0.92 ± 0.01	1.24
	0.10	0.15 ± 0.003	1.84	0.15 ± 0.01	4.86

Table 3: Analytical results of accuracy test

•	•				
Components	Spiked amount (µg/ml)	Measured amount (µg/ml)	RSD (%)	Recovery ^a (%)	
Paeoniflorin	4.00	4.00 ± 0.03	0.48	100.01	
	1.50	1.51 ± 0.03	0.68	100.49	
	1.40	1.37 ± 0.01	0.34	97.78	
6-Gingerol	4.50	4.51 ± 1.80	1.80	100.36	
	2.25	2.38 ± 0.01	1.48	105.88	
	2.00	2.07 ± 0.01	2.03	103.73	
Decursin	7.00	6.91 ± 0.004	0.10	98.68	
	3.00	3.05 ± 0.01	2.88	95.98	
	2.00	1.98 ± 0.01	0.37	99.09	
Glycyrrhizin	1.50	1.58 ± 0.002	0.37	105.64	
	6.50	6.30 ± 0.01	0.89	96.98	
	11.00	10.62 ± 0.10	4.82	96.58	
Cinnamic acid	2.40	2.43 ± 0.01	0.31	101.09	
	1.20	1.05 ± 0.01	1.00	95.89	
	1.00	1.01 ± 0.003	0.33	101.26	

aRecovery (%) = (amount found – original amount)/amount spiked ×100 %

Table 4: Contents of five marker compounds in commercial Ssanghwa tang samples

		Content (mg/g)			
	Paeoniflorin	6-Gingerol	Decursin	Glycyrrhizin	Cinnamic acid
SHT ^a (A)	7.95 ± 0.03	0.14 ± 0.01	3.21 ± 0.14	2.54 ± 0.43	3.16 ± 0.32
SHT (B)	3.63 ± 0.11	nd ^b	0.66 ± 0.09	2.23 ± 0.22	nd
SHT (C)	5.30 ± 0.21	nd	1.65 ± 0.32	2.67 ± 0.18	2.19 ± 0.35
SHT (D)	3.82 ± 0.14	nd	1.44 ± 0.10	2.88 ± 0.12	3.05 ± 0.21
SHT (E)	5.52 ± 0.21	nd	0.98 ± 0.14	2.53 ± 0.21	2.89 ± 0.38

^aSHT: commercial Ssanghwa tang samples made by five different pharmaceutical companies, ^bnd: not detected

acid in SHT. Validation of the method was carried out with linearity, accuracy and precision test. The results of validation indicated good precision and accuracy. The results of analysis of the commercial product suggest that this analysis method can be successfully applied for the quantification of marker compounds in SHT.

ACKNOWLEDGEMENTS

This research was supported by a grant [K09040] from the Korea Institute of Oriental Medicine.

REFERENCES

- 1. Normile D. Asian medicine. The new face of traditional Chinese medicine. Science 2003;299:188-90.
- 2. Xue T, Roy R. Studying traditional Chinese medicine. Science 2003;300:740-1.
- Lund ST, Bohlmann J. The molecular basis for wine grape quality: A volatile subject. Science 2006;311:804-5.
- 4. Lee MK, Park JH, Cho JH, Kim DH, Baek JH, Kim HJ, et al.

Simultaneous determination of hesperidin and glycyrrhizin in Pyungwi-san by HPLC/DAD. Kor J Pharmacogn 2008;39: 199-202.

- Lee MK, Lee KY, Kim SH, Park J, Cho JH, Oh MH, et al. Simultaneous determination of baicalin and glycyrrhizin in Eul-Ja-Tang by HPLC/DAD. Nat Prod Sci 2008;14:147-51.
- Ye J, Zhang X, Dai W, Yan S, Huang H, Liang X, et al. Chemical fingerprinting of Liuwei Dihuang Pill and simultaneous determination of its major bioactive constituents by HPLC coupled with multiple detections of DAD, ELSD and ESI-MS. J Pharm Biomed Anal 2009;49:638-45.
- Zhang H, Chen S, Qin F, Huang X, Ren P, Gu X. Simultaneous determination of 12 chemical constituents in the traditional Chinese Medicinal Prescription Xiao-Yao-San-Jia-Wei by HPLC coupled with photodiode array detection. J Pharm Biomed Anal 2008;48:1462-6.
- Song J, Han Q, Qiao C, Yip Y, Xu H. Simultaneous determination of multiple marker constituents in concentrated Gegen Tang granule by high performance liquid chromatography. Chin Med 2007;20:2-7.
- Ma JY, Park DH, Park KS, Do KT, Shin HK. Acute toxicity study on Ssanghwa tang in mice. Korean J Ori Med 2007;13:141-9.

Source of Support: Grant [K09040] from the Korea Institute of Oriental Medicine, Conflict of Interest: None declared