

# Toxicity evaluation of *Xanthorrhoea johnsonii* leaf methanolic extract using the *Artemia franciscana* bioassay

I. E. Cock<sup>1,2</sup>, F. R. Kalt<sup>1</sup>

<sup>1</sup>Biomolecular and Physical Sciences, Nathan Campus, Griffith University, 170 Kessels Road, Nathan, Queensland 4111, <sup>2</sup>Genomics Research Centre, Gold Coast Campus, Griffith University, Parklands Drive, Southport, Queensland 4222, Australia

Submitted: 13-01-2010

Revised: 05-02-2010

Published: 30-07-2010

## ABSTRACT

The toxicity of *Xanthorrhoea johnsonii* methanolic leaf extract was investigated using the *Artemia franciscana* nauplii bioassay. At 24 h, the extract produced an EC<sub>50</sub> of 361.0 ± 41.8 µg/ml, indicating that it was substantially more toxic than the pesticide Mevinphos (1346.2 ± 80.1 µg/ml) and approximately 4 fold less toxic than potassium dichromate (87.1 ± 5.2 µg/ml). Whilst potassium dichromate LC<sub>50</sub> values remained constant across the 72-h test period, these values decreased for the extract and Mevinphos to similar values (199.8 ± 60.5 and 114 ± 12.8 µg/ml, respectively), indicating their similar level of efficacy. Noteworthy was the apparent anesthetic effect of *X. johnsonii* leaf extract. Although the extract initially appeared to kill the *A. franciscana* nauplii, they were seen to temporarily recover by 48 h only to die by 72 h.

**Key words:** Australian plants, grass tree, methanol extracts, toxic plants, *Xanthorrhoea johnsonii*

## INTRODUCTION

Plants produce a wide variety of compounds, which in addition to giving them characteristic pigment, odor and flavor characteristics, may also have medicinal and therapeutic properties.<sup>[1]</sup> For thousands of years, traditional plant-derived medicines have been used in most parts of the world and their use in fighting disease is becoming the focus of intense study.<sup>[2,3]</sup> Much of the research into traditional medicinal plant use has focused on Asian,<sup>[4]</sup> African<sup>[5]</sup> and South American<sup>[6]</sup> plants. Relatively few studies have focused on the medicinal properties of Australian native plants.

*Xanthorrhoea* (grasstree) is a small genus of evergreen plants consisting of 28 species and 5 subspecies, all of which are endemic to Australia.<sup>[7]</sup> All species are long-lived with some plants being estimated at more than 550 years of age.<sup>[7-9]</sup> However, *Xanthorrhoea* species are also extremely slow growing. Indeed, the growth of *Xanthorrhoea johnsonii*

has been estimated to be at a rate of 0.88 cm/year.<sup>[9,10]</sup> Some species are stemless whilst others develop a thick, sometimes branching, trunk topped by a dense crown of rigid leaves. In species with a trunk, this trunk may take up to 20 years to develop.<sup>[11]</sup>

Australian Aboriginal people had many traditional uses for *Xanthorrhoea* species including use as tools and weapons<sup>[12-14]</sup> and as a resin.<sup>[12,15]</sup> *Xanthorrhoea* species also provided a minor food source for Aboriginals. Tubers<sup>[13]</sup> and shoots<sup>[12]</sup> were eaten whilst nectar from the flowers was used to make a sweet drink.<sup>[13]</sup> The seeds could also be ground to make dough. Since European settlement, *Xanthorrhoea* species were extensively harvested for their resin<sup>[7]</sup> until they were declared protected species.

Being such a slow growing group of plants, it is likely that *Xanthorrhoea* species have developed protective mechanisms to deter animal foraging which could potentially threaten their survival. Indeed, the list of animals that use *Xanthorrhoea* species as a major food source is relatively small.<sup>[7]</sup> Whilst several species of insects, birds and mammals are known to use *Xanthorrhoea* for food, they almost exclusively feed on the flowers and seeds, leaving all but the youngest leaves largely untouched. Indeed, *X. johnsonii* has been reported to poison cattle that graze on

### Address for correspondence:

Dr. Ian Cock, Biomolecular and Biomedical Sciences, Griffith University, Nathan Campus, 170 Kessels Road, Nathan, Queensland 4111, Australia. E-mail: I.Cock@griffith.edu.au

DOI: 10.4103/0973-1296.66929

its leaves.<sup>[16-19]</sup> Cattle poisoned by *Xanthorrhoea* become uncoordinated and lose condition, become dehydrated, and in severe cases die. It is possible that a chemical protectant/deterrent produced by *Xanthorrhoea* species discourages foraging by herbivores, allowing juvenile plants to reach maturity.

This study was undertaken to examine the toxicity of *X. johnsonii* leaves using the *A. franciscana* nauplii (brine shrimp larvae) lethality bioassay. This assay has been used to examine the toxicity of a wide variety of compounds.<sup>[20]</sup> It is an efficient, inexpensive and a relatively rapid way to detect toxic compounds, requiring only low amounts of sample (<20 mg). This test correlates well with cytotoxic activity of some human tumors and, therefore, has the potential to detect new antitumour agents.<sup>[21]</sup>

## MATERIALS AND METHODS

### Plant collection and extraction

*X. johnsonii* is one of the two species of *Xanthorrhoea* occurring naturally in Toohey Forest, Brisbane, Australia (along with *Xanthorrhoea macronema*) and is by far the dominant species. *X. johnsonii* leaves were collected from Toohey Forest and were identified with reference to a taxonomic key to Toohey Forest plants.<sup>[22]</sup> Samples were dried in a Sunbeam food dehydrator and then ground to a coarse powder. One gram of the powdered leaves was extracted extensively in 50 ml methanol (Ajax, AR grade) for 24 h at 4°C with gentle shaking. The extract was passed through filter paper (Whatman No. 54) under vacuum, followed by drying by rotary evaporation in an Eppendorf concentrator 5301. The resultant pellet was dissolved in 10 ml of 20% methanol. The extract was passed through 0.22 µm filter (Sarstedt) and stored at 4°C.

### Reference toxins for biological screening

Potassium dichromate (K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>) (AR grade, Chem-Supply, Australia) was prepared as a 1.6 mg/ml solution in distilled water and was serially diluted in synthetic seawater (described below) for use in the *A. franciscana* nauplii bioassay. Mevinphos (2-methoxycarbonyl-1-methylvinyl dimethyl phosphate) was obtained from Sigma-Aldrich as a mixture of *cis* (76.6%) and *trans* (23.0%) isomers and prepared as a 4 mg/ml stock in distilled water. The stock was serially diluted in artificial seawater for use in the bioassay.


### *A. franciscana* nauplii toxicity screening

Toxicity was tested using the *A. franciscana* nauplii lethality assay developed by Meyer *et al.*<sup>[20]</sup> for the screening of active plant constituents, with the following modifications.

*A. franciscana* cysts were obtained from North American Brine Shrimp, LLC, USA (harvested from the Great Salt Lake, Utah). Synthetic seawater was prepared using Reef Salt, AZOO Co., USA. Seawater solutions at 34 g/l distilled water were prepared prior to use. Two grams of *A. franciscana* cysts were incubated in 1 l synthetic seawater under artificial light (2000 Lux) at 25 °C with continuous aeration. Hatching commenced within 16–18 h of incubation. Newly hatched *A. franciscana* (nauplii) were used within 10 h of hatching. Nauplii were separated from the shells and remaining cysts and were concentrated to a suitable density by placing an artificial light at one end of their incubation vessel and the nauplii-rich water closest to the light was removed for biological assays. Four hundred microliters of seawater containing approximately 50 (mean 53, *n* = 168, SD 12) nauplii was added to wells of a 48-well plate and immediately used for bioassay. The plant extract was diluted to 2 mg/ml in seawater for toxicity testing, resulting in a 1 mg/ml concentration in the bioassay. Four hundred microliters of diluted plant extract and the reference toxins were transferred to the wells and incubated at 25 ± 1 °C under artificial light (1000 Lux). A negative control (400 µl seawater) was run in at least triplicate for each plate. All the treatments were performed in at least triplicate. The wells were checked at regular intervals and the number of dead counted. The nauplii were considered moribund if no movement of the appendages was observed within 10 s. After 72 h, all nauplii were sacrificed and counted to determine the total number per well. The LC<sub>50</sub> with 95% confidence limits for each treatment was calculated using probit analysis.<sup>[23]</sup>

## RESULTS AND DISCUSSION

One gram of powdered dried *X. johnsonii* leaves was extensively extracted with methanol and dried under vacuum, resulting in 39 mg of dried extracted material. Resuspension of the dried fraction in 10 ml of 10% methanol resulted in the crude test extract concentration of 3.9 mg/ml. The extract was diluted to 2000 µg/ml in artificial seawater for toxicity testing, resulting in a 1000 µg/ml concentration in the *A. franciscana* lethality bioassay. *A. franciscana* nauplii were also tested against methanol dilutions to determine the effect of residual methanol on toxicity. No increase in mortality above the control values was seen when testing methanol concentrations up to 5% in the assay, indicating that the residual methanol in the re-suspended extracts was not responsible for nauplii mortality (unpublished results). The results of *A. franciscana* bioassay screening of the *X. johnsonii* methanolic extract and the control toxins are shown in Figure 1.




**Figure 1:** Brine shrimp lethality of (a) *X. johnsonii* extract (1000 µg/ml), (b) Mevinphos (2000 µg/ml), (c) potassium dichromate (800 µg/ml). All bioassays were performed in at least triplicate and are expressed as mean  $\pm$  standard deviation

Two interesting features of *X. johnsonii* toxicity were noted. Firstly, *X. johnsonii* extract induced apparent moribundity much more rapidly than was observed for either of the positive control toxins. High levels of apparent moribundity ( $87.6 \pm 5.8\%$ ) were seen with *X. johnsonii* extract after 12-h exposure, with 100% mortality seen by 24 h. In contrast, both Mevinphos and potassium dichromate took much longer to achieve high levels of mortality. Following 12 h exposure, both Mevinphos and potassium dichromate had only induced low levels of mortality (0 and  $21.3 \pm 6.9\%$ , respectively). Indeed, 36 h was required by Mevinphos to achieve approximately 100% mortality ( $96.9 \pm 3.3\%$ ).

Perhaps more interesting was the decrease in apparent moribundity seen between 36 and 60 h of exposure. Throughout the course of these studies and previous studies within this laboratory, lack of movement of *A. franciscana* appendages for 10 s was considered as an indication of apparent moribundity. Many other studies have been less rigorous in their definition of apparent moribundity when determining  $LC_{50}/EC_{50}$  values. Several studies have defined death (for  $LC_{50}$  determination) as the lack of controlled forward movement. As this could indicate intoxication from which the brine shrimp could recover, we have defined a total lack of movement, over a period of 10 s, as our endpoint. Surprisingly, many of the *A. franciscana* nauplii used in these studies recovered from their apparent “death” and were seen to swim normally by 48 h exposure. These studies have been repeated several times using a longer time period (10 min) for lack of movement of appendages as a definition of death and the same trend is evident. It appears that the *X. johnsonii* extract has an anesthetic affect similar to the effect previously reported for curare,<sup>[24-26]</sup> from which the *A. franciscana* nauplii are able to temporarily recover. However, at 60 h, it became evident that the nauplii were experiencing difficulty in swimming (“strobe-like” swimming). Following 60 h, the percentage apparent moribundity again began to increase. The nauplii did not recover from this moribund state a second time.

A concentration–response analysis was undertaken to determine the dependence of percentage apparent moribundity on *X. johnsonii* extract concentration. Due to the previously described decrease in moribundity with time, we report both on the percentage dead (as defined by lack of movement of appendages for 10 s) [Figure 2] and percentage affected (lack of controlled forward movement) [Figure 3]. The same trend previously described for 1000 µg/ml *X. johnsonii* extract was also apparent in these concentration–response studies. Apparent moribundity of *A. franciscana* exposed to 1000 µg/ml *X. johnsonii* extract again decreased from 100% moribund at 24 h to 76.7% ( $\pm 3.6$ ) by 48 h and increased again to 87.9% ( $\pm 8.5$ ) by 72 h. The trend was even more apparent when percentage affected was monitored [Figure 3]. The percentage of affected *A. franciscana* exposed to 1000 µg/ml *X. johnsonii* extract decreased from 100% affected at 24 h to 63.2% ( $\pm 3.9$ ) by 48 h and increased again to 93.4% ( $\pm 6.6$ ) by 72 h.

$LC_{50}$  values for the *X. johnsonii* extract, potassium dichromate and mevinphos were calculated by probit




**Figure 2:** Dependence of *X. johnsonii* extract concentration on *A. franciscana* apparent moribundity following exposure for (a) 24 h, (b) 48 h, (c) 72 h. *X. johnsonii* extract was diluted freshly prior to use in deionized water; 400  $\mu$ l of juice dilution was added to 400  $\mu$ l of saline containing *A. franciscana nauplii*. Moribundity was defined as the lack of appendage movement for at least 10 s. All the bioassays were performed in at least triplicate

analysis [Table 1]. Similarly,  $EC_{50}$  values for *X. johnsonii* extract were also calculated. Both the extract and the control toxins displayed significant correlation between the concentration/dilution and the percentage affected. The  $R^2$  coefficients for the lines of regression of each test sample were greater than 0.9 with the response linear and statistically significant at  $P < 0.05$ .

## CONCLUSIONS

The results reported here demonstrate the toxicity of



**Figure 3:** Dependence of *X. johnsonii* extract concentration on percentage of affected *A. franciscana*, following exposure for (a) 24 h, (b) 48 h, (c) 72 h. *X. johnsonii* extract was diluted freshly prior to use in deionized water; 400  $\mu$ l of juice dilution was added to 400  $\mu$ l of saline containing *A. franciscana nauplii*. Affected nauplii were defined as the lack of controlled forward movement. All bioassays were performed in at least triplicate

methanolic extracts of *X. johnsonii* leaves, providing a possible explanation for the lack of use of the leaves as a food source for Australian native herbivores. Moreover, as the *A. franciscana nauplii* bioassay used in these studies correlates well with antitumor activity,<sup>[21]</sup> it is possible that *X. johnsonii* leaf extracts may also be effective as anti-tumor agents. Further studies are needed to determine whether this is the case and to identify the factors involved and their mechanism of action.

Perhaps more interesting was the apparent anesthetic

**Table 1: LC<sub>50</sub> and EC<sub>50</sub> determinations of *X. johnsonii* extract and the reference toxins Mevinphos and potassium dichromate; values indicate the mean of triplicate determinations ± standard deviation**

Treatment	LC <sub>50</sub> value in µg/ml (95% confidence interval) at time (h)		
	24 h	48 h	72 h
<i>X. johnsonii</i> extract	361.0 ± 41.8	402.1 ± 15.7	199.8 ± 60.5
<i>X. johnsonii</i> extract*	220.1 ± 8.6*	250.0 ± 9.8*	187.5 ± 19.8*
Mevinphos	1346.2 ± 80.1	523.3 ± 39.0	114 ± 12.8
Potassium dichromate	87.1 ± 5.2	81.9 ± 3.8	79.6 ± 4.6

\*Indicates EC<sub>50</sub> determinations (where effect is defined as lack of controlled forward movement)

effect of the methanol extract on the *A. franciscana* nauplii. This effect is similar to the effects previously described for tubocurarine, dimethyltubocurarine and alcuronium (collectively known as curare, a South American arrow poison) from *Chondrodendron tomentosum*.<sup>[24-26]</sup> Curare is a neuromuscular blocker that was commonly used as a muscle relaxant by anesthesiologists prior to the development of safer, more effective muscle relaxants. Curare functions by blocking neuronal nicotinic acetylcholine receptors (nAChR),<sup>[27]</sup> thus blocking neurotransmission. Determining whether the *X. johnsonii* leaf methanolic extract functions by a similar mechanism or not, was outside the scope of this study. Further studies are needed to isolate individual components from *X. johnsonii* leaf and examine their mechanism/s of action. It is possible that novel anesthetics/muscle relaxants may be discovered by such studies.

## ACKNOWLEDGMENTS

Financial support for this work was provided by the School of Biomolecular and Physical Sciences, Griffith University, Australia. The authors would like to thank Ella Merriman for assistance with collection, identification and processing of plant specimens.

## REFERENCES

- Cowan MM. Plant products as antibacterial agents. *Clin Microbiol Rev* 1999;12:564-82.
- Bhavnani SM, Ballow CH. New agents for Gram-positive bacteria. *Curr Opin Microbiol* 2000;3:528-34.
- Chiariandy CM, Seaforth CE, Phelps RH, Pollard GV, Khambay BP. Screening of medicinal plants from Trinidad and Tobago for antimicrobial and insecticidal properties. *J Ethnopharmacol* 1999;64:265-70.
- Patwardhan B, Warude D, Pushpangadan P, Bhatt N. Ayurveda and traditional Chinese medicine: A comparative overview. *Evid Based Complement Alternat Med* 2005;2:465-73.
- Hostettmann K, Marston A, Ndjoko K, Wolfender J. The potential of African plants as a source of drugs. *Curr Org Chem* 2000;4:973-1010.
- Alonso Paz E, Cerdeiras MP, Fernandez J, Ferreira F, Moyna P, Sobes M, *et al.* Screening of Uruguayan medicinal plants for antimicrobial activity. *J Ethnopharmacol* 1995;45:67-70.
- Borsboom AC. *Xanthorrhoea*: A review of current knowledge with a focus on *X. johnsonii* and *X. latifolia*, two Queensland protected plants-in-trade. Queensland: Environmental Protection Agency; 2005.
- Stanley TD, Ross EM. *Flora of South-eastern Queensland*. Vol. 3. Brisbane: Miscellaneous Publication QM88001, Queensland Department of Primary Industries; 1989. p. 45.
- Bulow-Olsen A, Just J, Liddle MJ. Growth and flowering history of *Xanthorrhoea johnsonii* Lee (Liliaceae) in Toohy Forest Queensland. *Bot J Linn Soc* 1982;84:195-207.
- Lamont BB, Downes S. The longevity, flowering and fire history of the grasstrees *Xanthorrhoea preissii* and *Kingia australis*. *J Appl Ecol* 1979;16:893-9.
- Burnie G, Forrester S, Greig D, Guest S, Harmony M, *et al.* *Botanica: The Illustrated A - Z of Over 10,000 Garden Plants for Australian Gardens and How to Cultivate Them*. In: Page S, Olds M, editors. Australia: Random House; 2004.
- Anderson ER. *Plants of Central Queensland: Their Identification and uses*. Information Series QI 92037. Queensland: Department of Primary Industries; 1993. p. 119.
- Gott B. Use of Victorian plants by Koories. In: Foreman DB, Walsh NG, editors. *Flora of Victoria*. Vol. 1. Melbourne: Inkarta Press; 1993. p. 195-211.
- Bedford DJ. Australian grasstrees. *Aus Nat Hist* 1992;24:42-50.
- Staff IA, Waterhouse JT. Arborescent monocotyledons. In: Pate JS, McComb AJ, editors. *The Biology of Australian Plants*. Nedlands: University of Western Australia Press; 1981. p. 238-57.
- McKenzie R. Australian native poisonous plants, Australian Society for Growing Australian Plants (Australia) Available from: <http://farrer.csu.edu.au/ASGAP?Apol7/sep97-4.html> [last updated on 1997] [last accessed on 2010 Jan 12].
- Everist SL. Botanical affinities of Australian poisonous plants. In: Keeler RF, Van Kampen KR, Lynn LJ, editors. *Effects of Poisonous Plants on Livestock*. London: Academic Press; 1978. p. 93-100.
- Hall WT. *Xanthorrhoea hastile* poisoning of cattle. *Queens J Agric Sci* 1956;13:1-10.
- Hurst E. *The poison plants of New South Wales*. Sydney: Snelling Printing Works; 1942.
- Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL. Brine shrimp: A convenient general bioassay for active plant constituents. *Planta Med* 1982;45:31-4.
- McLaughlin JL, Rogers LL, Anderson JE. The use of biological assays to evaluate botanicals. *Drug Inf J* 1998;32:513-24.
- Coutts RH, Catterall CP. *Identifying the plants of Toohy Forest*. Nambour, Australia: Ecos Educational Publishers; 1908.
- Finney DJ. *Probit Analysis*. 3<sup>rd</sup> ed. Cambridge: Cambridge University Press; 1971.
- Bisset NG. Arrow and dart poisons. *J Ethnopharmacol* 1989;25:1-41.
- Bisset NG. Curare. In: Pelletier WS, editor. *Alkaloids: Chemical*

and Biological Perspectives. Vol. 8. Berlin: Springer; 1992. p. 3-150.

26. Bisset NG. War and hunting poisons of the New World: Part 1: Notes on the early history of curare. J Ethnopharmacol 1992;36:1-26.
27. Strecker GJ, Jackson MB. Curare binding and the curare-induced subconductance state of the acetylcholine receptor

channel. Biophys J 1989;56:795-806.

**Source of Support:** School of Biomolecular and Physical Sciences, Griffith University, Australia,  
**Conflict of Interest:** None declared

### Author Help: Reference checking facility

The manuscript system ([www.journalonweb.com](http://www.journalonweb.com)) allows the authors to check and verify the accuracy and style of references. The tool checks the references with PubMed as per a predefined style. Authors are encouraged to use this facility, before submitting articles to the journal.

- The style as well as bibliographic elements should be 100% accurate, to help get the references verified from the system. Even a single spelling error or addition of issue number/month of publication will lead to an error when verifying the reference.
- Example of a correct style  
Sheahan P, O'leary G, Lee G, Fitzgibbon J. Cystic cervical metastases: Incidence and diagnosis using fine needle aspiration biopsy. Otolaryngol Head Neck Surg 2002;127:294-8.
- Only the references from journals indexed in PubMed will be checked.
- Enter each reference in new line, without a serial number.
- Add up to a maximum of 15 references at a time.
- If the reference is correct for its bibliographic elements and punctuations, it will be shown as CORRECT and a link to the correct article in PubMed will be given.
- If any of the bibliographic elements are missing, incorrect or extra (such as issue number), it will be shown as INCORRECT and link to possible articles in PubMed will be given.