

Antimicrobial and antifungal properties of the essential oil and methanol extracts of *Eucalyptus largiflorens* and *Eucalyptus intertexta*

Javad Safaei-Ghomi, Atefeh Abbasi Ahd

Essential Oils Research Institute, University of Kashan, Kashan, I. R. Iran

Submitted: 16-02-2010

Revised: 07-03-2010

Published: 30-07-2010

ABSTRACT

This study was conducted to evaluate the *in vitro* antimicrobial properties of essential oil, its major component, 1,8-cineole, and extracts of two *Eucalyptus* species, *Eucalyptus intertexta* and *Eucalyptus largiflorens*. Minimum inhibitory concentration (MIC) of the extracts was calculated by broth dilution method and the zone of inhibition was studied by agar disk diffusion method. Gentamicin (10 µg/disk) and rifampin (5 µg/disk) were used as reference controls for antibacterial studies and nystatin (100 µg/disk) for antifungal studies. The results of MIC study revealed that the essential oil has a stronger activity and broader spectrum than those of methanol extracts. It is interesting to point out that the oils had even greater potential of antimicrobial activities than those of 1,8-cineole as their main component.

Key words: Antimicrobial activity, *Eucalyptus largiflorens*, *Eucalyptus intertexta*, minimum inhibitory concentration, zone of inhibition

INTRODUCTION

The genus *Eucalyptus* is known for its rich source of bioactive compounds.^[1] It is a source for several unique secondary metabolites, which show a variety of biological activities, such as those of antioxidants, antibacterials, HIV inhibitors, attachment inhibitors, and others.^[2-9] Although reports on the essential oil composition of different *Eucalyptus* species are relatively common,^[10-15] investigations on their biological activities are still scarce. *Eucalyptus intertexta* and *Eucalyptus largiflorens* are two cultivated and adapted *Eucalyptus* species in warm regions of Iran, Kashan. To the best of our knowledge, the chemical composition of their essential oil is previously reported,^[16-18] but there is no report on antimicrobial profiles of these two species. Thus, in this study, the *in vitro* antimicrobial activities of their essential oil, its main component and extracts, were evaluated against a set of 11 microorganisms. Their activity potentials were qualitatively and quantitatively assessed by the presence or absence of inhibition zones, zone diameters and minimum inhibitory concentration (MIC) values.

Address for correspondence:

Dr. Javad Safaei-Ghomi, Essential Oils Research Institute,
University of Kashan, Kashan, I. R. Iran.
E-mail: safaei@kashanu.ac.ir

DOI: 10.4103/0973-1296.66930

MATERIALS AND METHODS

Plant material

Leaves of *E. largiflorens* and *E. intertexta* were collected from cultivated sample in Kashan Botanical Garden (Isfahan Province, Iran) at an altitude of ca. 1000 m in December 2006. The voucher specimen of the plant has been placed in the herbarium of Kashan Research Botanical Garden, Research Institute of Forests and Rangelands, Kashan, Iran.

Drugs and chemicals

1,8 Cineol was obtained from Sigma–Aldrich Chemie (Steinheim, Germany). Gentamicin, rifampin and nystatin were purchased from Himedia (Mumbai, India). Analytical grade methanol, dimethyl sulfoxide (DMSO), HPLC grade chloroform, anhydrous sodium sulfate, tween 40, and all culture media were obtained from Merck (Darmstadt, Germany). Ultrapure water was used for the experiments.

Isolation of the essential oil

A portion (100 g) of dried and finally ground plant material was subjected, for 3.5 h, to water distillation using a Clevenger-type apparatus as recommended by European Pharmacopoeia.^[19] The obtained essential oil was dried over anhydrous sodium sulfate and, after filtration, stored in amber vial at 4°C until analysis. The yield based on dry weight of the sample was calculated.

Preparation of methanol extracts

A portion (20 g) of the powdered plant material was Soxhlet-extracted with methanol for 8 h, at a temperature not exceeding the boiling point of the solvent. The extracts were concentrated using a rotary evaporator at 50°C to get crude extracts. Dried extracts were suspended in water and partitioned with chloroform to obtain polar (MW) and non-polar (MC) fractions. All the extracts were dried and kept in the dark at 4°C prior to use.

Antimicrobial activity

Microorganisms

The essential oil was tested against 11 microorganisms including *Aspergillus niger* ATCC 16404, *Candida albicans* ATCC 10231, *Pseudomonas aeruginosa* ATCC 27853, *Bacillus subtilis* ATCC 6633, *Staphylococcus aureus* ATCC 29737, *Escherichia coli* ATCC 10536, *Klebsiella pneumoniae* ATCC 10031, *Staphylococcus epidermidis* ATCC 12228, *Shigella dysenteriae* PTCC 1188, *Proteus vulgaris* PTCC 1182 and *Salmonella paratyphi-A* serotype ATCC 5702. All were provided by Iranian Research Organization for Science and Technology (IROST). Bacterial strains were cultured overnight at 37°C in nutrient agar (NA) and fungi were cultured overnight at 30°C in Sabouraud dextrose agar (SDA).

Disk diffusion assay

The *in vitro* antimicrobial activity of samples was evaluated by the disk diffusion method (NCCLS).^[20] The dried plant extracts were dissolved in DMSO to a final concentration of 30 mg/mL and filtered using 0.45 µm millipore filters for sterilization. Antimicrobial tests were carried out using the disk diffusion method^[21] and employing 100 µL of suspension containing 10⁸ CFU/mL of bacteria, 10⁶ CFU/mL of yeast and 10⁴ spore/mL of fungi spread on the NA, SDA and potato dextrose (PD) agar mediums, respectively. The disks (6 mm in diameter) impregnated with 10 µL of the essential oil, a commercial sample of 1,8-cineole or the extract solutions (300 µg/disk) and DMSO (as negative control) were placed on the inoculated agar. The inoculated plates were incubated for 24 h at 37°C for bacterial strains and for 48 and 72 h at 30°C for yeast and mold isolates, respectively. Gentamicin (10 µg/disk) and rifampin (5 µg/disk) were used as positive controls for bacteria and nystatin (100 IU) for fungi. The diameters of inhibition zones were used as a measure of antimicrobial activity and each assay was repeated two times.

Microwell dilution assay

MIC values were measured by microwell dilution assay method.^[22] The inocula of the microbial strains were prepared from 12 h broth cultures and suspensions were adjusted to 0.5 McFarland standard turbidity. The samples were dissolved in 10% DMSO and diluted to

the highest concentration (500 µg/mL) to be tested, and then serial twofold dilutions were made to a concentration ranging from 7.8 to 500 µg/mL in 10 mL sterile test tubes containing brain heart infusion (BHI) broth for bacterial strains and Sabouraud dextrose (SD) broth for yeast. The 96-well plates were prepared by dispensing 95 µL of the culture media and 5 µL of the inoculum into each well. A 100-µL aliquot from the stock solutions of the plant extracts initially prepared at the concentration of 500 µg/mL was added into the first well. Then, 100 µL from their serial dilutions was transferred into six consecutive wells. The last well containing 195 µL of the culture media without the test materials and 5 µL of the inoculum on each strip was used as the negative control. The final volume in each well was 200 µL. Gentamicin and rifampin for bacteria and nystatin for yeast were used as standard drugs for positive control in conditions identical to test materials. The plates were covered with sterile plate sealers. Contents of each well were mixed on plate shaker at 300 rpm for 20 s and then incubated at appropriate temperatures for 24 h. Microbial growth was determined by the presence of a white pellet on the well bottom and confirmed by plating 5 µL samples from clear wells on NA medium. The MIC value was defined as the lowest concentration of the plant extracts required for inhibiting the growth of microorganisms. All tests were performed in duplicate.

Minimum inhibitory concentration agar dilution assay

MIC values of 1,8-cineole for the fungus isolate sensitive to it were evaluated based on the agar dilution method.^[23] Appropriate amount of this compound was added aseptically to sterile, molten SDA medium containing Tween 20 (0.5%, v/v) to produce a concentration range of 7.8–500 µg/mL. The resulting SDA agar solutions were immediately mixed and poured into petri plates. The plates were spot inoculated with 5 µL (10⁴ spore/mL) of fungus isolate. Nystatin was used as reference antifungal drug and the inoculated plates were incubated at 30°C for 72 h. At the end of the incubation period, the plates were evaluated for the presence or absence of growth. The MIC was defined as the lowest concentration of the compound needed to inhibit the growth of microorganisms. Each test was repeated at least twice.

RESULTS AND DISCUSSION

Hydrodistillation of aerial parts of *E. largiflorens* and *E. intertexta* yields, respectively, 1.85 and 1.5% (w/w) of light yellowish oil.

According to the results given in Table 1, the essential oil of *E. intertexta* had great potential of antimicrobial activities

Table 1: Antimicrobial activity of the essential oil, its major components (1,8-cineole) and methanol fractions of *E. intertexta* and *E. largiflorens*

| Test microorganisms | Essential oil | | | | Main component | | | | Extract | | | | | | | | | | | | | | | |
|---------------------------------|----------------------|------|------------------------|------|----------------|------|----|------|----------------------|------|------|------------------------|----|------|-------------|------|----|-----|----|------|-----|-----|------|----|
| | <i>E. intertexta</i> | | <i>E. largiflorens</i> | | 1,8-cineole | | | | <i>E. intertexta</i> | | | <i>E. largiflorens</i> | | | Antibiotics | | | | | | | | | |
| | DDa | MICb | DD | MIC | DD | MIC | DD | MIC | CHCl3 | H2O | DD | MIC | DD | MIC | DD | MIC | DD | MIC | DD | MIC | | | | |
| <i>Ps. aeruginosa</i> | — | — | — | — | — | — | — | — | — | 10 | 31.2 | — | — | — | 10 | 31.2 | — | — | — | 23 | 500 | NA | NA | |
| <i>B. subtilis</i> | 11 | 15.6 | 28 | 125 | 30 | 250 | — | — | — | — | — | — | — | — | — | — | — | — | 13 | 15.6 | 21 | 500 | NA | NA |
| <i>E. coli</i> | 10 | 15.6 | — | — | 20 | 500 | — | — | — | — | — | — | — | — | — | — | — | — | 11 | 500 | 20 | 500 | NA | NA |
| <i>St. aureus</i> | 11 | 7.8 | 12 | 250 | 18 | 500 | — | — | — | — | — | — | — | — | — | — | — | — | 10 | 250 | 21 | 500 | NA | NA |
| <i>K. pneumoniae</i> | 15 | 7.8 | 20 | 125 | 20 | 500 | — | — | — | — | — | — | — | — | — | — | — | — | 7 | 250 | 22 | 250 | NA | NA |
| <i>St. epidermidis</i> | 11 | 7.8 | 15 | 125 | 22 | 250 | 17 | 62.5 | 15 | 31.2 | — | — | — | — | 16 | 31.2 | 40 | 250 | 35 | 500 | 35 | 500 | NA | NA |
| <i>Sh. dysenteriae</i> | 14 | 7.8 | — | — | 15 | 250 | — | — | — | — | — | — | — | — | 12 | 7.8 | 8 | 250 | 18 | 500 | 18 | 500 | NA | NA |
| <i>Pr. vulgaris</i> | 12 | 7.8 | 11 | 7.8 | 11 | 62.5 | — | — | — | — | — | — | — | — | 10 | 125 | 10 | 125 | 10 | 125 | 23 | 500 | NA | NA |
| <i>Sa. paratyphi-A</i> serotype | — | — | 10 | 62.5 | 12 | 500 | — | — | — | — | — | — | — | — | 10 | 62.5 | — | — | — | — | 21 | 500 | NA | NA |
| <i>C. albicans</i> | 12 | 7.8 | 31 | 125 | 11 | 31.3 | 16 | 62.5 | 17 | 62.5 | 14 | 125 | 11 | 31.2 | NA | NA | NA | NA | NA | NA | NA | 33 | 125 | |
| <i>A. niger</i> | — | — | 22 | 125 | — | — | — | — | 12 | 500 | 12 | 500 | — | — | NA | NA | NA | NA | NA | NA | NA | 27 | 31.3 | |

— indicates no antimicrobial activity, *DD (disk diffusion method), inhibition zones in diameter (mm) around the impregnated disks, ^bMIC (minimal inhibition concentrations as µg/ml), NA (not applicable)

against seven of the nine bacteria and a yeast species tested. Non-polar and polar fractions of the methanol extract were also found to be effective against one and four bacterial species, respectively, of the nine examined. However, essential oil and non-polar extract failed to show antifungal activity. The maximal inhibition zones and MIC values for bacterial strains which were sensitive to the essential oil, and non-polar and polar extracts of *E. intertexta*, were in the range of 10–12 mm and 7.8–15.6 µg /mL, 16–17 mm and 62.5 µg/mL, and 10–17 mm and 31.2–500 µg /mL, respectively.

The results of the bioassay [Table 1] also revealed that the essential oil of *E. largiflorens* exhibited moderate to high antimicrobial activity against all the bacteria, yeast and mold tested, except three microorganisms, *Ps. aeruginosa*, *E. coli* and *Sb. dysenteriae*. The evaluation of methanol fraction indicated that polar fraction showed strong activity against 7 out of 11 microorganisms while non-polar fractions did not possess any inhibitory action against the strains evaluated except *E. coli*.

Based on these results, it is possible to conclude that the essential oil has a stronger activity and broader spectrum than those of methanol extracts.

The relatively high antimicrobial activities of essential oils are most likely due to the presence of compounds with antimicrobial properties. A number of compounds present in relatively high concentrations in the essential oils are known to have antimicrobial properties. Particularly worth noting is 1,8-cineole (eucalyptol), which accounted for approximately 70.2% (v/v) of the *E. intertexta*^[17] and 37.5% (v/v) of the *E. largiflorens* essential oil,^[16] and which has been found to possess relatively strong antimicrobial properties against many important pathogens and spoilage organisms.^[24-26] These reports are further supported by our finding about 1,8-cineole which showed high inhibitory activities against *C. albicans* and *Pr. vulgaris* with MIC values of 31.3 and 62.5 µg/mL respectively.

However, a comparison showed that the oils has greater potential of antimicrobial activities than those of 1,8-cineole as their main component [Table 1]. Otherwise, other compounds such as limonene, α-pinene, *p*-cymene, and terpineol-4-ol, which have relatively strong antimicrobial activities,^[27-29] may be responsible for this activity. Therefore, the synergistic effects of these active chemicals with other constituents of the essential oils should be taken into consideration for the antimicrobial activity.

Moreover, as indicated in a previous report about the other *Eucalyptus* species, gram-positive bacteria are more sensitive

to the essential oils than gram-negative bacteria,^[30] while our results did not show any selective antimicrobial activity on the basis of the cell wall differences of bacteria.

REFERENCES

1. Elliot WR, Jones DL. Encyclopedia of Australian plants suitable for cultivation, vol. 4, Melbourne: Lothian; 1986.
2. Ghisalberti EL, Bioactive acylphloroglucinol derivatives from *Eucalyptus* species. *Phytochemistry* 1996;41:7-22.
3. Yamashita N, Etoh H, Sakata K, Ina H, Ina K. New Acylated Rhaponticin Isolated from *Eucalyptus rubida* as a Repellent against the Blue Mussel *Mytilus edulis*. *Agr Biol Chem* 1989;53:2827-9.
4. Nishizawa M, Emura M, Kan Y, Yamada H, Ogawa K, Hamanaka N. Macrocarpals: HIV-RTase inhibitors of *Eucalyptus globules*. *Tetrahedron Lett* 1992;33:2983-6.
5. Murata M, Yamakoshi Y, Homma S, Arai K, Nakamura Y. Macrocarpals, Antibacterial Compounds from *Eucalyptus*, Inhibit Aldose Reductase. *Biosci Biotech Biochem* 1992;56:2062-3.
6. Osawa T, Namiki M, A Novel Type of Antioxidant Isolated from Leaf Wax of *Eucalyptus* leaves. *Agr Biol Chem* 1981;45:735-9.
7. Jovin E, Beara I, Mimica-Dukić N, Grbovic S, Bugarin D, Balog K. *Eucalyptus* species in Montenegro. Chemistry and antioxidant activity. *Planta Med* 2007;73:956.
8. Negahban M, Moharrampour S. Fumigant toxicity of *Eucalyptus intertexta*, *Eucalyptus sargentii* and *Eucalyptus camaldulensis* against stored-product beetles. *J Appl Entomol* 2007;131:256-61.
9. Takahashi T, Kokubo R, Sakaino M. Antimicrobial activities of *Eucalyptus* leaf extracts and flavonoids from *Eucalyptus maculate*. *Let Appl Microbiol* 2004;39:60-4.
10. Dethier M, Nduwimana A, Cordier Y, Menut C, Lamaty G. Aromatic plants of tropical central Africa. XVI. Studies on essential oils of five *Eucalyptus* species grown in Burundi. *J Essent Oil Res* 1994;6:469-73.
11. Bignell CM, Dunlop PJ, Brophy JJ, Jackson JF. Volatile leaf oils of some South-western and Southern Australian species of the genus *Eucalyptus* part VI - subgenus symphyomyrtus, section adnataria. *Flav Fragr J* 1995;10:359-64.
12. Bignell CM, Dunlop PJ, Brophy JJ, Jackson JF. Volatile leaf oils of some South-western and Southern Australian species of the genus *Eucalyptus*. Part 7. Subgenus Symphyomyrtus, section Exsertaria. *Flav Fragr J* 1996;11:35-41.
13. Bignell CM, Dunlop PJ, Brophy JJ. Volatile leaf oils of some South-western and Southern Australian species of the genus *Eucalyptus* (Series I). Part XVIII. A. Subgenus Monocalyptus. B. Subgenus Symphyomyrtus: (i) section Guilfoyleanae; (ii) section Bisectaria, series Accedentes, series Occidentales, series Levispermae, series Loxophlebae, series Macrocarpae, series Orbifoliae, series Calycogonae; (iii) section Dumaria, series Incrassatae and series Ovulares. *Flav Fragr J* 1997;12:423-32.
14. Wirthensohn MG, Sedgley M, Jones GP. Epicuticular wax of juvenile *Eucalyptus* leaves and headspace analysis of leaf volatiles. *J Essent Oil Res* 2000;12:401-11.
15. Bignell CM, Dunlop PJ, Brophy JJ, Jackson JF. Volatile leaf oils of some south-western and Southern Australian species of the genus *Eucalyptus*. Part IX. Subgenus Symphyomyrtus. Section Bisectaria. (a) series Elongatae, (b) unpublished series Stricklandiae, (c) series Kruseanae and (d) series Orbifoliae. *Flav Fragr J* 1996;11:95-100.
16. Sefidkon F, Assareh MH, Abravesh Z, Barazandeh MM. Chemical composition of the essential oils of four cultivated *Eucalyptus* species in Iran as medicinal plants (*E-microtheca*, *E-spathulata*, *E-largiflorens* and *E-torquata*). *Iran J Pharm Res* 2007;6:135-40.
17. Sefidkon F, Assareh MH, Abravesh Z, Mirza M. Chemical composition of the essential oils of five cultivated *Eucalyptus* species in Iran: *E. intertexta*, *E. platypus*, *E. leucoxydon*, *E. sargentii* and *E. camaldulensis*. *J Essent Oil Bearing Plants* 2006;9:245-50.
18. Assareh MH, Jaimand K, Rezaee MB. Chemical composition of the essential oils of six *Eucalyptus* species (Myrtaceae) from south west of Iran. *J Essent Oil Res* 2007;19:8-10.
19. Anonymous. European Pharmacopoeia. 3rd ed. Strasburg: Council of Europe; 1996. p. 121-2.
20. Wayne PA, National Committee for Clinical Laboratory Standard. In Performance standards for antimicrobial disk susceptibility test. 6th approved standard; 1997, M2-A6.
21. Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover RH. Manual of clinical microbiology. 7th ed. Washington: ASM; 1995. p.1773.
22. Gulluce M, Sokmen M, Sahin F, Sokmen A, Adiguzel A, Ozer H. Biological activities of the essential oil and methanolic extract of *Micromeria fruticosa* (L) Druce ssp serpyllifolia (Bieb) PH Davis plants from the eastern Anatolia region of Turkey. *J Sci Food Agric* 2004;84:735-41.
23. Gul HL, Ojanen T, Hanninen O. Antifungal evaluation of bis Mannich bases derived from acetophenones and their corresponding piperidinols and stability studies. *Biol Pharm Bull* 2002;25:1307-10.
24. Rosato A, Vitali C, De Laurentis N, Armenise D, Milillo MA. Antibacterial effect of some essential oils administered alone or in combination with Norfloxacin. *Phytomedicine* 2007;14:727-32.
25. Pitarokili D, Tzakou O, Loukis A, Harvala C. Volatile metabolites from *Salvia fruticosa* as antifungal agents in soilborne pathogens. *J Agric Food Chem* 2003;51:3294-301.
26. Sonboli A, Babakhani B, Mehrabian AR. Antimicrobial activity of six constituents of essential oil from *Salvia*. *Z Naturforsch C* 2006;61:160-4.
27. Dorman HJD, Deans SG. Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. *J Appl Microbiol* 2000;88:308-16.
28. Carson CF, Riley TV. Antimicrobial activity of the major components of the essential oil of *Melaleuca alternifolia*. *J Appl Bacteriol* 1995;78:264-9.
29. Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils - A review. *Food Chem Toxicol* 2008;46:446-75.
30. Gilles M, Zhao J, An M, Agboola S. Chemical composition and antimicrobial properties of essential oils of three Australian *Eucalyptus* species. *Food Chem* 2010;119:731-7.

Source of Support: Nil, Conflict of Interest: None declared