

45°C to give 1.91 g of residue. The residue was stored at -70°C prior to use as a test sample (RCE).

Extraction of tyrosinase from Melan-A cells

Melan-a cells were disrupted via resuspension in a tyrosinase buffer (80 mM PO₄ buffer + 1% Triton-X100 + 100 mg/ml PMSF), followed by sonication in an ice bath. After 15 minutes of centrifugation at 12,500 rpm, the supernatants were utilized in the tyrosinase assays. 150 mg of proteins were required for each of the reactions.^[7]

Tyrosinase activity assay methods

The dopa-oxidase activity of tyrosinase was spectrophotometrically determined as described previously, with minor modifications.^[8] Each concentration of the test substance was dissolved in MeOH. 120 ml of L-dopa (8 mM, dissolved in 67 mM phosphate buffer, pH 6.8) and 40 ml of either the same buffer or of the test sample, were added to a 96-well microplate, after which 40 ml of mushroom tyrosinase (125 U) or melan-a cell tyrosinase (150 mg) were added. After 30 minutes of incubation at 37°C, the quantity of dopachrome in the reaction mixture was determined based on optical density at a wavelength of 492 nm (OD 492). Kojic acid was used as a reference agent.^[9]

Cell line and culture procedures

In order to evaluate the effects of RCE on melanocytes, we utilized a melan-a cell model. A pigmented melanocyte cell line, 'melan-a', was previously derived from the normal epidermal melanoblasts of embryos of inbred C57BL mice.^[10] The melan-a cells were cultured in RPMI 1640 medium with 10% fetal bovine serum (FBS) in 200 nM TPA. 10 ml of the medium was added to a 100 mm culture dish, and the cells were seeded at a density of approximately 5×10^5 cell/dish. The cells were grown to confluency, seeded at a concentration of 10^5 cells/well on 24-well plates, then incubated for an additional 24 hours. Each well received a daily exchange of 990 ml of medium and was treated with 10 ml of the sample for three days.

Determination of cell viabilities

The percentage of viable cells was determined by staining the cell population with crystal violet. After the media was removed from each of the wells, the cells were washed with PBS. 200 ml of crystal violet (CV 0.1%, 10% EtOH, the rest is PBS) was then added. This was incubated at room temperature for 5 minutes and washed twice in water. After the addition of 1 ml of EtOH, the samples were shaken at room temperature. The UV absorption of the resultant supernatant was measured at a wavelength of 590 nm.

Determination of melanin contents in cells

After the media was removed from each of the wells, the wells were washed in PBS. After this step, 1 ml of 1 N

NaOH was added in order to dissolve the melanin. UV absorption was then measured at 400 nm, and the melanin content per well was calculated.

Measurement of depigmenting activities on brown guinea pig skin

Brown guinea pig skin is an excellent pigmentation model because its pigmentation system is resemble to human skin.^[11,12] Three female brown guinea pigs weighing approximately 450 g were used in this study. The animals were purchased from OrientBio Inc., at Seoul, Korea. They were housed in an environment of constant temperature, humidity and light/dark cycle. UVB-induced hyper pigmentation elicited on the shaven backs of the animals. 500 mJ/cm² of UV-B was exposed on separated areas of the back once a week three times. Test samples or vehicle (70% ethanol) were then topically applied to the hyper pigmented areas (10 ml/circle) once a day for 8 weeks.^[13,14]

Statistical analysis

The data was expressed as the means \pm S.E. Statistical significance was evaluated via one-way ANOVA.

RESULTS

Tyrosinase Inhibitory effect

The RCE inhibited melan-a and mushroom tyrosinase activity by 49% and 52%, respectively, at 500 mg/ml [Figure 1]. Although this inhibitory effect was lower than that of kojic acid at identical concentrations, the extract produced a graded dose response.

Inhibitory effect on melanin biosynthesis in Melan-A cells

The results of the inhibitory experiments on melanin biosynthesis in melan-a cells are presented in [Table 1]. Although RCE induced a slight decrease (4.9%) in cell viability, treatment with 100 μ g/ml of RCE resulted in a 20.6% reduction in melanin production in the melan-a cells.

Depigmenting effect on hyper pigmented brown guinea pig skin

Figure 2 shows the depigmenting activity of RCE in the hyper pigmented brown guinea pig skin. Hyper pigmentation was elicited on the dorsal skin of the brown guinea pigs using UV-B radiation. UV-B irradiation induces the tanning or burning of skin.^[15,16] After treatment with 3% RCE for 6 weeks, although DL-value was not statistically significant, a visible decrease in hyper pigmentation was observed when compared to the vehicle treat group. During all experimental days, visible erythema was not observed on the dorsal skin treated with RCE.




Figure 1: The effects of RCE on tyrosinase activity (a) Inhibitory effects on melan-a cell tyrosinase (b) Inhibitory effects on mushroom tyrosinase, Each value represents the mean \pm S.E. of three experiments, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared to vehicle

Table 1: Effects of RCE on cell growth and melanin production in melan-a cells

Samples	Concentrations (µg/ml)	Melanin production (%)	Cell viability (%)
RCE	1	102.3 \pm 5.0	99.9 \pm 2.3
	10	92.7 \pm 4.2	95.5 \pm 5.1
	100	81.7 \pm 6.0*	95.2 \pm 5.9
Kojic acid	1	107.1 \pm 9.9	96.6 \pm 7.5
	10	96.0 \pm 2.6	98.6 \pm 6.4
	100	91.9 \pm 4.0	84.3 \pm 5.8*
PTU (Phenyl thiourea)	1	88.9 \pm 7.6	97.3 \pm 1.3
	10	41.4 \pm 9.3***	80.1 \pm 9.3*
	100	25.3 \pm 8.4***	72.3 \pm 9.4**

Kojic acid and PTU were used as reference materials. Each value is representative of the mean \pm S.E. of three experiments, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared to vehicle

DISCUSSION

The inhibitory effects of the root of RCE on melanin production were evaluated by assessing its tyrosinase inhibitory effects, melanin production-inhibitory properties in melan-a cells and depigmenting ability in brown guinea pig skin. Overall, the results of the present study revealed that the root of *C. fortunei* inhibits melanogenesis. RCE was shown to inhibit tyrosinase activity. In melan-a cells, the melanin inhibitory effect of RCE was lower than that of Phenyl thiourea (PTU), but cell toxicity of RCE was vastly lower at same concentrations [Table 1]. In addition, RCE treatment reduced the melanin content in brown guinea pig skin. Based on these results, RCE can be expected to inhibit melanogenesis. The results of the present study indicate that RCE may prove useful as a skin depigmenting material. Furthermore, the root of *C. fortunei* might prove to be a new valuable herbal source for the development of skin depigmenting agents.

REFERENCES

- No JK, Soung DY, Kim YJ, Shim KH, Jun YS, Rhee SH, *et al.* Inhibition of tyrosinase by green tea components. *Life Sci* 1999;65:241-6.




Figure 2: Depigmenting effects of RCE on UVB-induced hyperpigmentation in brown guinea pigs skin. Photographs showing changes in hyper pigmented guinea pig skin treated with vehicle alone (a) or 3% RCE in vehicle (b) after six weeks. The degree of the pigmentation decrease was represented as the DL-value (c)

- Choi SY, Kim S, Kim H, Suk K, Hwang JS, Lee BG, *et al.* (4-Methoxy-benzylidene)-(3-methoxy-phenyl)-amine, a nitrogen analog of stilbene as a potent inhibitor of melanin production. *Chem Pharm Bull (Tokyo)* 2002;50:450-2.
- Baurin E, Arnoult T, Scior T, Do QT, Bernard P. Preliminary screening of some tropical plants for anti-tyrosinase activity. *J Ethnopharmacol* 2002;82:155-8.
- Iozumi K, Hoganson GE, Pennella R, Everett MA, Fuller BB. Role of tyrosinase as the determinant of pigmentation in cultured human melanocytes. *J Invest Dermatol* 1993;100:806-11.
- Bae KH. The medicinal plants of Korea. Seoul: Kyohak-sa Press; 2001.
- Guo JP, Pang J, Wang XW, Shen ZQ, Jin M, Li JW. *In vitro* screening of traditionally used medicinal plants in China against enteroviruses. *World J Gastroenterol* 2006;12:4078-81.

7. Fuller BB, Drake MA, Spaulding DT, Chaudhry F. Down regulation of tyrosinase activity in human melanocyte cell culture by yohimbine. *J Invest Dermatol* 2000;114:268-76.
8. Koketsu M, Choi SY, Ishihara H, Lim BO, Kim H, Kim SY. Inhibitory effects of 1, 3-selenazol-4-one derivatives on mushroom tyrosinase. *Chem Pharm Bull (Tokyo)* 2002;50:1594-6.
9. Parvez S, Kang M, Chung HS, Cho C, Hong MC, Shin MK, *et al.* Survey and mechanism of skin depigmenting and lightening agents. *Phytother Res* 2006;20:921-34.
10. Bennett DC, Cooper PJ, Hart IR. A line of non-tumorigenic mouse melanocytes, syngeneic with the B16 melanoma and requiring a tumor promoter for growth. *Int J Cancer* 1987;39:414-8.
11. Yoshida M, Hirotsu S, Nakahara M, Uchiwa H, Tomita Y. Histamine is involved in ultraviolet B-induced pigmentation of guinea pig skin. *J Invest Dermatol* 2002;118:255-60.
12. Park KT, Kim JK, Hwang D, Yoo Y, Lim YH. Inhibitory effect of mulberroside A and its derivatives on melanogenesis induced by ultraviolet B irradiation. *Food Chem Toxicol* 2011;49:3038-45.
13. Ando H, Ryu A, Hashimoto A, Oka M, Ichihashi M. Linoleic acid and α -linoleic acid lightens ultraviolet-induced hyper pigmentation of the skin. *Arch Dermatol Res* 1998;290:375-81.
14. Imokawa G, Kawai M, Mishima Y, Motegi I. Differential analysis of experimental hypermelanosis induced by UVB, PUVA and allergic dermatitis using a brownish guinea pig model. *Arch Dermatol Res* 1986;278:352-62.
15. Choi SY, Kim S, Hwang JS, Lee BG, Kim H, Kim SY. Benzylamide derivative compound attenuates the ultraviolet B-induced hyper pigmentation in the brownish guinea pig skin. *Biochem Pharmacol* 2004;67:707-15.
16. Schempp CM, Winghofer B, Muller K, Schulte-Monting J, Mannel M, Schopf E, *et al.* Effect of oral administration of *Hypericum perforatum* extract (St. John's Wort) on skin erythema and pigmentation induced by UVB, UVA, visible light and solar simulated radiation. *Phytother Res* 2003;17:141-6.

Cite this article as: Choi SY. Inhibitory effects of *Cyrtomium fortunei* J. Smith root extract on melanogenesis. *Phcog Mag* 2013;9:227-30.

Source of Support: Nil. **Conflict of Interest:** None declared.

Author Help: Reference checking facility

The manuscript system (www.journalonweb.com) allows the authors to check and verify the accuracy and style of references. The tool checks the references with PubMed as per a predefined style. Authors are encouraged to use this facility, before submitting articles to the journal.

- The style as well as bibliographic elements should be 100% accurate, to help get the references verified from the system. Even a single spelling error or addition of issue number/month of publication will lead to an error when verifying the reference.
- Example of a correct style
Sheahan P, O'leary G, Lee G, Fitzgibbon J. Cystic cervical metastases: Incidence and diagnosis using fine needle aspiration biopsy. *Otolaryngol Head Neck Surg* 2002;127:294-8.
- Only the references from journals indexed in PubMed will be checked.
- Enter each reference in new line, without a serial number.
- Add up to a maximum of 15 references at a time.
- If the reference is correct for its bibliographic elements and punctuations, it will be shown as CORRECT and a link to the correct article in PubMed will be given.
- If any of the bibliographic elements are missing, incorrect or extra (such as issue number), it will be shown as INCORRECT and link to possible articles in PubMed will be given.