

Ingenine F: A New Cytotoxic Tetrahydro Carboline Alkaloid from the Indonesian Marine Sponge *Acanthostrongylophora ingens*

Sabrin Ibrahim^{1,2}, Gamal Mohamed^{3,4}, Rwaida Al Haidari¹, Amal El-Kholy^{5,6}, Mohamed Zayed^{1,7}

¹Department of Pharmacognosy and Pharmaceutical Chemistry, College of Pharmacy, Taibah University, Al Madinah Al Munawwarah 41477, Saudi Arabia, ²Department of Pharmacognosy, Faculty of Pharmacy, Assiut University, Assiut 71526, Egypt, ³Department of Natural Products and Alternative Medicine, Faculty of Pharmacy, King Abdulaziz University, Jeddah, 21589, Saudi Arabia, ⁴Department of Pharmacognosy, Faculty of Pharmacy, Al-Azhar University, Assiut Branch, Assiut 71524, Egypt, ⁵Department of Clinical and Hospital Pharmacy, College of Pharmacy, Taibah University, Al Madinah Al Munawwarah 30078, Saudi Arabia, ⁶Department of Clinical Pharmacy, Faculty of Pharmacy, Ain-Shams University, Cairo 11566, ⁷Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Al-Azhar University, Cairo, Egypt

Submitted: 23-10-2017

Revised: 23-11-2017

Published: 10-04-2018

ABSTRACT

Background: Marine organisms are established to be a wealthy source of bioactive compounds with diverse chemical structures and bioactivities. *Acanthostrongylophora ingens* is known to be rich with pyrimidine β -carboline and manzamine-type alkaloids. The goal of the present work is to isolate and identify new alkaloids from *A. ingens* as well as to assess the cytotoxic potential of these metabolites towards various cancer cell lines. **Methods:** The crude MeOH extract of the sponge was separated by vacuum liquid chromatography (VLC), using *n*-hexane, EtOAc, and MeOH. The EtOAc fraction was chromatographed on VLC, SiO₂, sephadex LH-20, and RP₁₈ columns, affording four metabolites. Their structures were identified using infrared, ultraviolet, high-resolution mass spectrometry, and nuclear magnetic resonance spectroscopic techniques, as well as comparison with the published data. **Results:** A new 1,2,3,4-tetrahydro- β -carboline (TH β Cs) alkaloid, ingenine F (4) and three known compounds: Annomontine (1), acanthomine A (2), and 1-oxo-1,2,3,4-TH β Cs (3) were isolated and identified. Ingenine F (4) exhibited cytotoxic activity toward hormone-dependent breast carcinoma (MCF7), colon carcinoma (HCT116), and lung carcinoma (A549) cell lines with IC₅₀ values of 2.82, 1.00, and 2.37 μ M, respectively, compared to doxorubicin (IC₅₀ 0.012, 0.036, and 0.102 μ M, respectively). **Conclusion:** It is the first report for the isolation of TH β Cs alkaloids from *A. ingens*. The TH β Cs alkaloid with *N*-methylbutyramide unit as found in ingenine F is very rarely encountered in nature. Ingenine F may provide new promising candidates for potential cytotoxic agent.


Key words: *Acanthostrongylophora ingens*, alkaloid, cytotoxic activity, Ingenine F, Tetrahydro- β -carboline

SUMMARY

- Ingenine F, a new 1,2,3,4-TH β Cs derivative (4) and three known alkaloids (1-3) were isolated from *A. ingens*. Their structures were verified by various spectroscopic analyses. Compound 4 had potent cytotoxic effect toward MCF7, HCT116, and A549 cancer cell lines.

Abbreviations used: 1D: One-dimensional; 2D: Two-dimensional; CC: Column chromatography; COSY: Correlations spectroscopy; DMSO: Dimethyl sulfoxide; HMBC: Heteronuclear multiple bond

correlation experiment; HRESIMS: High resolution electrospray ionization mass spectrometry; HSQC: Heteronuclear single quantum correlation; IR: Infrared; LCQ: Liquid chromatography quadrupole; LTO: Linear trap quadrupole; NMR: Nuclear magnetic resonance; RP: Reversed phase; SiO₂: Silica gel; TLC: Thin-layer chromatography; UV: Ultraviolet; VLC: Vacuum liquid chromatography.



Correspondence:

Prof. Sabrin Ibrahim,
Department of Pharmacognosy and
Pharmaceutical Chemistry, College of Pharmacy,
Taibah University, Al Madinah Al Munawwarah
41477, Saudi Arabia.
E-mail: sbrahim@taibahu.edu.sa; sabrinshaur@
gmail.com
DOI: 10.4103/pm.pm_489_17

Access this article online

Website: www.phcog.com

Quick Response Code:



INTRODUCTION

The ability to use nature as an inspiring framework for devolving novel products is an interesting area of scientific research. In fact, large numbers of current pharmacopoeias derive directly or indirectly from natural products. Marine organisms are established to be fruitful origin of bioactive novel compounds with diverse chemical structures that are of great usefulness as new drug leads for treating various ailments. *Acanthostrongylophora ingens* (order Haplosclerida; family Petrosiidae) is considered to be a rich source of pyrimidine β -carboline^[1-3] and manzamine-type alkaloids.^[4-8] These compounds exhibit a diverse range of bioactivities, including antimalarial, antileishmanial, antituberculosis,^[4-6] cytotoxic,^[1,2,9-11]

and anti-HIV.^[4-6] In our further search for structurally interesting and bioactive compounds from the sponge *A. ingens*, a new 1,2,3,4-tetrahydro- β -carboline (TH β Cs) alkaloid, ingenine

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Ibrahim S, Mohamed G, Al Haidari R, El-Kholy A, Zayed M. Ingenine F: A new cytotoxic tetrahydro carboline alkaloid from the Indonesian marine sponge *Acanthostrongylophora ingens*. Phcog Mag 2018;14:231-4.




Figure 1: Structures of isolated compounds 1-4

F (4), and three known compounds were isolated [Figure 1]. The structural identities of the isolated metabolites were proven by interpreting the spectroscopic data. Herein, the isolation, structural characterization, and cytotoxic potential of 4 were described.

MATERIALS AND METHODS

General experimental procedures

A JASCO polarimeter was utilized for measuring optical rotation (Jasco Co., Tokyo, Japan). An electrothermal-9100 digital melting point instrument was used to get the melting points (Perkin-Elmer, Waltham, MA, USA). Ultraviolet (UV) data were recorded by Hitachi-300 spectrophotometer (Kyoto, Japan). The infrared (IR) absorbance was obtained by a Shimadzu Infrared 400 spectrophotometer (Shimadzu, Kyoto, Japan). A liquid chromatography quadrupole (LCQ) DECA mass spectrometer (ThermoFinnigan, Bremen, Germany) was utilized to obtain the electrospray ionization mass spectroscopy (ESIMS) spectrum. High resolution electrospray ionization mass spectrometry (HRESIMS) was assessed by linear trap quadrupole (LTQ)-Orbitrap MS spectrometer. A Bruker Avance DRX 500 MHz spectrometer was used for measuring nuclear magnetic resonance (NMR) data (Bruker BioSpin, Billerica, MA, USA). High-performance liquid chromatography (HPLC) was carried out on a C-18 column Eurospher 100 (300 mm × 8 mm, Knauer, Berlin, Germany) at 5.0 ml/min flow rate. Sephadex LH-20 (0.25–0.1 mm), silica gel (0.063–0.200 mm), and RP₁₈ (0.04–0.063 mm) (Merck, Darmstadt, Germany) were utilized for column chromatography. For TLC, precoated plates silica gel (60 F₂₅₄, 0.2 mm, Merck, Darmstadt, Germany) were used. RP₁₈ six mL standard extraction tube (40–63 μm) (Merck, Darmstadt, Germany) was used for the purification of compounds.

Animal material

The sponge material was obtained in 2010 from Sulawesi Island by scuba diving at 12 m depth. The collected material was immediately freeze-dried after collection. A specimen (ZMAPOR. 17795-A) was preserved in the Zoological Museum (Amsterdam University).

Extraction and isolation

The dried sponge (1.3 kg) was extracted with MeOH (4 × 2 L) at room temperature (25°C). The combined extract was concentrated to get a brown residue (39.0 g). The latter was submitted to vacuum liquid chromatography (VLC), utilizing *n*-hexane, EtOAc, and MeOH (2 × 1.0 L, each) to get *n*-hexane (5.9 g), EtOAc (7.8 g), and MeOH (20.8 g) fractions. The EtOAc (7.8 g) fraction was chromatographed, using CHCl₃:MeOH gradient on VLC to get 13 subfractions: EA-1 to EA-13. Fraction EA-3 (380 mg) was separated using MeOH: CHCl₃ (90:10) on a sephadex LH-20 CC (50 g × 50 cm × 3 cm) to give 3 subfractions: EA-3A to EA-3C. Subfraction EA-3B (71 mg) was submitted to SiO₂ CC (40 g × 50 × 3 cm), utilizing CHCl₃:MeOH (95:5–80:20) to get impure 1. Purification of 1 was performed using a semi-preparative HPLC to yield 1 (10.2 mg, yellow needles). Fraction EA-4 (148 mg) was separated on a SiO₂ CC (40 g × 50 cm × 2 cm), utilizing a CHCl₃:MeOH gradient to give 2 subfractions: EA-4A and EA-4B. Subfraction EA-4A (69 mg) was subjected to SiO₂ CC (40 g × 50 cm × 2 cm) with CHCl₃:MeOH gradient to afford impure 2 and 3. Each compound was purified separately on RP₁₈ extraction tube, using H₂O: acetonitrile gradient elution to yield 2 (9.5 mg, red needles) and 3 (7.1 mg, white amorphous powder). Fraction EA-5 (65 mg) was subjected to RP₁₈ CC (20 g × 50 cm × 2 cm) eluting with H₂O: MeOH gradient to get 2 subfractions: EA-5A and EA-5B. Subfraction EA-5A (29 mg) was separated on HPLC to yield 4 (2.7 mg, pale yellow powder).

Spectral data

Ingenine F (4): pale yellow powder; [α]_D²⁰ + 13.9 (*c* 0.3, CH₃OH); UV (MeOH) λ_{\max} (log ϵ) 226 (4.37), 300 (3.98) nm; IR (KBr) ν_{\max} 3210, 2924, 1649, 1564, 1058 cm⁻¹; NMR data [Table 1]; HRESIMS *m/z* 272.1764 (M + H)⁺ (calcd for C₁₆H₂₂N₃O, 272.1763).

Cytotoxicity assay

The *in vitro* cytotoxic activity of 4 was assessed toward colon carcinoma (HCT116), hormone-dependent breast carcinoma (MCF7), human cervix carcinoma (Hela), lung carcinoma (A549), and brain tumor (PC12) cell lines (ATCC, Rockville, MD) as previously stated.^[12–15] DMSO and doxorubicin were utilized as negative and positive standards, respectively.


RESULTS AND DISCUSSION

Compound 4 was separated as pale yellow powder and afforded positive tests for alkaloids.^[16] Its molecular formula C₁₆H₂₁N₃O was estimated by the pseudomolecular ion peak at *m/z* 272.1764 (M + H)⁺ (calcd for C₁₆H₂₂N₃O, 272.1763) in HRESIMS spectrum, requiring eight DBE. Its UV absorbances at 226, and 300 nm specified the existence of a TH β C chromophore.^[17,18] Its IR showed bands at 3210 (NH), 2924 (C-H), and 1649 (C = O) cm⁻¹. The chemical shifts together with NMR experiments suggested 4 to have a C-1 substituted 1,2,3,4-tetrahydro β -carboline skeleton.^[18] ¹³C NMR and heteronuclear single quantum correlation (HSQC) spectra of 4 exhibited 16 carbons, including 8 aromatic carbons, an amide carbonyl (δ_{C} 156.9), 5 methylenes, NH-bonded methine, and N-methyl signal (δ_{C} 48.6). The ¹H NMR spectrum showed four-coupled aromatic protons at δ_{H} 7.45 (H-5), 7.01 (H-6), 7.11 (H-7), and 7.37 (H-8), suggesting the presence of an ABCD spin system characteristic for a 1,2-disubstituted benzene. They correlated to the carbons, resonating at δ_{C} 118.1, 119.1, 121.9, and 111.4, respectively, in the HSQC spectrum. In addition, signals for a NH-bonded methine at δ_{H} 4.71 (H-1)/ δ_{C} 51.9 (C-1), two coupled methylenes at δ_{H} 3.58 (m, H-3)/ δ_{C} 40.8 (C-3) and δ_{H} 2.91 (m, H-4)/ δ_{C} 18.2 (C-4), and NH group at δ_{H} 8.21 (brs, 2-NH) were attributable to 1,2,3,4-tetrahydro-pyridine ring of the β -carboline moiety. The HMBC correlations of H-1 to C-1a, C-4a, and C-3, H-3 to C-1 and C-4a, H-4 to C-1a, C-4a, and C-3, H-5 to C-7 and C-8a, H-6 to C-5a and C-8, H-7 to C-5 and C-8a, and H-8 to C-5a

Table 1: NMR data of compound 4 (DMSO- d_6 , 500 and 125 MHz)

Position	δ_H multiplicity (J in Hz)	δ_C	COSY	HMBC
1	4.71 t (7.2)	51.9 CH	1'	1a, 3, 4a, 3'
1a	-	130.1 C	-	-
3	3.58 m	40.8 CH ₂	4	1, 4a
4	2.91 m	18.2 CH ₂	3	1a, 3, 4a
4a	-	105.9 C	-	-
5	7.45 d (7.6)	118.1 CH	6	7, 8a
5a	-	125.8 C	-	-
6	7.01 t (7.6)	119.1 CH	5, 7	5a, 8
7	7.11 t (7.6)	121.9 CH	6, 8	5, 8a
8	7.37 d (7.6)	111.4 CH	7	5a, 6
8a	-	136.2 C	-	-
4'	2.25 m	28.7 CH ₂	2'	1, 1a, 2'
	1.95 m			
3'	1.75 m	24.2 CH ₂	1', 3'	1, 1'
2'	3.21 m	40.3 CH ₂	3'	1', 4'
1'	-	156.9 C	-	-
1'-NH-CH ₃	3.17 s	48.6 CH ₃	-	1'
NH	11.22 s	-	-	4a, 5a, 8a
2-NH	8.21 brs	-	-	-

NMR: Nuclear magnetic resonance; DMSO: Dimethyl sulfoxide

**Figure 2:** ^1H - ^1H COSY and HMBC correlations of 4

and C-6 established the TH β C moiety of 4 [Figure 2]. In the ^1H and ^{13}C spectra, signals for three-coupled aliphatic methylenes (δ_H 3.21 [m, H-2']/ δ_C 40.3 [C-2'], 1.75 [m, H-3']/ δ_C 24.2 [C-3'], and 2.25 and 1.95 [each m, H-4']/ δ_C 28.7 [C-4']), and NH-bonded methyl (δ_H 3.17/ δ_C 48.6), and an amide carbonyl (δ_C 156.9) characteristic for *N*-methylbutyramide moiety were observed. This moiety was assured by the cross peaks of H-2'/C-1' and C-4', H-3'/C-1', H-4'/C-2', and 1'-NH-CH₃/C-1' in HMBC. The connectivity of the *N*-methylbutyramide moiety at C-1 of the TH β C moiety was assured by the HMBC relations of H-2' and H-3'/C-1, and H-4'/C-1 and C-1a. Due to the scarcity of the compound, C-1 relative configuration was assigned to be *R*-form based on the comparison ^1H and ^{13}C shifts and *J* value of H-1 as well as optical rotation of 4 with those of series of analogous compounds.^[19,20] On the basis of the above evidence, 4 was identified as a TH β C alkaloid substituted at C-1 by *N*-methylbutyramide. Thus, compound 4 was finally assigned as (*R*)-*N*-methyl-4-(2,3,4,9-tetrahydro-1*H*-pyrido [3,4-*b*] indol-1-yl) butanamide and named ingenine F.

The NMR data of compounds 1-3 were in accordance with those reported previously for anomontine (1),^[1,2] acanthomine A (2),^[1] and 1-oxo-1,2,3,4-TH β Cs (3).^[1]

Ingenine F (4) was assessed for its cytotoxic potential toward A549, MCF7, HCT116, PC12, and Hela cancer cell lines. It displayed cytotoxic activity against MCF7, HCT116, and A549 with IC₅₀ values of 2.82, 1.00, and 2.37 μM , respectively, compared to doxorubicin (IC₅₀ 0.012, 0.036, and 0.102 μM , respectively). However, it was inactive toward Hela and PC12 cancer cell lines.

CONCLUSION

A new 1,2,3,4-TH β Cs derivative (4) and three known compounds (1-3) were separated from the Indonesian marine sponge *A. ingens*. Their structures were identified by the aid of comprehensive spectroscopic analysis. Compound 4 exhibited cytotoxic effect toward MCF7, HCT116, and A549 cancer cell lines.

Acknowledgements

We are thankful to Dr. Rob vanSoest for the identification of the sponge (Zoological Museum, University of Amsterdam).

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Ibrahim SR, Ebel RA, Ebel R, Proksch P. Acanthomine A, a new pyrimidine- β -carboline alkaloid from the sponge *Acanthostrongylophora ingens*. *Nat Prod Commun* 2008;3:175-8.
- Ibrahim SR, Mohamed GA, Zayed MF, Sayed HM. Ingenines A and B, two new alkaloids from the Indonesian sponge *Acanthostrongylophora ingens*. *Drug Res (Stuttg)* 2015;65:361-5.
- Ibrahim SR, Mohamed GA. Ingenine E, a new cytotoxic β -carboline alkaloid from the Indonesian sponge *Acanthostrongylophora ingens*. *J Asian Nat Prod Res* 2017;19:504-9.
- Peng J, Hu JF, Kazi AB, Li Z, Avery M, Peraud O, *et al.* Manadomanzamines A and B: A novel alkaloid ring system with potent activity against mycobacteria and HIV-1. *J Am Chem Soc* 2003;125:13382-6.
- Rao KV, Kasanah N, Wahyuono S, Tekwani BL, Schinazi RF, Hamann MT, *et al.* Three new manzamine alkaloids from a common Indonesian sponge and their activity against infectious and tropical parasitic diseases. *J Nat Prod* 2004;67:1314-8.
- Rao KV, Donia MS, Peng J, Garcia-Palmero E, Alonso D, Martinez A, *et al.* Manzamine B and E and ircular A related alkaloids from an Indonesian acanthostrongylophora sponge and their activity against infectious, tropical parasitic, and Alzheimer's diseases. *J Nat Prod* 2006;69:1034-40.
- El-Desoky AH, Kato H, Eguchi K, Kawabata T, Fujiwara Y, Losung F, *et al.* Acantholactam and pre-neo-kaulamine, manzamine-related alkaloids from the Indonesian marine sponge *Acanthostrongylophora ingens*. *J Nat Prod* 2014;77:1536-40.
- Furusato A, Kato H, Nehira T, Eguchi K, Kawabata T, Fujiwara Y, *et al.* Acanthomanzamines A-E with new manzamine frameworks from the marine sponge *Acanthostrongylophora ingens*. *Org Lett* 2014;16:3888-91.
- Kong F, Andersen RJ. Ingenamines A and B, new cytotoxic alkaloids from the marine sponge *Xestospongia ingens*. *Tetrahedron* 1994;50:6137-44.
- Kong F, Andersen RJ. Ingenamine alkaloids isolated from the sponge *Xestospongia ingens*: structures and absolute configurations. *Tetrahedron Lett* 1995;51:2895-906.
- Andersen RJ, van Soest RW, Kong F. In: Pelletier SW, editor. *Alkaloids: Chemical and Biological Perspectives*. New York: Pergamon; 1996;10:301-55.
- Borenfreund E, Babich H, Martin-Alguacil N. Rapid chemosensitivity assay with human normal and tumor cells *in vitro*. *In Vitro Cell Dev Biol* 1990;26:1030-4.
- Mohamed GA, Ibrahim SR, Ross SA. New ceramides and isoflavone from the Egyptian *Iris germanica* L. rhizomes. *Phytochem Lett* 2013;6:340-4.
- Ibrahim SR, Abdallah HM, Mohamed GA, Ross SA. Integracides H-J: New tetracyclic triterpenoids from the endophytic fungus *Fusarium* sp. *Fitoterapia* 2016;112:161-7.

15. Ibrahim SR, Elkhayat ES, Mohamed GA, Khedr AI, Fouad MA, Kotb MH, *et al.* Aspernolides F and G, new butyrolactones from the endophytic fungus *Aspergillus terreus*. *Phytochem Lett* 2015;14:84-90.
16. Cannell RJ, editor. How to approach the isolation of a natural product. In: *Methods in Biotechnology: Natural Products Isolation*. Vol. 4. Totowa New Jersey USA: Humana Press; 1998. p. 1-51.
17. El-Shazly A, Wink M. Tetrahydroisoquinoline and beta-carboline alkaloids from haloxylon articulatum (Cav.) bunge (Chenopodiaceae). *Z Naturforsch C* 2003;58:477-80.
18. Prinsep MR, Blunt JW, Munro MH. New cytotoxic beta-carboline alkaloids from the marine bryozoan, *Cribricellina cribraria*. *J Nat Prod* 1991;54:1068-76.
19. Megyesi R, Forró E, Fülöp F. Enzymatic strategy for the resolution of new 1-hydroxymethyl tetrahydro- β -carboline derivatives in batch and continuous-flow systems. *ChemistryOpen* 2016;5:254-60.
20. Arazy Z, Czarnocki Z, Wojtasiewicz K, Maurin JK. Enantioselective synthesis of (1R)-1-(hydroxymethyl)-2-acetyl-1,2,3,4-tetrahydro- β -carboline from L-(+)-tartaric acid. *Tetrahedron Asymmetry* 2000;11:2793-800.