

- Gard 1992;79:53-64.
14. Jiménez-González FJ, Veloza LA, Sepúlveda-Arias JC. Anti-infectious activity in plants of the genus *Tabebuia*. *Univ Sci (Bogota)* 2013;18:257-67.
 15. Gómez-Estrada H, Díaz-Castillo F, Franco-Ospina L, Mercado-Camargo J, Guzmán-Ledezma J, Medina JD, *et al.* Folk medicine in the Northern coast of Colombia: An overview. *J Ethnobiol Ethnomed* 2011;7:27.
 16. Waksmundzka-Hajnos M, Sherma J, Kowalska T. *Thin Layer Chromatography in Phytochemistry*. Boca Raton, FL: CRC Press; 2008.
 17. Singleton V, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am J Enol Vitic* 1965;16:144-58.
 18. Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT Food Sci Technol* 1995;28:25-30.
 19. Ou B, Hampsch-Woodill M, Prior RL. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *J Agric Food Chem* 2001;49:4619-26.
 20. Cao G, Prior RL. Measurement of oxygen radical absorbance capacity in biological samples. *Methods Enzymol* 1999;299:50-62.
 21. Mosmann T. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983;65:55-63.
 22. Hemamalini K, Lavanya CH, Bhargava A, Vasireddy U. Anti-ulcer activity of methanolic extracts of *Wattakaka volubilis* and *Tabebuia rosea* in rats. *Asian J Pharm Clin Res* 2012;5:242-6.
 23. Sathiya M, Muthuchelian K. Studies on phytochemical profile and antibacterial activity of ethanolic leaf extract of *Tabebuia rosea* (Bertol.) DC. *Ethnobot Leaf* 2008;2008:152.
 24. Joselin J, Brintha TS, Florence AR, Jeeva S. Phytochemical evaluation of Bignoniaceae flowers. *J Chem Pharm Res* 2013;5:106-11.
 25. Compadre CM, Jáuregui JF, Nathan PJ, Enriquez RG. Isolation of 6-O-(p-coumaroyl)-catalpol from *Tabebuia rosea*. *Planta Med* 1982;46:42-4.
 26. Lino von Poser G, Schripsema J, Henriques AT, Rosendal Jensen S. The distribution of iridoids in Bignoniaceae. *Biochem Syst Ecol* 2000;28:351-66.
 27. Joshi KC, Prakash L, Shah RK. Chemical examination of the roots of *Tabebuia rosea* and heartwood of *Oroxylum indicum*. *Planta Med* 1977;31:257-8.
 28. Joshi K, Prakash L, Singh P. Quinones and other constituents from *Tabebuia rosea*. *Planta Med* 1973;12:942-3.
 29. Sichaem J, Kaennakam S, Siripong P, Tip-Pyang S. Tabebuialdehydes A-C, cyclopentene dialdehyde derivatives from the roots of *Tabebuia rosea*. *Fitoterapia* 2012;83:1456-9.
 30. Franco Ospina LA, Castro Guerrero JP, Ocampo Buendía YC, Pájaro Bolívar IB, Díaz Castillo F. Antiinflammatory, antioxidant and antibacterial activity of two species of *Tabebuia* genus. *Rev Cubana Plant Med* 2013;18:34-46.
 31. Suo M, Ohta T, Takano F, Jin S. Bioactive phenylpropanoid glycosides from *Tabebuia avellanedae*. *Molecules* 2013;18:7336-45.
 32. Rahman MM, Islam MB, Biswas M, Khurshid Alam AH. *In vitro* antioxidant and free radical scavenging activity of different parts of *Tabebuia pallida* growing in Bangladesh. *BMC Res Notes* 2015;8:621.
 33. Murray PJ, Wynn TA. Protective and pathogenic functions of macrophage subsets. *Nat Rev Immunol* 2011;11:723-37.
 34. Laskin DL, Pendino KJ. Macrophages and inflammatory mediators in tissue injury. *Annu Rev Pharmacol Toxicol* 1995;35:655-77.
 35. Hemamalini K, Bhargava A. Anti-inflammatory effect of methanolic extract of leaves of various plants. *Int J Pharma Sci* 2013;4:P477-81.
 36. Regalado AI, Sánchez LM, Mancebo B. Anti-inflammatory activity of the methanolic extracts of leaves and stems from *Tabebuia hypoleuca* (C. Wright) Urb. *J Pharm Pharmacogn Res* 2015;3:109-17.
 37. Zhang L, Hasegawa I, Ohta T. Iridoid esters from *Tabebuia avellanedae* and their *in vitro* anti-inflammatory activities. *Planta Med* 2017;83:164-71.